

**DIAGNOSES OF FOUR NEW SQUIRRELS FROM SIAM.**

By C. BODEN KLOSS, F. Z. S.

During my recent visit to Siam ( October and November 1916 ) I obtained the four new races of squirrels of which brief diagnoses are given below. Detailed descriptions will be published later in a full account of my collection of mammals.

1. **Sciurus finlaysoni trotteri**, subsp. nov.

Like *S. f. folletti* of Koh Phai, but darker and practically without the buffy tinge of that race ; tail above almost clear creamy, only the extreme tip darkened.

Typical locality Koh Lan, Inner Gulf of Siam.

2. **Sciurus atrodorsalis tachin**, subsp. nov.

Like *S. a. atrodorsalis* from the latitude of Moulmein but considerably smaller and with a grizzled line down the centre of the chest.

Typical locality Tachin, west of Bangkok, Central Siam.

3. **Sciurus epomophorus inexpectatus**, subsp. nov.

Like *S. e. milleri* from Trang, Peninsular Siam, but paler ; much paler than *S. e. davisoni* from Southern Tenasserim.

Typical locality Koh Lak, Pran, S. W. Siam.

4. **Sciurus erythræus pranis.**, subsp. nov.

Like *S. e. rubeculus* from Trang, Peninsular Siam, with a grizzled median line, but the undersurface ochraceous, never more than partially tinged with chestnut, and the tip of the tail frequently albescent.

Typical locality Koh Lak, Pran, S. W. Siam.