

THE
JOURNAL
OF THE
SIAM SOCIETY,
Natural History Supplement.

Vol. VII, No. 4.

BANGKOK.

1930.

A REVISED LIST OF THE BUTTERFLIES OF SIAM, WITH
NOTES ON THEIR GEOGRAPHICAL DISTRIBUTION.

By E. J. GODFREY, B.Sc., F.E.S.

WITH A MAP.

Since the publication in 1916 of my paper entitled "The Butterflies of Siam,"* much progress has been made in our knowledge of the butterfly fauna of the country.

The following list is based mainly on my own collection which was started in 1908 but, as will be seen, a number of the species and subspecies recorded have been obtained by other collectors.

Papilio arcesilaus, *P. cocles*, *P. periander* and *P. allica* were described by Fabricius from Siam in 1787, but I am unable to state from what parts of the country these butterflies came or by whom they were obtained.

H. Druce was in the country in 1874, W. H. Doherty in 1891, and H. Fruhstorfer in 1900-01.

In more recent years, Messrs. W. A. Graham and C. J. Aagaard have collected in Bangkok and Peninsular Siam; Colonel C. H. Stockley in W. Siam; Mr. H. M. Pendlebury (Systematic Entomologist of the F. M. S. Museums) in Nakon Sritamarat, P. Siam; Mr. M. B. Tennent in the Chiengmai district, N. Siam; Major W. R. S. Ladell in C. Siam and on the Pran River, W. Siam and Mr. J. J. McBeth in Sriracha.

Dr. W. H. Beach has kindly sent me specimens from Chieng-

* Journ. Nat. Hist. Soc. Siam, ii, pp. 106-147 (1916).

rai, N. Siam, and I have also received specimens, obtained by their collectors in various parts of the country, from Dr. Malcolm Smith and Sir Walter Williamson.

At various times during the last nine months Dr. A. F. G. Kerr has very kindly collected for me on inspection tours in Peninsular Siam and, as a result of his activities, over twenty species and subspecies have been added to the list.

Mr. Tennent's collection was very kindly presented to me by Mrs. Harris of Chiengmai.

My list of 1916 included 371 species; the present one deals with 754 named forms (specific and subspecific) recorded under 692 species.

The following table shows how the total of the species is made up:—

	Species
Papilionidæ	99
Danaidæ	36
Satyridæ	81
Nymphalidæ	139
Erycinidæ	19
Lycænidæ	179
Hesperiidæ	139
<hr/>	
	692

Of the species, 8 are known only from Siam and of the subspecies, 48. Sixty-two species are represented by 2 subspecies, and four by 3.

The following species and subspecies originally described from Siam have since been found in Burma: *Hestia leuconoe siamensis* Godf.; *Stichophthalma godfreyi* Roths.; *Azanus urios*, *Arhopala dispar* and *Arhopala andamanica ignara* Ril. and Godf.

Orthophaetus barroni Ril. and Godf. has proved to be synonymous with *Capila mackwoodi* Evans, and *Everes rileyi* Godf. is now regarded as a subspecies of *Everes parhasius* Fabr.

The types of all new species and subspecies obtained by me in Siam are incorporated in the main collection of the British Museum, together with examples (δ and φ) of practically every butterfly of which I have obtained more than one specimen.

Colonel Stockley's collections (three in number) were worked out by Captain N. D. Riley and the majority of his specimens are also in the British Museum.

Mr. Pendlebury's collection* from Peninsular Siam is incorporated in the collections of the F. M. S. Museums.

The zoo-geographical divisions marked on the map, and used throughout the paper, are those proposed by Mr. C. Boden Kloss in the Journal of the Natural History Society of Siam (Vol. i, p. 250). The need for such divisions was felt owing to the fact that Siam is a country which is almost unique in the number of different minor faunal regions which meet within its political boundaries.

Two striking examples of differences in the butterfly fauna are seen in the cases of *Elymnias nesaea* and *Elymnias hypernestra* both of which are represented by three good and distinct subspecies.

In my lists of the Lycænidæ and Hesperiidæ I have followed Colonel Evans in his excellent series of papers entitled "The Identification of Indian Butterflies", and published in the Journal of the Bombay Natural History Society (Vols. XXVIII-XXXI). In dealing with the remaining Families I have followed Dr. A. Seitz in "The Macrolepidoptera of the World". Throughout the paper I have given references under each species to Dr. Seitz' great work.

As the list of Siamese butterflies is now so far advanced, I have added an index, but this does not imply that the list is to be considered complete.

I take this opportunity of acknowledging my great indebtedness to Captain Riley for the very generous aid he has accorded me at all times since 1913 in working out my material and in recording my results.

* The paper dealing with this collection, entitled "Notes on the Lepidoptera Rhopalocera (Butterflies) taken in Nakon Sritamarat, P. Siam," was published in the Journ. F. M. S. Museums, Vol. xi, pp. 21-48, April 1923.

PAPILIONIDAE.

PAPILIONINAE.

ORNITHOPTERA-GROUP.

1. *Papilio helena cerberus* Feld.*Papilio helena* Linnaeus, Syst. Nat. x, p. 461 (1758).*Papilio helena cerberus* Jordan, in Seitz, Macrolep. ix, p. 24 (1909).Khao Ram (2,000 ft.), Nakon Sritamarat, P. Siam (*Pendlebury*).*Distribution*: Sikkim to Burma; Siam; Malay Peninsula; Borneo.

Other subspecies are found in Java, Sumatra and the Celebes.

2. *Papilio aeacus* (Feld.).*Ornithoptera aeacus* Felder, Wien. Ent. Monats. iv, p. 225 (1860);Jordan (*Papilio*), in Seitz, Macrolep. ix, p. 25 (1909).*Distribution*: Sikkim to China and Formosa, south to the Malay Peninsula.(a) *Papilio aeacus praecox* Fruhst.*Papilio aeacus praecox* Fruhstorfer, Iris, p. 134 (1913).

Widely spread over Siam proper.*

Distribution: Fruhstorfer (*l. c.*) restricts this subspecies to Siam.(b) *Papilio aeacus thomsoni* Bates.*Papilio aeacus thomsoni* Jordan, in Seitz, Macrolep. ix, p. 25 (1909).2 ♀ ♀: Wat Kiriwong, Nakon Sritamarat, P. Siam, 3. 5. 28 (*Kerr*).*Distribution*: P. Siam; Malay Peninsula.

Nox-Group.

3. *Papilio sycorax* Grose-Smith.*Papilio sycorax* Grose-Smith, Ent. Month. Mag. xxi, p. 284 (1885);

Jordan, in Seitz, Macrolep. ix, p. 29, pl. 17 a, b (1909).

Khao Ram (1,000 ft.) and Khao Luang (2,000 ft.), P. Siam, (*Pendlebury*); Patalung and Nakon Sritamarat, P. Siam, (*Kerr*).*Distribution*: P. Siam; Malay Peninsula; Sumatra.4. *Papilio varuna* White.*Papilio varuna* White, Entomol. i, p. 280 (1842).

* That is, Siam excluding the Peninsula.

Distribution: Sikkim; Assam; Burma; Tonkin; Siam; Malay Peninsula; Penang.

(a) *Papilio varuna astorion* Westw.

Papilio varuna astorion Jordan, in Seitz, Macrolep. ix, p. 30 (1909).

1 ♂: Me Wong W. Siam (*Stockley*); 1 ♂: Chiengmai, N. Siam (*Tennent*).

Distribution: Sikkim to Tavoy; Tonkin; N. and W. Siam.

(b) *Papilio varuna varuna* White.

Papilio varuna varuna Jordan, in Seitz, Macrolep. ix, p. 30 (1909).

Bangnara (*Godfrey*); Ban Lan Sah and Khao Ram (*Pen-dlebury*); Wat Kiriwong, Nakon Sritamarat (*Kerr*)—all in P. Siam.

Distribution: Mergui; P. Siam; Malay Peninsula; Penang.

5. *Papilio zaleucus* Hew.

Papilio zaleucus Hewitson, Exot. Butt. iii, Pap. pl. 7, figs. 24, 25 (1865); Jordan, in Seitz, Macrolep. ix, p. 30, fig. 17 c (1909).

Me Song forest, Prae, N. Siam and Sriracha forest, S. E. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*); Me Wong, W. Siam (*Stockley*).

Distribution: Shan States to S. Burma; Siam; Tonkin.

LATREILLEI-Group.

6. *Papilio adamsoni* Gr-Sm.

Papilio adamsoni Grose-Smith, Ann. Mag. Nat. Hist. (5) xviii p. 149 (1886); Jordan, in Seitz, Macrolep. ix, p. 31, fig. 19 c (1909).

Chiengmai, N. Siam (*Tennent*); Pechaburi River, W. Siam (*Godfrey*); Pang Yao, W. Siam (*Stockley*).

Distribution: Burma; Shan States; Siam.

7. *Papilio philoxenus polyeuctes* Dbldy.

Papilio philoxenus Gray, Zool. Misc. p. 132 (1831).

Papilio philoxenus polyeuctes Jordan, in Seitz, Macrolep. ix, p. 32 (1909).

Doi Chang (4,500 ft.) and Me Song forest, Prae, N. Siam (*Godfrey*); Pang Yao, W. Siam (*Stockley*).

Distribution: Sikkim to Burma; Tonkin; Siam.

The typical form occurs from Kashmir to Nepal, and a sub-species is found in Formosa.

8. *Papilio dasarada barata* Roths.

Papilio dasarada Moore, Cat. Lep. Mus. E. I. C. p. 96 (1857).

Papilio dasarada barata Jordan, in Seitz, Macrolep ix, p 32 (1909).

1 ♂ : Doi Chang (4,500 feet), N. Siam, 13. 4. 21 (*Godfrey*) ;

2 ♂ ♂ : Me Wong (2,500 feet), W. Siam, 13. 4. 26 (*Stockley*).

Distribution : Burma to Siam.

The typical form occurs from Sikkim to Assam.

9. *Papilio laos* Ril. & Godf.

Papilio laos Riley and Godfrey, Journ. Nat. Hist. Soc. Siam, iv, p. 168, pl 4 (1921).

1 ♂ : Mouth of Menam Songkram, Sayaburi (Saniaburi), E. Siam.

Also seen on the French Laos bank of the Mekong River.

Distribution : Siam ; French Laos.

This species is nearest to *P. alcinous* Klug.

COON-Group.

10. *Papilio neptunus neptunus* Guer.

Papilio neptunus Guerin, Rev. Zool., p. 43 (1840).

Papilio neptunus neptunus Jordan, in Seitz, Macrolep. ix, p. 33 (1910).

1 ♂ , 1 ♀ : Ban Prakawp, P. Siam, 17. 7. 28 (*Kerr*).

Distribution : S. Burma ; P. Siam ; Malay Peninsula.

Subspecies are found in Sumatra, Nias and Borneo.

11. *Papilio coon doubledayi* Wall.

Papilio coon Fabricius Syst. Ent. iii (1), p. 10 (1793).

Papilio coon doubledayi Jordan, in Seitz, Macrolep. ix, p. 34, fig. 166 (1909).

Khao Ram and Ban Lan Sah Kah, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : Burma ; Siam ; Malay Peninsula.

The species also occurs in Sumatra and Java.

HECTOR-Group.

12. *Papilio aristolochiae* Fabr.

Papilio aristolochiae Fabricius, Syst. Ent. p. 443 (1775).

Distribution : China ; India ; Ceylon ; Malay Peninsula and Archipelago as far as Flores.

(a) *Papilio aristolochiae goniopeltis* Roths.

Papilio aristolochiae goniopeltis Jordan, in Seitz, Macrolep. ix, p. 36 (1909).

Widely spread over Siam proper.

Distribution: Hongkong southwards to Burma and Siam.

- (b) *Papilio aristolochiae aristolochiae* Fabr.

Papilio aristolochiae Jordan, in Seitz, Macrolep. ix, p. 38 (1909).

1 ♂, 1 ♀: Nakon Sritamarat, P. Siam (Pendlebury).

Distribution: India; P. Siam.

Pendlebury states that both the specimens he records agree with the form *diphilus* Esp.

AGESTOR-Group.

13. *Papilio agestor agestor* Gray.

Papilio agestor Gray, Zool. Misc. p 32 (1832).

Papilio agestor agestor Jordan, in Seitz, Macrolep. ix, p. 41 (1909).

Khun Than, 2,500 ft. N. Siam (Godfrey); Doi Sutep, 5,000 ft.

N. Siam (*Tennent*).

Distribution: Sikkim to N. Burma; Tonkin; Siam.

The species extends south to the Malay Peninsula.

14. *Papilio epycides epycides* Hew.

Papilio epycides Hewitson, Exot. Butt. iii, pl. 6, fig. 16 (1864).

Papilio epycides epycides Jordan, in Seitz, Macrolep. ix, p. 41, fig. 29c (1909).

2 ♂♂: Chiengmai, N. Siam, Feb. 1921 (*Tennent*).

Distribution: Sikkim to N. Burma; Siam.

Replaced from the Shan States to the Karen Hills by *P. e. hypochra* Jord.

15. *Papilio slateri marginata* Oberth.

Papilio slateri Hewitson, Exot. Butt. ii, pl. 4 (1859).

Papilio slateri race *tavoyanus* Bingham, Faun. Br. Ind. Butt. ii, p. 73.

Papilio slateri slateri, marginata et tavoyanus Jordan, in Seitz, Macrolep. ix, p. 41 (1909).

3 ♂♂, 1 ♀: Doi Sutep, Chiengmai, N. Siam, March 1921 (*Tennent*).

Distribution: Tonkin; Shan States to Karen Hills; N. Siam

The species is distributed from Sikkim to Sumatra and Borneo.

Two of the males recorded above, without white marks on the hind wing upperside, approximate to *P. slateri slateri* Hew.; the other is referable to *P. slateri tavoyanus* Butl. Riley * is of the opinion

* Riley, *in litt.*, 9. 8. 27.

that all should be called *P. slateri marginata* Oberth., which covers all intermediate zone specimens. He says:—"These two subspecies (*slateri slateri* and *slateri tavoyanus*) seem moderately constant in their own areas, but between these areas is a large tract, apparently, in which either form occurs".

CLYTIA-Group.

16. *Papilio clytia* Linn.

Papilio clytia Linnaeus, Syst. Nat. x, p. 479 (1758).

Papilio clytia f. *clytia*, f. *papone*, f. *janus*, f. *onpape*, f. *panope*, et f. *dissimilis* Jordan, in Seitz, Macrolep. ix, p. 42 (1909).

This very variable species is widely spread over Siam and occurs in the following forms: f. *clytia* Linn., f. *papone* Westw., f. *janus* Fruhst., f. *panope* Linn., f. *onpape* M. and f. *dissimilis* Linn. According to my observations, the *clytia* form is by far the commonest. I have not met with the *janus* form.

Distribution: India to Tonkin, Hainan, China, Formosa, and the Philippines, south to Malacca, Singapore and Timor.

17. *Papilio paradoxa telearchus* f. *danisepa* Butl.

Zelema paradoxa Zinken-Sommer, Nov. Act. Nat. Cur. xv., pl. 15, figs. 9, 10 (1832).

Papilio paradoxa telearchus f. *danisepa* Jordan, in Seitz, Macrolep. ix, p. 44, fig. 32b (1909).

Me Song forest, Prae, N. Siam and Pechaburi River, W. Siam (*Godfrey*); Klong Pong Kapo, W. Siam (*Stockley*).

Distribution: Assam to Burma; Siam; Tonkin.

The species ranges from N. India through the Malay Peninsula to the Greater Sunda Isles and Palawan.

CASTOR-Group.

18. *Papilio mahadeva* Moore.

Papilio mahadeva Moore, Proc. Zool. Soc. Lond. p. 840, pl. 51 (1878); Evans, Journ. Bomb. N. H. S. xxx, p. 237 (1923).

Papilio castor mahadeva Jordan, in Seitz, Macrolep. ix, p. 45 (1909).

Chiengrai, N. Siam (*Tennent*); Muak Lek, E. Siam and Pechaburi River, W. Siam (*Godfrey*); Nakon Sritamarat P. Siam (*Pendlebury*).

Distribution: Burma; Siam; Malay Peninsula.

Jordan (*l. c.*) treats *mahadeva* as a subspecies of *castor* Westw.

19. Papilio castor mehala Grose-Smith.

Papilio castor Westwood, Ann. Mag. Nat. Hist. ix, p. 37 (1842).

Papilio castor mehala Jordon, in Seitz, Macrolep. ix, p. 45 (1909).

1 ♂ : Me Song forest, Prae, N. Siam, April 1918.

This specimen is now in the British Museum and is the only example of *mehala* deposited there.

Distribution : S. Burma; Siam.

The species ranges from Sikkim to Hainan and Formosa, south to the Malay Peninsula.

DEMOLEUS-Group.

20. Papilio demoleus malayanus Wall.

Papilio demoleus Linnaeus, Syst. Nat. x, p. 464 (1758).

Papilio demoleus malayanus Jordon, in Seitz, Macrolep. ix, p. 48 (1909).

Widely spread over Siam,

Distribution : S. Burma; Siam; Annam; Malay Peninsula; Singapore.

The species ranges from India westwards to Persia, eastwards to China and Tonkin, southwards to the Lesser Sunda Isles and N. Australia. It apparently does not occur in the Greater Sunda Isles or the Philippines.

HELENUS-Group.

21. Papilio demolion demolion Cram.

Papilio demolion Cramer, Pap. Exot. i, p. 40, pl. 89, figs. A, B (1776).

Papilio demolion demolion Jordon, in Seitz, Macrolep. ix, p. 51, fig. 21a (1909).

Me Song forest, Prae, N. Siam and Sriracha forest, S. E. Siam. (*Godfrey*); Ban Prakawp, Songkla, P. Siam (*Kerr*).

Distribution : S. Burma; Siam; Malacca; Sumatra; Borneo; Java.

Subspecies are found in Nias and Palawan.

22. Papilio noblei de Nicév. (=henricus Oberth.).

Papilio noblei de Nicéville, Journ. As. Soc. Beng. p. 287 (1888);

Jordan, in Seitz, Macrolep. ix, p. 52, fig. 216 (1909).

1 ♂ : Me Song forest, Prae, N. Siam.

Distribution: This very rare species is represented in the British Museum by 1 ♂ from Upper Burma (*Capt. Broughton*, 1900) and 6 ♂♂ (in the Adams Coll.) from Tonkin.

P. henricus Oberth. must be regarded as an absolute synonym of *P. noblei*.

23. *Papilio nephelus saturnus* Guer.

Papilio nephelus Boisduval, Spec. Gen. Lep. i, p. 210 (1836).

Papilio nephelus saturnus Jordan, in Seitz, Macrolep. ix, p. 52, fig. 22a (1909).

Bangnara (*Godfrey*); Nakon Sritamarat (*Pendlebury*)—P. Siam.

Distribution: P. Siam; Malay Peninsula; Penang.

The typical form occurs in Java, and subspecies are found in Sumatra and Borneo.

24. *Papilio chaon chaon* Westw.

Papilio chaon Westwood, Arcana Ent. ii, p. 97 (1845).

Papilio chaon chaon Jordan, in Seitz, Macrolep. ix, p. 53, figs. 22b, c (1909).

Me Song forest and Chiengrai, N. Siam; Pak Jong, E. Siam; Sriracha, S. E. Siam; Pechaburi River, W. Siam (*Godfrey*); Ban Prakawp, Songkla, P. Siam (*Kerr*).

Distribution: Nepal to N. Burma; Tonkin; Siam; Cochin China.

A subspecies is found in Hainan, China and Formosa.

25. *Papilio helenus helenus* Linn.

Papilio helenus Linnaeus, Syst. Nat. x, p. 459 (1758).

Papilio helenus helenus Jordan, in Seitz, Macrolep. ix, p. 54, fig. 216 (1909).

Prae, N. Siam; Muak Lek, E. Siam; Sriracha, S. E. Siam; (*Godfrey*); Chiengrai, N. Siam (*Tennent*); Nakon Sritamarat, P. Siam (*Kerr*).

Distribution: N. India to S. China, Tonkin and Hainan, south to Burma, Siam and the Malay Peninsula.

The species extends through the Malay Archipelago to Timor.

26. *Papilio iswara iswara* White.

Papilio iswara White, Entomolog. i, p. 280 (1842).

Papilio iswara iswara Jordan, in Seitz, Macrolep. ix, p. 55, fig. 21c (1910).

Bangnara (*Godfrey*) ; Nakon Sritamarat (*Pendlebury*)—P. Siam.

Distribution : Mergui ; P. Siam ; Malay Peninsula ; Penang ; Sumatra.

A subspecies is found in Borneo and the Natunas.

27. **Papilio hippoönous** Feld.,

Papilio hippoönous Felder, Wien. Ent. Mon. vi, p. 283 (1862).

Distribution ; S. Burma ; Siam ; Luzon ; Palawan ; Mindano ; Bazilan ; Sangir and Talaut Islands.

(a) **Papilio hippoönous pitmani** Elw.

Papilio hippoönous pitmani Jordan, in Seitz, Macrolep. ix, p. 59 (1909).

Pak Jong, E. Siam.

Distribution : S. Burma ; E. Siam.

(b) **Papilio hippoönous siamensis** Godf.

Papilio hippoönous siamensis Godfrey, Journ. Nat. Hist. Soc. Siam, ii, p. 110 (1916).

Petchaburi River, W. Siam.

Distribution : Known only from W. Siam.

POLYTES-Group.

28. **Papilio polytes polytes** Linn

Papilio polytes Linnaeus, Syst. Nat. x, p. 460 (1758).

Papilio polytes polytes Jordan, in Seitz, Macrolep. ix, p. 60 (1909).

Widely spread over Siam.

Distribution : The species ranges from India to China and the Loo Choo Islands, south to the Moluccas and Timor.

MEMNON-Group.

29. **Papilio memnon agenor** Linn.

Papilio memnon Linnaeus, Syst. Nat. x, p. 460 (1758).

Papilio memnon agenor Jordan, in Seitz, Macrolep. ix, p. 72, fig. 30a (1909).

Widely spread over Siam.

Distribution : Sikkim to Siam.

The species ranges from India to China and Japan, south to Malaya as far as Borneo and Flores.

Three forms of the female occur in Siam:—(a) ♀-f. *agenor* Linn., (b) ♀-f. *distantius* Roths., and (c) ♀-f. *butlerianus* Roths. The first two forms are fairly common in Bangkok, the third I know of only from a single specimen obtained by Aagaard at Bangnara, P. Siam.

30. *Papilio protenor euprotenor* Fruhst.

Papilio protenor Cramer, Pap. Exot. i, p. 77 (1775).

Papilio protenor euprotenor Jordan, in Seitz, Macrolep. ix, p. 75 (1909).

Me Song forest and Chiengmai, N. Siam; Pechaburi River, W. Siam.

Distribution: Sikkim to Burma; Tonkin; Hainan; Siam.

The typical form occurs from Kashmir to Kumaon.

31. *Papilio rhetenor rhetenor* Westw.

Papilio rhetenor Westwood Arc. Ent. i, p. 59 (1842).

Papilio rhetenor rhetenor Jordan, in Seitz, Macrolep. ix, p. 76, fig. 29b (1909).

1 ♂ : Chiengmai, N. Siam, April 1921 (*Tennent*); 1 ♀ : Chiengrai, N. Siam, May 1921 (*Godfrey*); 1 ♂ : Pan Yao, W. Siam, Feb. 1920 (*Stockley*).

Distribution: Nepal to S. Burma; Siam.

A subspecies is found in West and Central China.

PARIS-Group.

32. *Papilio elephenor elephenor* Doubl.

Papilio elephenor Doubleday, Ann. Mag. N. H. xvi, p. 305 (1845); Jordan, in Seitz, Macrolep. ix, p. 77, fig. 34 a, (1909).

Papilio elephenor elephenor Evans, Journ. Bomb. N. H. S. xxix, p. 235 (1923).

1 ♂, 1 ♀ : Doi Sutep (4,500ft.), N. Siam, March 1921 (*Tennent*).

Distribution: Assam; N. Siam.

A subspecies is found in the Shan States.

33. *Papilio polyctor triumphator* Fruhst.

Papilio polyctor Boisduval, Spec. Gen. Lep. i, p. 205 (1836).

Papilio polyctor triumphator Jordan, in Seitz, Macrolep. ix, p. 79 (1909).

Papilio polyctor ganesa Evans, Journ. B. N. H. S. xxix, p. 235 (1923).

Chantabun, S. E. Siam (*Fruhstorfer*).

Distribution: Jordan (*l. c.*) restricts this subspecies to Siam and Tonkin and gives the range of the species as Kashmir to Tonkin and Siam.

Evans (*l. c.*) treats *triumphator* as a synonym of *ganesa* Moore (Sikkim-N. Burma).

34. Papilio paris paris Linn.

Papilio paris Linnaeus, Syst. Nat. x, p. 459 (1758).

Papilio paris paris Jordan, in Seitz, Macr. ix, p. 79, fig. 34 e (1909).

Prae, N. Siam; Pak Song and Muak Lek, E. Siam; Pechaburi River, W. Siam (*Godfrey*); Hue Sak, Chumphorn, P. Siam (*Aagaard*).

Distribution: Hongkong to Malacca and N. W. India.

Subspecies are found in S. India, China, Formosa, Sumatra, and Java.

35. Papilio arcturus arcturus Westw.

Papilio arcturus Westwood, Ann. Mag. Nat. Hist. ix, p. 37 (1842).

Papilio arcturus arcturus Jordan, in Seitz, Macrolep. ix, p. 80 (1909).

1 ♂: Hot Spring, Me Lamoung (2,000 ft.), W. Siam, 15. 4. 20 (Stockley); 1 ♂: Doi Chang (4,500 ft.), N. Siam, April 1921 (*Godfrey*).

Distribution: Sikkim to Burma; Siam.

A subspecies is found in West and Central China.

PALINURUS-Group.

36. Papilio palinurus palinurus Fabr.

Papilio palinurus Fabricius, Mant. Ins. ii, p. 2 (1787).

Papilio palinurus palinurus Jordan, in Seitz, Macrolep. ix, p. 81 (1909).

Pechaburi River, W. Siam (*Godfrey*); Bangnara, P. Siam (*Aagaard*).

Distribution: S. Burma; Siam; Malacca; Borneo; Sumatra. Subspecies are found in Nias, Palawan, and the Philippines.

ANTIPHATES-Group.

37. Papilio agetes agetes Westw.

Papilio agetes Westwood, Arc. Ent. ii, p. 23, pl. 55, figs. 1, 2 (1843).

Papilio agetes agetes Jordan, in Seitz, Macrolep. ix, p. 87 (1910).

Me Song forest, Prae, N. Siam; Sriracha forest, S. E. Siam; Pechaburi River, W. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*).

Distribution: Burma; Tonkin; Hainan; Siam.

The species extends to the Malay Peninsula, Sumatra and Borneo.

38. *Papilio nomius swinhoei* Moore.

Papilio nomius Esper, Ausl. Schmett. p. 210, pl. 52 (1785-98).

Papilio nomius swinhoei Jordan, in Seitz, Macrolep. ix, p. 87, fig. 41 c (1909).

Widely spread over Siam.

Distribution: Assam to Burma; Tonkin, Hainan; Siam.

The typical form occurs from Sikkim to S. India and Ceylon.

39. *Papilio aristaeus hermocrates* Feld.

Papilio aristaeus Cramer, Pap. Exot. iv, p. 60, pl. 318, figs. E. F. (1782).

Papilio aristaeus hermocrates Jordan, in Seitz, Macrolep. ix, p. 88, fig. 41 a (1909).

Me Song forest, Prae, N. Siam; Muak Lek, E. Siam; Sriracha, S. E. Siam; Pechaburi River, W. Siam.

Distribution: Burma to the Malay Peninsula; Borneo; Sumatra.

The species ranges from N. India to the Bismarck Isles and N. Australia.

40. *Papilio antiphates pompilius* Fabr.

Papilio antiphates Cramer, Pap. Exot. i, p. 113, pl. 72 (1775).

Papilio antiphates pompilius Jordan, in Seitz, Macrolep. ix, p. 89 (1909).

Widely spread over Siam.

Distribution: Sikkim to Burma; Hainan; Siam; Annam.

The species ranges from India and China to Malaya.

PAYENI-Group.

41. *Papilio gyas gyas* Westw.

Papilio gyas Westwood, Arcana Ent. i, p. 41 (1841).

Papilio gyas gyas Jordan, in Seitz, Macr. ix, p. 91, fig. 39 c (1909).

1 ♂ : Chieng Dao (2,100 m.), N. Siam, Jan. 1920.

Distribution : Sikkim to Assam ; Siam.

A subspecies is found in N. Burma.

42. **Papilio payeni langsonensis** Fruhst.

Papilio payeni Boisduval, Spec. Gen. Lep. i, p. 235 (1836).

Papilio payeni langsonensis Jordan, in Seitz, Macrolep. ix, p. 91 (1909).

1 ♂ : Me Song forest, Prae, N. Siam.

Distribution : Tonkin ; Siam.

The typical form is found in Java and subspecies occur from Sikkim to Assam, and from N. Burma to the Karen.

EURYPYLVUS-Group.

43. **Papilio cloanthus cloanthus** Westw.

Papilio cloanthus Westwood, Arc. Ent. i, p. 42 (1841).

Papilio cloanthus cloanthus Jordan, in Seitz, Macrolep. ix, p. 94 (1909).

3 ♂♂ : Phar Mieng, Chiengrai, N. Siam, 17. 4. 21 (Godfrey);

2 ♂♂ : Hot spring, Me Lamoung (2,000 ft.), W. Siam, 15. 4. 20 (Stockley).

Distribution : Kashmir to Burma ; Siam.

Subspecies are found in China and Sumatra.

44. **Papilio sarpedon sarpedon** Linn.

Papilio sarpedon Linnaeus, Syst. Nat. x, p. 46 (1758).

Papilio sarpedon sarpedon Jordan, in Seitz, Macrolep. ix, p. 95, fig. 44 d (1909).

Widely spread over Siam.

Distribution : Kashmir to Tonkin, Hainan and the Philippines, south to the Malay Archipelago as far as Lombok.

The species ranges from China and Japan to the Solomon Islands.

45. **Papilio doson axion** Feld.

Papilio doson Felder, Verh. Zool-bot. Ges. Wien. xiv, p. 305 (1864).

Papilio doson axion Jordan, in Seitz, Macrolep. ix, p. 97, fig. 43c (1909).

Widely spread over Siam.

Distribution : Kumaon to Burma ; Tonkin ; Hainan ; Siam.

The species ranges from India and China to Malaya.

46. Papilio eurypylus cheronus Fruhst.

Papilio eurypylus Linnaeus, Syst. Nat. x, p. 464 (1758).

Papilio eurypylus cheronus Jordan, in Seitz, Macrolep. ix, p. 98 (1909).

Prae, N. Siam ; Muak Lek, E. Siam ; Sriracha, S. E. Siam.

Distribution : Sikkim to Burma ; Tonkin ; Hainan ; Siam.

The species ranges from India and China to Malaya and Australia.

47. Papilio bathycles Zinken-Sommer.

Papilio bathycles Zinken-Sommer, Nov. Act. Ac. Cur. p. 157, pl. 14, figs. 6, 7 (1831).

Distribution : India to the Greater Sunda Isles and Palawan.

(a) Papilio bathycles chiron Wall.

Papilio bathycles chiron Jordan, in Seitz, Macrolep. ix, p. 100 (1909).

Prae and Chiengrai, N. Siam (*Godfrey*) ; Chiengmai, N. Siam (*Tennent*) ; Me Lamoung, W. Siam (*Stockley*).

Distribution : Sikkim to Burma ; Siam ; Annam.

(b) Papilio bathycles bathycloides Honrath.

Papilio bathycles bathycloides Jordan, in Seitz, Macrolep. ix, p. 100 fig. 44 c (1909).

1 ♂ : Kao Den, Patalung, P. Siam (*Kerr*).

Distribution : P. Siam ; Malay Peninsula ; Sumatra ; Borneo and Palawan.

48. Papilio agamemnon agamemnon Linn.

Papilio agamemnon Linnaeus, Syst. Nat. x, p. 462 (1758).

Papilio agamemnon agamemnon Jordan, in Seitz, Macrolep. ix, p. 101, fig. 45 d (1909).

Widely spread over Siam.

Distribution : From India and China to Bali and the Philippines.

Subspecies are found in the Papuan region down to Queensland and the Solomon Islands.

49. Papilio arycles arycleoides Fruhst.

Papilio arycles Boisduval, Spec. Gen. Lep. i, p. 231 (1836).

Papilio arycles arycleoides Fruhstorfer, Iris, p. 344 (1901); Jordan, in Seitz, Macrolep. ix, p. 102 (1909).

Muak Lek, E. Siam ; Sriracha, S. E. Siam.

Distribution : Confined to Siam.

The species ranges from Assam to the Greater Sunda Islands and Palawan.

MACAREUS-Group.

50. **Papilio macareus indochinensis** Fruhst.

Papilio macareus Godart, Ency. Méth. ix, p. 76 (1819).

Papilio macareus indochinensis Jordan, in Seitz, Macrolep. ix, p. 104 (1909).

Me Song forest, Prae, N. Siam ; Pak Jong, E. Siam ; Pechaburi River, W. Siam (*Godfrey*) ; Chiengmai, N. Siam (*Tennent*).

Distribution : Tonkin ; Siam ; Annam.

The species ranges from Sikkim to Hainan and the Philippines.

51. **Papilio xenocles** Doubl.

Papilio xenocles Doubleday, in Gray's Zool. Misc., p. 74 (1842).

Distribution : Sikkim to Burma ; Tonkin ; Hainan ; Siam.

(a) **Papilio xenocles kephisos** Fruhst.

Papilio xenocles kephisos Jordan, in Seitz, Macrolep. ix, p. 105 (1909).

Pechaburi River (*Godfrey*) ; Hot spring, Me Lamoung (*Stockley*)—W. Siam.

Distribution : Burma ; Tonkin ; W. Siam.

(b) **Papilio xenocles lindos** Fruhst.

Papilio xenocles lindos Fruhstorfer, Soc. Entomol., p. 142 (1901).

Muak Lek and Pak Jong, E. Siam.

Distribution : Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

52. **Papilio leucothoe leucothoe** Westw.

Papilio leucothoe Westwood, Arc. Ent. ii, p. 128 (1845).

Papilio leucothoe leucothoe Jordan, in Seitz, Macrolep. ix, p. 105, fig. 47 b (1909).

2 ♂ ♂ : Klawng Ton, Satul, P. Siam, 10.3.28 (*Kerr*).

Distribution : P. Siam ; Malay Peninsula ; N. E. Sumatra.

Subspecies are found in W. Sumatra and Borneo.

53. **Papilio delesserti delesserti** Guér.*Papilio delesserti* Guérin, Rev. Zool. p. 233 (1839).*Papilio delesserti delesserti* Jordan, in Seitz, Macrolep. ix, p. 105, fig. 47 c (1910).1 ♂ : Khao Luang (2,000 ft.), Nakon Sritamarat, 25.3.22 (*Pendlebury*) ; 1 ♂ : Ban Ton, Satul, 14.3.28 (*Kerr*)—P. Siam.*Distribution* : P. Siam ; Malay Peninsula ; Penang ; Sumatra ; Java ; Borneo.

Subspecies are found in Nias and Palawan.

54. **Papilio megarus similis** Lathy.*Papilio megarus* Westwood, Arc. Ent. ii, p. 98, pl. 72, fig. 2 (1845).*Papilio megarus similis* Jordan, in Seitz, Macrolep. ix, p. 106 (1910).

Widely spread over Siam.

Distribution : S. Burma ; Tonkin ; Hainan ; Siam ; Malay Peninsula.

The typical form occurs from Assam to N. Burma.

GENUS *Leptocircus*.55. **Leptocircus curius curius** (Fabr.).*Papilio curius* Fabricius, Mant. Ins. ii, p. 9 (1787).*Leptocircus curius* Jordan, in Seitz, Macrolep. ix, p. 108 (1910).Prae, N. Siam ; Muak Lek, E. Siam ; Sriracha S. E. Siam (*Godfrey*) ; Kambengpet and Me Wong, W. Siam (*Stockley*).*Distribution* : Assam to the Greater Sunda Isles and Palawan.

Subspecies are found in China and Nias.

56. **Leptocircus meges viriscens** Butl.*Papilio meges* Zinken-Sommer, Nov. Act. Ac. Nat. Cur., p. 161 (1832).*Leptocircus meges viriscens* Jordan, in Seitz, Macrolep. ix, p. 108 (1910).Prae, N. Siam ; Pak Jong and Muak Lek, E. Siam ; Sriracha, S. E. Siam (*Godfrey*). Nakon Sritamarat, P. Siam (*Pendlebury* ; *Kerr*).*Distribution* : Burma ; Tonkin ; Hainan ; Siam ; Malay Pen.

The species ranges from India and China to Malaya and the Philippines.

PIERINAE.

57. **Leptosia xiphia xiphia** (Fabr.).*Papilio xiphia* Fabricius, Spec. Ins. ii, p. 43 (1781).

Leptosia xiphia xiphia Fruhstorfer, in Seitz, Macrolep. ix, p. 123 (1910).

Widely spread over Siam.

Distribution : Ceylon ; India ; Burma to Annam.

The species ranges from N. India to China and the Malayan subregion.

58. *Delias agostina* (Hew.).

Pieris agostina Hewitson, Exot. Butt. i, pl. 1, figs. 1, 2 (1862).

Delias agostina, infumata et annamitica Fruhstorfer, in Seitz, Macrolep. ix, p. 123 (1910).

1 ♂ : Khun Than (3,000 ft.), N. Siam, 1 ♂ : Me Song forest, Prae, N. Siam (Godfrey) ; 1 ♀ : Pan Yao, W. Siam, (Stockley).

Fruhstorfer (*l. c.*) splits this species into three races : (a) *D. a. agostina* Hew. from Sikkim, (b) *D. a. infumata* from Assam and Tenasserim, and (c) *D. a. annamitica* Fruhst. from S. Annam. From the material in the British Museum it would appear that these races are not really separable, and that all are better regarded as true *agostina*.

59. *Delias agoranis* H. G. Smith.

Delias agoranis H. G. Smith, Ann. Mag. N. H. (5) xx, p. 226 (1887) ; Moore (*Piccarda*), Lep. Ind., p. 182, pl. 535, figs. 2, 2a (1905) ; Riley and Godfrey, Journ. Nat. Hist. Soc. Siam, iv, p. 169 (1921).

Delias singhapura agoranis, Fruhstorfer, in Seitz, Macrolep. ix, p. 124 (1910).

Delias agostina agoranis Evans, Journ. Bomb. N. H. S. xxix, p. 250 (1923).

Me Song forest, Prae, N. Siam.

Distribution : Burma ; Siam.

Riley and Godfrey (*l. c.*) regard *agoranis* as a good species, equally distinct from both *singhapura* and *agostina*.

60. *Delias hyparete* (Linn.).

Papilio hyparete Linnaeus, Syst. Nat. (10) i, p. 469 (1758).

Distribution : N. India to Formosa and the Philippines, south to the Malay Peninsula and Archipelago.

(a) ***Delias hyparete ciris* Fruhst.**

Delias hyparete ciris Fruhstorfer, in Seitz, Macrolep. ix, p. 125, pl. 51 b (1910).

Widely spread over Siam proper.

Distribution: Tonkin; Cochin-China; Siam.

(b) ***Delias hyparete metarete* Butl.**

Delias hyparete metarete Fruhstorfer, op. cit., p. 125.

Nakon Sritamarat, P. Siam (*Pendlebury*; *Kerr*).

Distribution: P. Siam; Malay Peninsula; Singapore.

61. ***Delias belladonna ithiela* (Butl.).**

Papilio belladonna Fabricius, Ent. Syst. iii, p. 180 (1793).

Thyca ithiela Butler, Ann. Mag. N. H. (4) iv, p. 242 (1869).

Delias belladonna ithiela Fruhstorfer, in Seitz, Macrolep. ix, p. 130 (1911).

Me Wong, W. Siam (*Stockley*); Doi Sutep, N. Siam (*Tennent*).

Distribution: Sikkim; Siam.

The species ranges from N. India to Sumatra.

62. ***Delias descombesi leucacantha* Fruhst.**

Pieris descombesi Boisduval, Spec. Gén. Lep. i, p. 465 (1836).

Delias descombesi leucacantha Fruhstorfer, in Seitz, Macrolep. ix, p. 131, fig. 56 c (1910).

Me Song forest, Prae, N. Siam; Muak Lek, E. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam.

Distribution: Sikkim to Siam.

The typical form occurs in Cochin-China and a subspecies is found in the Malay Peninsula.

63. ***Delias aglaia* (Linn.).**

Papilio aglaia Linnaeus, Syst. Nat. i, p. 465 (1758).

Distribution: Formosa; China; India; Malay Peninsula; Sumatra; Nias; Borneo; Palawan.

(a) ***Delias aglaia thyra* Fruhst.**

Delias aglaia thyra Fruhstorfer, in Seitz, Macrolep. ix, p. 133, fig. 56 c (1911).

Me Song forest, Prae, N. Siam and Sriracha forest, S. E. Siam (*Godfrey*); Samson, Bangkok, C. Siam (*Aagaard*).

Distribution: Siam; Annam.

(b) **Delias aglaia parthenope** (Wall.).

Thyca parthenope Wallace, Trans. Ent. Soc. Lond., p. 347 (1867).

Delias aglaia parthenope Fruhstorfer, op. cit., p. 133.

1 ♀ : Bangnara, P. Siam, September 1913.

Distribution : P. Siam ; Malay Peninsula ; Singapore.

64. Delias ninus (Wall.).

Thyca ninus Wallace, Trans. Ent. Soc. Lond., p. 347 (1867).

Delias ninus Fruhstorfer, in Seitz, Macrolep. ix, p. 134 (1910).

Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula ; N. Borneo.

65. Delias thysbe pyramus (Wall.).

Papilio thysbe Cramer, Pap. Exot. iii, p. 70 (1782).

Thyca pyramus Wallace, Trans. Ent. Soc. Lond., p. 347 (1867).

Delias thysbe pyramus Fruhstorfer, in Seitz, Macrolep. ix, p. 134 (1910).

Me Song forest, Prae and Khun Than (2,500 ft), N. Siam (*Godfrey*) ; Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : Sikkim to Burma ; Siam ; Malay Peninsula.

The typical form occurs in China.

66. Prioneris thestylis jugurtha Fruhst.

Pieris thestylis Doubleday, in Gray's Zool. Misc., p. 76 (1842).

Prioneris thestylis jugurtha Fruhstorfer, in Seitz, Macrolep. ix, p. 137 (1910).

Chiengrai, N. Siam and Pechaburi River, W. Siam (*Godfrey*) ;

Me Wong, W. Siam (*Stockley*).

Distribution : Siam ; Annam ; Tonkin.

The typical form occurs from Mussoorie to Burma, and subspecies are found in the Malay Peninsula and Formosa.

67. Prioneris clemantine helferi Fruhst.

Pieris clemantine Doubleday, Ann. Mag. N. H. xvii, p. 23, (1846).

Prioneris clemantine helferi Fruhstorfer, in Seitz, Macrolep. ix, (1910).

Pak Jong, S. E. Siam (*Godfrey*) ; Pan Yao, W. Siam (*Stockley*).

Distribution : Tonkin ; Shan States ; Siam.

The typical form occurs from Sikkim to Burma, and subspecies are found in Hainan, the Malay Peninsula and Sumatra.

63. **Pieris canidia** (Sparr.).

Papilio canidia Sparrman, Amoen. Acad. vii, p. 504 (1768); Fruhstorfer (*Pieris*), in Seitz, Macrolep. ix, p. 139 (1910).

Me Song forest, Prae, and Chiengrai, N. Siam.

Distribution: Chitral to N. Burma; Hainan; Formosa; N. Siam.

69. **Huphina nerissa dapha** (Moore).

Papilio nerissa Fabricius, Syst. Ent., p. 471 (1775).

Appias dapha Moore, Proc. Zool. Soc., p. 838 (1878).

Huphina nerissa dapha Fruhstorfer, in Seitz, Macrolep. ix, p. 141 fig. 64 a (1919).

Widely spread over Siam.

Distribution: Burma; Siam.

The typical form occurs in China and Tonkin, and subspecies are found in Sikkim (and Assam), Formosa, Sumatra and Java.

70. **Huphina nadina nadina** (Luc.).

Pieris nadina Lucas, Rev. et Mag. Zool. (2), iv, p. 333 (1852).

Huphina nadina nadina Fruhstorfer, in Seitz, Macrolep. ix, p. 142, (1910).

Widely spread over Siam.

Distribution: Sikkim to Burma; Tonkin; Hainan; Siam.

The species ranges from Formosa to the Malay Peninsula and Sumatra.

71. **Huphina lea** (Doubt.).

Pieris lea Doubleday, Ann. Mag. N. H. xvii, p. 23 (1846).

Distribution: S. Burma to the Malay Peninsula, Sumatra the Natunas and Borneo.

(a) **Huphina lea siamensis** (Butl.).

Huphina lea siamensis Fruhstorfer, in Seitz, Macrolep. ix, p. 145 (1910).

Prae, N. Siam; Pak Jong, E. Siam; Sriracha, S. E. Siam.

Distribution: This subspecies is confined to Siam.

(b) **Huphina lea malaya** Fruhst.

Huphina lea malaya Fruhstorfer, op. cit., p. 145.

Bangnara (*Godfrey*); Nakon Sritamarat (*Pendlebury*)—P. Siam.

Distribution : P. Siam ; Malay Peninsula.

72. **Appias libythea** (Fabr.).

Papilio libythea Fabricius, Syst. Ent., p. 471 (1775).

Appias libythea race *zelmira* Bingham, Faun. Br. Ind., ii, p. 200.

Appias libythea libythea, *zelmira* et *peducaeae* Fruhstorfer, in Seitz, Macrolep. ix, p. 148 (1910).

This species is widely spread over Siam and occurs in two forms *libythea* Fabr. and *zelmira* Cram., both of which may be obtained in Bangkok, the former being quite common. Bingham treated *libythea* and *zelmira* as separate races which met in India in the plains of Bengal. Fruhstorfer (*l. c.*) divides up the species into three races : *libythea* in India and Ceylon ; *zelmira* "in Tenasserim, Annam, Indo-China, also in Assam and Sikkim and even sometimes in the plains of Bengal" ; and *peducaeae* Fruhst. in the Philippines.

If a subspecific name is to be given to examples from Siam, it must be *libythea* as this appears to be the predominant form but, considering its variability, it would seem better not to split the species at all.

73. **Appias lyncida** (Cram.).

Papilio lyncida Cramer, Pap. Exot. ii, pl. 131, fig. 13 (1779).

Distribution : India and Ceylon eastwards to Formosa and the Philippines, southwards to Celebes, Flores and Java.

(a) **Appias lyncida hippoides** Moore.

Appias hippoides Moore, Trans. Ent. Soc., p. 312 (1881).

Appias lyncida hippoides Fruhstorfer, in Seitz, Macrolep. ix, p. 148 (1910).

Widely spread over Siam proper.

Distribution : Sikkim to Burma ; Siam ; Annam.

(b) **Appias lyncida vasava** Fruhst.

Appias lyncida vasava Fruhstorfer, op. cit., p. 149 (1910).

Nakon Sritamarat, P. Siam (*Pendlebury* ; *Kerr*).

Distribution : P. Siam ; Malay Peninsula.

74. **Appias nero galba** (Wall.).

Papilio nero Fabricius, Ent. Syst. iii, p. 153 (1793).

Tachyris galba Wallace, Trans. Ent. Soc. Lond. (3) iv, p. 378 (1867).

Appias nero galba Fruhstorfer, in Seitz, Macrolep. ix, p. 150 (1910).

Prae, N. Siam ; Muak Lek, E. Siam ; Sriracha, S. E. Siam (*Godfrey*) ; Nakon Sritamarat, P. Siam (*Pendlebury* ; *Kerr*).

Distribution : Sikkim to Siam ; Tonkin.

The species ranges from Sikkim to the Malay Peninsula and Archipelago.

75. *Appias indra thronion* Fruhst.

Pieris indra Moore, Cat. Lep. Mus. E. I. C., p. 74 (1857).

Appias indra thronion Fruhstorfer, in Seitz, Macrolep. ix, p. 153 (1910).

Muak Lek, E. Siam (*Godfrey*) ; Me Lamoung, W. Siam (*Stockley*).

Distribution : Confined to Siam.

The species ranges from Sikkim southwards to the Malay Peninsula, eastwards to China.

76. *Appias lalage* (Doubl.).

Papilio lalage Doubleday, in Gray's Zool. Misc., p. 76 (1842).

Distribution : From the Himalayas southwards to Siam and the Malay Peninsula, eastwards to China.

(a) *Appias lalage argyridina* (Butl.).

Hyposcritia argyridina Butler, A. M. N. H. (5) xvi, p. 340 (1885).

Appias lalage argyridina Fruhstorfer, in Seitz, Macrolep. ix, p. 153 (1910).

Prae and Chiengrai, N. Siam (*Godfrey*) ; Me Lamoung, W. Siam (*Stockley*).

Distribution : N. Burma ; Shan States ; Siam.

(b) *Appias lalage lagela* (Moore).

Catophaea lagela Moore, Proc. Zool. Soc., p. 838 (1878).

Appias lalage lagela Fruhstorfer, op. cit., p. 153.

Chiengrai, N. Siam (*Godfrey*) ; Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : S. Burma to Mergui, P. Siam and the Malay Peninsula.

77. *Appias lalassis* (Grose-Smith).

Lade lalassis Grose-Smith, Ann. Mag. N. H. (5) xx, p. 265 (1887).

Appias lalassis Fruhstorfer, in Seitz, Macrolep. ix, p. 154 (1910).

1 ♂ : Me Song forest, Prae, N. Siam, April 1918 ; 2 ♂ ♂ : Phar Mieng, near Chiengrai, N. Siam, April 1921.

Distribution : Hitherto recorded only from Burma.

This interesting species was placed by de Nicéville under the genus *Lade* on account of the absence in the males of the anal tufts of hair which are so conspicuous in the males of *Appias*.

78. *Appias albina darada* (Feld.).

Pieris albina Boisduval, Sp. Gén. Lep. i, p. 480 (1836).

Pieris darada Felder, Novara Reise, Lep. ii, p. 165 (1865).

Appias albina confusa (= *darada* auct. nec Feld.) Fruhstorfer, in Seitz, Macrolep. ix, p. 154 (1910).

Widely spread over Siam, sometimes occurring in enormous numbers.

Distribution : S. India ; Sikkim to Siam ; Tonkin.

The species ranges from India to Malaya, the Philippines and Moluccas.

Three ♀ forms are found in Siam : (a) ♀-f *darada* Feld. (d.s.f.), (b) ♀-f *principalis* Fruhst. (interm. form), and (c) ♀-f *semiflava* Fruhst. (w. s. f.).

Fruhstorfer's assumption that the name *darada* has been wrongly used is incorrect, and his *confusa* therefore falls.

79. *Appias melania* (Fabr.).

Papilio melania Fabricius, Syst. Ent. i, p. 475 (1755).

Distribution : India to the Philippines and Australia.

(a) *Appias melania pseudoleis* Fruhst.

Appias melania pseudoleis Fruhstorfer, in Seitz, Macrolep. ix, p. 155 (1910).

Sriracha forest, S. E. Siam.

Distribution : Confined to Siam.

(b) *Appias melania distanti* Moore.

Appias melania distanti Fruhstorfer, op. cit., p. 155.

Padang Besar, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula.

80. *Ixias pyrene verna* Druce.

Papilio pyrene Linnaeus, Mus. Ulr., p. 241 (1764).

Ixias pyrene verna Fruhstorfer, in Seitz, Macrolep. ix, p. 159 (1910).

Widely spread over Siam.

Distribution : Confined to Siam.

The species ranges from India to China, south to the Malay Peninsula.

81. *Dercas verhuelli doubledayi* Moore.

Colias verhuelli Vander Hoeven, Tijdsch. voor Nat. Ges. V, p. 341 (1839).

Dercas verhuelli doubledayi Fruhstorfer, in Seitz, Macrolep. ix, p. 160 (1910).

2 ♂♂ : Me Song forest, Prae, N. Siam, April 1918.

Distribution : Sikkim to N. Burma ; N. Siam.

The typical form occurs in China and a subspecies is found in S. Burma.

82. *Catopsilia pyranthe* (Linn.).

Papilio pyranthe Linnaeus, Syst. Nat. x, p. 469 (1758).

Catopsilia pyranthe Fruhstorfer, in Seitz, Macrolep. ix, p. 162 (1910).

Widely spread over Siam.

Distribution : India to China and Formosa, south to Malaya and Australia.

83. *Catopsilia florella* (Fabr.).

Papilio florella Fabricius, Syst. Ent., p. 479 (1775) ; Bingham (*Catopsilia*), Faun. Br. Ind. Butt. ii, p. 224 ; Bell (*Catopsilia*), Journ. Bomb. N. H. S. xxii, p. 523 ; Fruhstorfer, in Seitz, Macrolep. ix, p. 162, fig. 69 d (1910).

Although both Bingham and Bell record this species from Siam, I have neither taken it myself nor seen any specimen obtained in the country.

Distribution : India and China over Persia to Aden.

84. *Catopsilia crocale* (Cram.).

Papilio crocale Cramer, Pap. Exot. i, p. 87 (1775) ; Fruhstorfer (*Catopsilia*), in Seitz, Macrolep. ix, p. 162 (1910).

Common everywhere in Siam.

Distribution : From India through the Malayan subregion to Australia.

85. **Catopsilia pomona** (Fabr.).

Papilio pomona Fabricius, Syst. Ent., p. 479 (1775); Fruhstorfer (*Catopsilia*), in Seitz, Macrolep. ix, p. 163 (1910).

Widely spread over Siam.

The female form *catilla* Cram. is not uncommon.

Distribution : The same as that of *C. crocale*.

86. **Catopsilia scylla sidra** Fruhst.

Papilio scylla Linnaeus, Cent. Ins., p. 20 (1763).

Catopsilia scylla sidra Fruhstorfer, in Seitz, Macrolep. ix, p. 163 (1910).

Widely spread over Siam.

Distribution : Siam ; Malay Peninsula ; Sumatra.

The species ranges from S. Burma through the Malayan subregion to Australia.

87. **Terias libythea libythea** (Fabr.).

Papilio libythea Fabricius, Ent. Syst. Suppl., p. 427 (1798).

Terias libythea libythea Fruhstorfer, in Seitz, Macrolep. ix, p. 160 (1910).

2 ♂♂ : Chiengrai, N. Siam, 29.4.21.

Distribution : India ; Burma ; Siam.

Subspecies are found in the Malay Peninsula, Java, Sumatra and Australia.

88. **Terias venata venata** Moore.

Terias venata Moore, Cat. Lep. Mus. E. I. C. (1), p. 65 (1857).

Terias venata venata Fruhstorfer, in Seitz, Macrolep. ix, p. 166 (1910).

Bangkok, C. Siam.

Distribution : India ; Ceylon ; Siam.

A subspecies occurs from Sikkim to N. Burma.

89. **Terias laeta pseudolaeta** Moore.

Terias laeta Boisduval, Spec. Gén. Lep. i, p. 614 (1836).

Terias laeta pseudolaeta Fruhstorfer, in Seitz, Macrolep. ix, p. 166 (1910).

Chiengrai, N. Siam ; Pak Jong, E. Siam ; Bangkok, C. Siam.

Distribution : S. Burma ; Siam.

The species ranges from the Himalayas to Annam, Tonkin and Hainan.

90. **Terias hecabe hecabe** (Linn.).

Papilio hecabe Linnaeus, Syst. Nat. x, p. 470 (1758).

Terias hecabe hecabe Fruhstorfer, in Seitz, Macrolep. ix, p. 167, fig. 73 f (1910).

Widely spread over Siam.

Distribution: Fruhstorfer (*l. c.*) gives the range of the typical form as "Southern China to Anterior India."

The species is distributed from Corea and Japan south to Australia.

91. **Terias blanda davidsoni** Moore.

Terias blanda Boisduval, Spec. Gén. Lep. i, p. 672 (1836).

Terias blanda davidsoni Fruhstorfer, in Seitz, Macrolep. ix, p. 169, fig. 73 e (1910).

Widely spread over Siam.

Distribution: S. Burma; Tonkin; Siam; Annam.

The species ranges from Sikkim to the Malay Peninsula and Archipelago.

92. **Terias lacteola lacteola** Dist.

Terias lacteola Distant, Rhop. Malay., p. 466 (1886).

Terias lacteola lacteola Fruhstorfer, in Seitz, Macrolep. ix, p. 170, fig. 73 d (1910).

Muak Lek, E. Siam (*Fruhstorfer*).

Distribution: Siam; Annam; Malay Peninsula.

Fruhstorfer (*l. c.*) recognizes subspecies from Borneo, Sumatra, Java, the Natunas and Palawan.

93. **Terias sari sodalis** Fruhst.

Terias sari Horsfield, Cat. Lep. Mus. E. I. C., p. 136 (1829).

Terias sari sodalis Fruhstorfer, in Seitz, Macrolep. ix, p. 170 (1910).

Muak Lek, E. Siam (*Godfrey*); Nakon Sritamarat, P. Siam (*Kerr*).

Distribution: Siam; Mergui; Malay Peninsula.

The species ranges from India to the Greater Sunda Isles and the Philippines.

94. **Terias tilaha** Horsf.

Terias tilaha Horsfield, Cat. Lep. Mus. E. I. C., p. 136 (1829);

Fruhstorfer, in Seitz, Macrolep. ix, p. 170, fig. 73 g (1910).

J

Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula ; Borneo ; Java ; Sumatra.

95. *Gandaca harina burmana* Moore.

Terias harina Horsfield, Cap. Lep. Mus. E. I. C. p., 137 (1829).

Gandaca harina burmana Fruhstorfer, in Seitz, Macrolep. ix p. 175 (1910).

Widely spread over Siam.

Distribution : Burma ; Siam ; Annam.

The species ranges from Sikkim to Hainan and the Philippines, south through Malaya to New Guinea.

96. *Hebomoia glaucippe glaucippe* (Linn.).

Papilio glaucippe Linnaeus, Syst. Nat. x, p. 469 (1758).

Hebomoia glaucippe glaucippe Fruhstorfer, in Seitz, Macrolep. ix, p. 75 (1910).

Widely spread over Siam.

Distribution : Nepal to Burma ; Siam ; Malay Peninsula.

The species ranges from India to China and the Philippines, south to the Moluccas.

97. *Pareronia avatar paravata* Bingham.

Eronia avatar Moore, Cat. Lep. Mus. E.I.C., p. 61 (1857).

Pareronia avatar paravata Fruhstorfer, in Seitz, Macrolep. ix, p. 178, pl. 66a (1910).

Pak Jong, E. Siam.

Distribution : S. Burma ; Siam.

The typical form occurs from Sikkim to N. Burma.

98. *Pareronia valeria hippia* Fruhst.

Papilio valeria Cramer, Pap. Exot. i, pl 85, fig. a (1779).

Pareronia valeria hippia Fruhstorfer, in Seitz, Macrolep. ix, p. 178 pl. 66c (1910).

Widely spread over Siam.

The rare ♀-f *philomela* Fabr. has been obtained at Pak Jong, E. Siam and on the Pechaburi river, W. Siam.

Distribution : India ; Burma ; Siam.

The species extends south to the Malay Peninsula and Archipelago.

99. ***Udaina cynis cynis* Hew.***Udaina cynis* Hewitson, Exot. Butt. iii, pl. 8, fig. 54 (1866).*Udaina cynis cynis* Fruhstorfer, in Seitz, Macrolep. ix, p. 183 (1910).

3 ♂♂ : Bangnara, P. Siam, September 1923.

Distribution : P. Siam ; Malay Peninsula ; Sumatra.

A subspecies is found in Borneo.

DANAIDAE.

DANAINE.

100. ***Hestia lynceus reinwardti* Moore.***Hestia lynceus* Drury, Ill. Exot. Ent. ii, pl. 7, fig. 1 (1773).*Hestia lynceus reinwardti* Fruhstorfer in Seitz, Macrolep. ix, p. 220 (1910).Khao Ram and Khao Luang, Nakon Sritamarat, P. Siam (*Penillebury*) ; Khao Kalakiri, Pattani, P. Siam (*Kerr*).*Distribution* : P. Siam ; Malay Peninsula.

Other subspecies are found in Sumatra, Nias, Borneo, Java and the Natuna Islands.

101. ***Hestia hypermnestra linteata* Butl.***Hestia linteata* Butler, Trans. Linn. Soc. ser. 2, 1., p. 535 (1879).*Hestia hypermnestra linteata* Fruhstorfer, in Seitz, Macrolep. ix, p. 221 (1910)1 ♂ : Khao Pang-nga, c. 300 m., P. Siam, 27.2.29 (*Kerr*).*Distribution* : Mergui ; P. Siam ; Malay Peninsula.

The typical form occurs in Borneo and the Natuna Islands, and subspecies are known from Sumatra and Java.

102. ***Hestia hadeni* W-M. & de Nic.***Hestia hadeni* Wood-Mason and de Nicéville, Journ. As. Soc. Beng. ii, p. 242 (1880) ; Fruhstorfer, in Seitz, Macrolep. ix, p. 219, pl. 756 (1910).*Hestia lynceus hadeni* Evans, Journ. Bomb. N. H. S. xxxi (2), p. 519 (1923).

This record is based on a single male in the British Museum labelled " Bangkok, Siam."

Fruhstorfer (*l. c.*) states that the species is a very rare one, known only from Bassein at the estuary of the Irawaddi.

Evans (*l. c.*) treats *hadeni* as a subspecies of *H. lynceus*.

103. *Hestia leuconoe siamensis* Godf.

Idea leuconoe Erichson, Nov. Act. Ac. Nat. Cur. xvi, p. 283 (1834); Fruhstorfer (*Hestia*), in Seitz, Macrolep. ix, p. 221 (1910).

Hestia leuconoe siamensis Godfrey, Journ. N. H. S. Siam, ii, p. 17 (1916); Riley and Godfrey, op. cit. iv, pl. 7, fig. 6 (1921); Moulton, Journ. F.M.S. Mus. x, p. 161 (1921).

Nong Khor and Hup Bon, Sriracha forest, S. E. Siam. (Godfrey); Klung, Chantabun, S. E. Siam (Kerr).

Distribution : Siam ; Mergui Archipelago.

The species ranges from the Loo Choo Islands, Formosa and the Philippines to Malaysia.

H. l. siamensis is replaced in the southern half of the Malay Peninsula by *H. l. chersonesia* Fruhst. Of these two subspecies Moulton (*l. c.*) writes :—"It is probable that the original continental range of *leuconoe* has long been interrupted and the species has died out for some reason or other in the north of the Malay Peninsula leaving a gap between (and at the same time giving rise to) the present-day subspecies *chersonesia* and *siamensis*."

104. *Ideopsis daos perakana* Fruhst.

Ideopsis daos Boisduval, Spec. Gén. i, pl. 24, fig. 3 (1836).

Ideopsis daos perakana Fruhstorfer, in Seitz, Macrolep. ix, p. 26, pl. 76a (1910).

Khao Ram and Khao Luang, Nakon Sritamarat (Pendlebury); Khao Ton and Khao Keo Range, Satul (Kerr)—all in P. Siam.

Distribution : P. Siam ; Malay Peninsula.

The typical form occurs in Borneo and subspecies are found in Sumatra, Nias, the Natunas, and Palawan.

105. *Danaida chrysippus* (Linn.).

Papilio chrysippus Linnaeus, Syst. Nat. x, p. 471 (1758).

Distribution : India to China, south through the Malay Archipelago to Australia.

(a) *Danaida chrysippus chrysippus* (Linn.).

Danaida chrysippus chrysippus Fruhstorfer, in Seitz, Macrolep. ix, p. 193 (1910).

Widely spread over Siam proper.

Distribution: Ceylon; India; Burma; China; Siam.

(b) **Danaida chrysippus bataviana** Moore.

Danaida chrysippus bataviana Fruhstorfer, op. cit., p. 194.

Distribution: P. Siam; Malay Peninsula; Sumatra; Java.

106. Danaida plexippus plexippus (Linn.).

Papilio plexippus Linnaeus, Syst. Nat. x, p. 471 (1758).

Danaida plexippus plexippus Fruhstorfer, in Seitz, Macrolep. ix, p. 194 (1910).

Widely spread over Siam and very common.

Distribution: India to China and the Malay Peninsula.

The species extends through the Malay Archipelago to Australia.

107. Danaida melanippus hegesippus (Cram.).

Papilio melanippus Cramer, Pap. Exot. ii, pl. 127, figs. A. B. (1779).

Danaida melanippus hegesippus Fruhstorfer, in Seitz, Macrolep. ix, p. 196 (1910).

Muak Lek, E. Siam; Sriracha S. E. Siam; Bangkok, C. Siam.

Distribution: Siam; Malay Peninsula; Borneo; Sumatra; Natuna Isles.

The species ranges from India through Malaya to Java and the Celebes.

108. Danaida affinis malayana Fruhst.

Papilio affinis Fabricius, Syst. Ent. p. 511 (1775).

Danaida affinis malayana Fruhstorfer, in Seitz, Macrolep. ix, p. 201, pl. 77d (1910); Moulton, Journ. F. M. S. Mus. ix, p. 177, (1921).

Bangkok, C. Siam; Sriracha S. E. Siam.

Distribution: Siam; Malay Peninsula.

Padang Besar (Pendlebury); Kaw Samui (Kerr)—P. Siam.

The species extends east and south to the Philippines, Australia, and the Solomon Islands,

Moulton (1921) obtained *malayana* from Johore and Kuala Selangor, thus establishing the original habitat of the subspecies which for many years was doubtful.

109. **Danaida melissa septentrionis** (Butl.).

Papilio melissa Cramer, Pap. Exot. iv, pl. 377, figs. C. D (1782).

Danaida melissa septentrionis Fruhstorfer, in Seitz, Macrolep. ix, p. 202 (1910).

Widely spread over Siam.

Distribution: Kashmir to China and Formosa, south to the Malay Peninsula.

The typical form occurs in Java, and subspecies are found in Nias, Borneo, Fiji and the New Hebrides.

110. **Danaida gautama gautama** (Moore).

Danais gautama Moore, Ann. Mag. N. H. (4) xx, p. 43 (1877).

Danaida gautama gautama Fruhstorfer, in Seitz, Macrolep. ix, p. 203 (1910).

1 ♀ : Huey Tak Sao, W. Siam, 23.2.24 (*Stockley*).

Distribution: Burma; Mergui Archipelago; Siam.

A subspecies is found in the Nicobar Islands.

111. **Danaida limniace limniace** (Cram.).

Papilio limniace Cramer, Pap. Exot. i (1775).

Danaida limniace limniace Fruhstorfer, in Seitz, Macrolep. ix, p. 204 (1910).

Sriracha, S. E. Siam; Bangkok, C. Siam; Pechaburi River, W. Siam.

Distribution: From China south to Siam and the Malay Peninsula.

Subspecies are found in India, Borneo and the Celebes.

112. **Danaida aspasia aspasia** (Fabr.).

Papilio aspasia Fabricius, Mant. Ins. ii, p. 15 (1787).

Danaida aspasia aspasia Fruhstorfer, in Seitz, Macrolep. ix, p. 205 pl. 76 d (1910).

Pak Jong, E. Siam and Sriracha, S. E. Siam (*Godfrey*); Nakon Sritamarat, P. Siam (*Pendlebury*; *Kerr*).

Distribution: Burma; Siam; Malay Peninsula; Sumatra; Palawan.

Subspecies are found in Nias, Borneo, and Java.

113. *Danaida eryx eryx* (Fabr.).

Papilio eryx Fabricius, Ent. Syst. Suppl., p. 423 (1798).

Danaida eryx eryx Fruhstorfer, in Seitz, Macrolep. ix, p. 207, pl. 77 b (1910).

Distribution: Siam; Cochin-China; Malay Peninsula; Sumatra.

Subspecies are found in Java and Nias.

114. *Danaida aglea melanoides* (Moore).

Papilio aglea Cramer, Pap. Exot. iv, p. 377 (1781).

Parantica melanoides Moore, P. Z. S., p. 247 (1883).

Danais grammica Boisduval, Spéc. Gén. Lép. i. (1836).

Danaida aglea grammica Fruhstorfer, in Seitz, Macrolep. ix, p. 208 (1910).

Widely spread over Siam.

Distribution: Kashmir to Burma; Siam.

The typical form *aglea* is restricted to Ceylon, S. and C. India; and subspecies are known from Tonkin and Formosa.

Grammica Boisd. (1836) must be regarded as a synonym of *aglea* Cram. (1781).

115. *Danaida melaneus plataniston* Fruhst.

Papilio melaneus Cramer, Pap. Exot. i, pl. 30, fig. D (1775).

Danaida melaneus plataniston Fruhstorfer, in Seitz, Macrolep. ix, p. 210 (1910).

Me Song forest, Prae, and Chiengrai, N. Siam; Sriracha, S. E. Siam (*Godfrey*); Hat Sanuk, S. W. Siam (*Boden Kloss*); Me Wong, W. Siam (*Stockley*); Nakon Sritamarat, P. Siam (*Pendleberry*; *Kerr*).

Distribution: Sikkim to Burma; Siam; Malay Peninsula.

The typical form occurs in China and a subspecies is found in Java.

116. *Danaida sita* (Koll.).

Danais sita Kollar, in Hügel's Kashmir, iv, p. 424 (1844).

Distribution: Kashmir to China and Formosa, south to the Malay Peninsula.

(a) ***Danaida sita tira* Fruhst.**

Danaida sita tira Fruhstorfer, in Seitz, Macrolep. ix, p. 211 (1910).

Khun Than (3,500 ft.) and Doi Chang (4,500 ft.), N. Siam (*Godfrey*) ; Me Lamoung (2,000 ft.) W. Siam (*Stockley*).

Distribution : Assam ; S. Burma ; Siam.

(b) **Danaida sita ethologa** (Swinh.).

Danaida sita ethologa Fruhstorfer, op. cit., p. 241.

Khao Luang (5,760–5,800 ft.), Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula.

117. Danaida similis (Linn.).

Papilio similis Linnaeus, Mus. Ulr. p. 299 (1764).

Distribution : Loo Choo Islands to Palawan and Flores, and west through S. China and Malaysia to India and Ceylon.

(a) **Danaida similis persimilis** (Moore).

Danaida similis persimilis Fruhstorfer, in Seitz, Macrolep., ix, p. 211 (1910).

Widely spread over Siam and very common.

Distribution : Confined to Siam.

(b) **Danaida similis vulgaris** (Butl.).

Danaida similis vulgaris Fruhstorfer, op. cit., p. 211.

Bangnara, P. Siam (*Godfrey*) ; Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula ; Borneo ; Sumatra ; Natuna Islands.

EUPLOEINAE.

118. Euploea crameri marsdeni Moore.

Euploea crameri Lucas, Rev. Zool., p. 318 (1853).

Euploea crameri marsdeni Fruhstorfer, in Seitz, Macrolep. ix, p. 230 (1910).

Bangnara and Chumporn (*Godfrey*) ; Nakon Sritamarat (*Pendlebury*) ; Satul (*Kerr*)—all in P. Siam.

Distribution : P. Siam ; Malay Peninsula.

The species ranges from India, through Malaysia, to Bali.

119. Euploea malayica malayica (Butl.).

Crastia malayica Butler, Journ. Linn. Soc. xiv, p. 297 (1878).

Euploea malayica malayica Jordan, in Seitz, Macrolep. ix, p. 230, fig. 80b (1910).

1 ♂ : Wat Kiriwong, Nakon Sritamarat, P. Siam, 28.4.28 (*Kerr*).

Distribution : P. Siam ; Malay Peninsula ; Sumatra.

Subspecies are found in Nias, Java and Borneo.

120. **Euploea modesta modesta** Butl.

Euploea modesta Butler, Proc. Zool. Soc. Lond., p. 271 (1866).

Euploea modesta modesta Fruhstorfer, in Seitz, Macrolep. ix, p. 231 (1910).

Widely spread over Siam.

Distribution : Burma ; Siam ; Annam ; Malay Peninsula.

Subspecies are found in Sumatra and Borneo.

121. **Euploea camaralzaman** Butl.

Euploea camaralzaman Butler, Proc. Zool. Soc. Lond., p. 271 (1866); Fruhstorfer, in Seitz, Macrolep. ix, p. 231 (1910).

Muak Lek, E. Siam ; Chantabun, S. E. Siam ; Petchaburi River, W. Siam.

Distribution : Siam ; S. Burma.

122. **Euploea core circuita** Swin.

Papilio core Cramer, Pap. Exot. iii, p. 266 (1780).

Euploea circuita Swinhoe, Ann. Mag. N. H. iv, p. 499 (1903).

Euploea distanti circuita Fruhstorfer, in Seitz, Macrolep. ix, p. 236 (1910).

1 ♂ , 2 ♀ ♀ : Paknam Langsuan, Chumporn, P. Siam, Sep. 1 918

Distribution : Tonkin ; Cochin-China ; Siam.

The species ranges from India to the Malay Peninsula and Sumatra.

123. **Euploea godarti** Luc.

Euploea godarti Lucas, Rev. et Mag. Zool., p. 319 (1853); Fruhstorfer in Seitz, Macrolep. ix, p. 236 (1910).

Widely spread over Siam.

Distribution : Burma, Tenasserim ; Siam.

124. **Euploea layardi** Druce.

Euploea layardi Druce, Proc. Zool. Soc. Lond., p. 103, pl. 16, fig. 1 (1874) ; Bingham, Faun. Br. Ind. Butt. i, p. 34 (1905) ; Moulton, Journ. F. M. S. Mueums, x (3), p. 183 (1921).

Euploea godarti layardi Fruhstorfer, in Seitz, Macrolep. ix, p. 236 (1910).

Occurs with *E. godarti*, but is not so common.

Distribution : S. Burma ; Siam.

This species, which was originally described from Siam, differs from the preceding one as follows :—

(1). The violaceous-white patch in the apex of the fore wing so prominent in *godarti* is entirely absent and the spots are more obsolescent.

(2). The sexual brand on the fore wing of the male is quite distinct, both in length and breadth, from that of the male of *godarti*.

E. godarti is, however, an extremely variable species and it is quite probable that *layardi* may prove to be merely a seasonal form of it.

125. *Euploea orontobates* Fruhst.

Euploea orontobates Fruhstorfer, in Seitz, Macrolep. ix, p. 236 (1910).

Distribution : Known only from a single male taken by Fruhstorfer at Chantabun, S. E. Siam, in January 1901.

Fruhstorfer does not figure this species and his description is very meagre, but it is apparently intermediate between *E. godarti* and *E. camaralzaman*.

126. *Euploea alcathoe* (Godt.).

Danais alcathoe Godart, Ency. Méth. ix, p. 178 (1819).

Distribution : India to Siam, and through Malaysia to Lombok.

(a) Euploea alcathoe aesatia Fruhst.

Euploea alcathoe aesatia Fruhstorfer, in Seitz, Macrolep. ix, p. 237, pl. 85 b (1910).

Pak Jong, E. Siam ; Pechaburi River, W. Siam.

Distribution : S. Burma ; Siam.

Two males taken by Aagaard at Samsen, Bangkck, seem inseparable from those of *E. a. doubledayi* Feld. (Sikkim to the Shan States).

(b) Euploea alcathoe monticola Moulton.

Euploea alcathoe monticola Moulton, Journ. F. M. S. Museums x, p. 183 (1921); Pendlebury, op. cit., xi, p. 30 (1923).

Khao Ram (750 ft.), Nakon Sritamarat, P. Siam (Pendlebury).

Distribution : P. Siam Malay Peninsula.

Moulton (*l. c.*) describes *monticola* as a montane form of *E. alcathoe*, but expresses the opinion that it may prove to be a distinct species.

Pendlebury (*l. c.*) points out that its occurrence in P. Siam at the low altitude of 750 ft. tends to support this view.

127. *Euploea deione* Westw.

Euploea dione Westwood, Cab. Or. Ent., p. 76, pl. 37, fig. 3 (1848).

Distribution: Sikkim to the Malaya Peninsula, and through the Malayan Islands to Sumbawa and Palawan.

(a) ***Euploea deione limborgi* Fruhst.**

Euploea deione limborgi Fruhstorfer, in Seitz, Macrolep ix, p. 238 (1910).

Pak Jong, E. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam (*Godfrey*); Hat Sanuk, S. W. Siam (*F. M. S. Museums' Coll.*).

Distribution: Karens to S. Burma; Mergui; Siam; S. Annam.

(b) ***Euploea deione menetriesi* Feld.**

Euploea deione menetriesi Fruhstorfer, op. cit., p. 238.

Khao Ram and Khao Luang, P. Siam (*Pendlebury*).

Distribution: S. Burma; P. Siam; Malay Peninsula.

128. *Euploea dufresne harrisi* Feld.

Danais dufresne Godart, Ency. Méth. ix, Suppl., p. 815 (1823).

Euploea harrisi harrisi Fruhstorfer, in Seitz, Macrolep. ix, p. 246 (1910).

Euploea dufresne harrisi Moulton, Journ. F. M. S. Mus. x, p. 185 (1921).

Widely spread over Siam.

Distribution: Burma; Siam; Cochin-China; Malay Peninsula. Fruhstorfer (*l. c.*) recognizes as separate species: (a) *E. harrisi* Feld., from the Continent, (b) *E. lacordairei* Moore, from the Indo-Malayan Islands, and (c) *E. dufresne* Godt. from the Philippine Islands. He remarks:—"Whether all the three species now regarded as separate belong to one species is a question which can only be solved by further anatomical research".

Moulton (*l. c.*) treats all as one species, "subspecifically distinct from one another in their own particular regions"; and adopts the oldest name *dufresne* Godt. for the species.

129. **Euploea mulciber mulciber** (Cram.).

Papilio mulciber Cramer, Pap. Exot. ii, pl. 127 (1799).

Euploea mulciber mulciber Fruhstorfer, in Seitz, Macrolep. ix, p. 250 (1910).

Widely spread over Siam.

Distribution : India to China ; Siam ; Malay Peninsula.

Several well-marked subspecies are found in the Malay Archipelago.

130. **Euploea mazares ledereri** Feld.

Euploea mazares Moore, Cat. Lep. Mus. E. I. C. i, p. 127 (1857).

Euploea mazares ledereri Fruhstorfer, in Seitz, Macrolep. ix, p. 252 (1910).

Bangnara (*Godfrey*) ; Satul and Pattani (*Kerr*) ; Nakon Sritamarat (*Pendlebury*)—all in P. Siam.

Distribution : S. Burma ; P. Siam ; Malay Peninsula.

The species ranges from Formosa to the Solomon Islands.

131. **Euploea corus drucei** Moore.

Papilio corus Fabricius, Ent. Syst. iii, p. 41 (1793).

Euploea corus drucei Fruhstorfer, in Seitz, Macrolep. ix, p. 258 (1910).

Chantabun and Sriracha, S. E. Siam ; Bangkok, C. Siam.

Distribution : Known only from Siam.

The species ranges from Ceylon and Burma, through Malaysia, to Palawan and the Celebes.

132. **Euploea leucostictos leucogonyx** (Butl.).

Euploea leucostictos Gmelin, Syst. Nat. i (5), p. 289 (1788-1791).

Euploea leucostictos leucogonyx Fruhstorfer, in Seitz, Macrolep. ix, p. 263 (1910).

Pak Jong, E. Siam and Pechaburi River, W. Siam (*Godfrey*) ; Hat Sanuk, S. W. Siam (*F. M. S. Museums' Coll.*) ; Nakon Sritamarat, P. Siam (*Pendlebury*).

Distrbtion : Burma ; Siam ; Malay Peninsula.

The species ranges from India to Formosa and the Mariannes, south to Malaysia.

133. **Euploea midamus** (Linn.).

Papilio midamus Linnaeus, Mus. Ulr., p. 251 (1764).

Distribution: Sikkim to S. China and Tonkin, south to the Malay Peninsula.

(a) *Euploea midamus brahma* (Moore).

Euploea midamus brahma Fruhstorfer, in Seitz, Macrolep. ix, p. 270 (1910).

Me Song forest, Prae, N. Siam; Pak Jong and Muak Lek, E. Siam.

Distribution: Karens to S. Burma; Siam.

(b) *Euploea midamus dejani* (Moore).

Euploea midamus dejani Fruhstorfer, op. cit., p. 271.

Hat Sanuk and Tasan, S. W. Siam (*F. M. S. Museums' Coll.*).

Distribution: S. W. Siam and the Malay Peninsula.

134. *Euploea crassa crassa* Butl.

Euploea crassa Butler, Proc. Zool. Soc. Lond., p. 278 (1866).

Euploea klugi crassa Fruhstorfer, in Seitz, Macrolep. ix, p. 271, fig. 79 c (1910); id. (*E. crassa crassa*), op. cit., p. 278.

Widely spread over Siam.

Distribution: S. Burma; Siam.

Subspecies are found in India and Ceylon.

135. *Euploea diocletianus diocletianus* (Fabr.).

Papilio diocletiana Fabricius, Ent. Syst. iii, p. 40 (1793).

Euploea diocletianus diocletianus Fruhstorfer, in Seitz, Macrolep. ix, p. 271 (1910).

Widely spread over Siam.

Distribution: Assam to Burma; Siam; Malay Peninsula; Sumatra.

Subspecies are found in Nias, Borneo, Java and the Natuna Isles.

SATYRIDAE.

SATYRINAE.

136. *Ypthima huebneri* Kirby.

Ypthima huebneri Kirby, Cat. Di. Lep., p. 95 (1871); Fruhstorfer, in Seitz, Macrolep. ix, p. 287, fig. 99 e (1910).

Me Song forest, Prae, N. Siam; Pak Jong, E. Siam.

Distribution: Sikkim to Burma; Siam; Annam; Tonkin.

137. *Ypthima avanta* Moore.

Ypthima avanta Moore, Proc. Zool. Soc. Lond., p. 567 (1874).

Ypthima avanta avanta Fruhstorfer, in Seitz, Macrolep. ix, p. 288 (1910).

Me Song forest, Prae, N. Siam ; Muak Lek, E. Siam.

Distribution : Kashmir to Sikkim ; Siam.

Subspecies are found in W. China, in Ceylon and S. India ; and in Burma.

138. *Ypthima methora* sobrina Elw. & Edw.

Ypthima methora Hewitson, Trans. Ent. Soc. (3) ii, p. 291 (1864).

Ypthima sobrina Elwes and Edwards, Trans. Ent. Soc. Lond., p. 29 (1893) ; Fruhstorfer, in Seitz, Macrolep. ix, p. 288 (1910).

Ypthima methora sobrina Evans, Journ. Bomb. N. H. S. xxix, p. 787 (1923).

1 ♂ : Me Wong (600 ft.), W. Siam, 18. 2. 24 (*Stockley*).

Distribution : Karen Hills ; Siam.

The typical form occurs from Sikkim to Assam.

Fruhstorfer (*l. c.*) treats *sobrina* as a separate species confined to the Karen Hills.

139. *Ypthima baldus* baldus (Fabr.).

Papilio baldus Fabricius, Syst. Ent. App., p. 829 (1775).

Ypthima baldus baldus Fruhstorfer, in Seitz, Macrolep. ix, p. 289, pl. 99 d (1910).

Widely spread over Siam.

Distribution : Sikkim to Burma ; Tonkin ; Siam ; Annam.

The species ranges from Japan, Hongkong and India, south to the Greater Sunda Islands.

140. *Ypthima watsoni* (Moore).

Pandima watsoni Moore, Lep. Ind. ii, p. 89, pl. 113 (1893-96) ;

Fruhstorfer (*Ypthima*), in Seitz, Macrolep. ix, p. 291 (1910).

Me Wong (600 ft.) and Klong Klung (400 ft.), W. Siam (*Stockley*).

Distribution : Manipur to S. Burma ; Siam.

141. *Ypthima dohertyi* (Moore).

Thympia dohertyi Moore, Lep. Ind. ii, p. 65, pl. 108, figs. 1, 1 a (1893) ; Fruhstorfer, in Seitz, Macrolep. ix, p. 292 (1911).

1 ♂ : Klong Klung, W. Siam, 4. 4. 20 (*Stockley*).

Distribution: Shan States to Karen; Siam.

Fruhstorfer (*l. c.*) suggests that *dohertyi* may be merely a race of *Y. methora* Hew.

142. *Ypthima savara savara* H. G. Smith.

Ypthima savara H. G. Smith, A. M. N. H. (5) xx, p. 267 (1887).

Ypthima savara savara Fruhstorfer, in Seitz, Macrolep. ix, p. 292 (1911).

Doi Chang (4,500 ft.), Khun Than (2,500 ft.), and Me Song forest Prae—all in N. Siam (*Godfrey*); Umpang, W. Siam (*Stockley*).

Distribution: Burma; Siam.

Fruhstorfer (*l. c.*) recognizes a subspecies from Tonkin.

143. *Erites argentina* Butl.

Erites argentina Butler, Cat. Satyr. Brit. Mus., p. 188 (1868);

Fruhstorfer, in Seitz, Macrolep. ix, p. 302 (1911).

Moore (Lep. Ind. II, p. 114) states that this species has been taken on the Siam-Tenasserim border.

Distribution: S. Burma; Siam; Malay Peninsula; Borneo; Sumatra; Java.

144. *Erites angularis angularis* Moore.

Erites angularis Moore, Proc. Zool. Soc. Lond., p. 825 (1878).

Erites angularis angularis Fruhstorfer, in Seitz, Macrolep. ix, p. 303 (1911).

Kanburi, W. Siam (*Fruhstorfer*); Me Song forest, Prae, N. Siam (*Godfrey*).

Distribution: S. Burma; Siam; Malay Peninsula.

A subspecies is found in Sumatra.

145. *Mycalesis anapita* Moore.

Mycalesis anapita Moore, Cat. Lep. Mus. E. I. C. i, p. 232, No. 495 (1857); Moulton, Sarawak Mus. Journ. ii, No. 6, p. 206 (1915).

Mycalesis anapita anapita et fucentia Fruhstorfer, in Seitz, Macrolep. ix, p. 334 (1911).

1 ♂: Rai Chawng, Takuapa, P. Siam, 7. 2. 29 (*Kerr*).

Distribution: P. Siam; Malay Peninsula; Borneo; Sumatra; Bangka; Billiton.

Of *fucentia* Fruhst., Moulton (*l. c.*) writes:—"The black distal border of the hind wing is variable in a long Sarawak series before

Erratum.

Mycalesis anapita* Moore** should follow ***Coelites nothis
Boisd., on p. 248.

me, and is insufficient in development and constancy to warrant Fruhstorfer's separation as a separate race."

146. *Lethe rohria rohria* (Fabr.).

Papilio rohria Fabricius, Mant. Insec. ii, p. 45 (1787).

Lethe rohria rohria Fruhstorfer, in Seitz, Macrolep. ix, p. 315 (1911); Evans, Journ. Bomb. N. H. S. xxix (2), p. 533 (1923).

Khun Than (3,500 ft.), N. Siam (*Godfrey*); Pang Yao, W. Siam (*Stockley*).

Distribution : Sikkim to Burma ; Siam.

Evans (*l. c.*) recognizes as good subspecies : *rohria nilgiriensis* Guer. (Ceylon ; S. India) and *rohria dyrta* Feld. (Kashmir—Kumaon).

147. *Lethe europa niladana* Fruhst.

Papilio europa Fabricius, Syst. Ent. p. 500 (1775).

Lethe europa niladana Fruhstorfer, in Seitz, Macrolep. ix, p. 315 (1911).

Distribution : Sikkim to Burma ; Siam.

The species ranges from India and China to the Malay Peninsula, Borneo, Sumatra, Java and the Philippines.

148. *Lethe confusa gambara* Fruhst.

Lethe confusa Aurivillius, Ent. Tids, p. 142 (1897).

Lethe confusa gambara Evans, Journ. Bomb. N. H. S. xxix (2), p. 533 (1923).

Lethe rohria gambara Fruhstorfer, in Seitz, Macrolep ix, p. 315 (1911).

Me Song forest, Prae, N. Siam (*Godfrey*); Me Lamoung (2,000 ft.) and Pang Yao, W. Siam (*Stockley*).

Distribution : Assam to Burma ; Siam.

The typical form occurs from Murree to Sikkim.

Fruhstorfer (*l. c.*) treats *gambara* as a subspecies of *rohria* Fabr., but Evans (*l. c.*) refers it to *confusa*.

149. *Lethe minerva tritogeneia* Fruhst.

Papilio minerva Fabricius, Syst. Ent. p. 493 (1775).

Lethe minerva tritogeneia Fruhstorfer, in Seitz, Macrolep. ix, p. 318 (1911).

Me Song forest, Prae, N. Siam.

Distribution : Chin Hills to S. Burma ; Siam.

The typical form occurs in Java and Sumatra.

150. *Lethe kansa* (Moore).

Debis kansa Moore, Cat. Lep. Mus. E.I.C., p. 220 (1857); Fruhstorfer (*Lethe*), in Seitz, Macrolep. ix, p. 318 (1911); Evans (*Lethe*), Journ. Bomb. N. H. S. xxix (2), p. 536 (1923).

Distribution: Sikkim to Burma; Siam.

Fruhstorfer's races *zeugitana* (from Assam) and *vaga* (from Tenasserim) are not recognized by Evans.

151. *Lethe vindyha* (Feld.).

Debis vindyha Felder, Wien. ent. Monats. iii, p. 402 (1859); Fruhstorfer (*Lethe*), in Seitz, Macrolep. ix, p. 319, fig. 98 b (1911).

Chiengrai (*Godfrey*); Chiengmai (*Tennent*)—N. Siam.

Distribution: Sikkim to Burma; Siam.

152. *Lethe mekara zuchara* Fruhst.

Debis mekara Moore, Cat. Lep. Mus. E. I. C., p. 291 (1857).

Lethe mekara zuchara Fruhstorfer, in Seitz, Macrolep. ix, p. 319 (1911).

Me Song forest, Prae, N. Siam (*Godfrey*); Me Wong, W. Siam (*Stockley*).

Distribution: Assam to Burma; Siam.

The species ranges from Sikkim to the Malay Peninsula, Borneo and Sumatra.

153. *Lethe chandica flanona* Fruhst.

Debis chandica Moore, Cat. Lep. Mus. E. I. C. p. 219 (1857).

Lethe chandica flanona Fruhstorfer, in Seitz, Macrolep. ix, p. 320 (1911).

1 ♂: Me Lamoung, W. Siam, April 1920 (*Stockley*); 1 ♀: Me Song forest, Prae, N. Siam, April 1918 (*Godfrey*).

Distribution: Assam to Burma; Siam.

The species ranges from India to Hainan, Formosa and the Philippines, south to the Malay Peninsula, Sumatra and Borneo.

154. *Lethe verma sintica* Fruhst.

Satyrus verma Kollar, in Hügel's Kaschmir, iv, p. 447 (1844).

Lethe verma sintica Fruhstorfer, in Seitz, Macrolep. ix, p. 324 (1911).

Me Song forest, Prae, N. Siam.

Distribution: Sikkim to N. Burma; N. Siam.

The species ranges from India to China, Tonkin, Hainan and Formosa, south to Siam.

155. **Lethe muirheadi bhima** (Marsh.).

Lassiommata muirheadi Felder, Wien. ent. Monats. p. 28 (1862).

Neope muirheadi bhima Fruhstorfer, in Seitz, Macrolep. ix, p. 326 (1911).

Me Song forest, Prae, N. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*).

Distribution : N. Burma to Karen; Siam.

The typical form occurs in Central and West China and a subspecies is found in Tonkin.

156. **Neorina lowi neophyta** Fruhst.

Cyllo lowii Doubleday and Hewitson, Gen. Diurn. Lep. p. 369, (1851).

Neorina lowi neophyta Fruhstorfer, in Seitz, Macrolep. ix, p. 326 (1911).

1 ♂ : Kamala, Puket (c. 200 m.), P. Siam, 10.3.29 (*Kerr*).

Distribution : P. Siam; Perak.

Fruhstorfer (*l. c.*) refers the typical form to Borneo, and recognizes subspecies from Nias, Sumatra, Palawan and Paragua.

157. **Anadebis himachala** (Moore).

Theope himachala Moore, Cat. Lep. Mus. E. I. C, i, p. 234 (1857);

Fruhstorfer, in Seitz, Macrolep. ix, p. 328, fig. 94 a (1911).

1 ♂ : Chiengrai, N. Siam, April 1921.

Distribution : Sikkim to N. Burma; N. Siam.

158. **Anadebis diademooides diademooides** (Moore).

Zethera diademooides Moore, P. Z. S., p. 824 (1878).

Anadebis diademooides diademooides Fruhstorfer, in Seitz, Macrolep. ix, p. 328 (1911).

2 ♂♂ : Klong Klung, W. Siam, 4.4.20 (*Stockley*).

Distribution : Karen to S. Burma; Siam.

159. **Anadebis batmara** Fruhst.

Anadebis batmara Fruhstorfer, Iris, xx, p. 253 (1907).

Anadebis diademooides batmara Fruhstorfer, in Seitz, Macrolep. ix, p. 328, pl. 94 a (1911).

Me Song forest, Prae, N. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*).

Distribution : Tonkin; Siam.

Fruhstorfer (*l. c.*) splits the species *diademooides* into three subspecies: *diademooides* Moore from Burma, Tenasserim and Pegu; *batmara* Fruhst. from Tonkin; and *henrici* Holl. from Hainan.

Riley* is of the opinion that *batmara* is a good and distinct species. Fruhstorfer himself remarks:—"batmara is very distinct from all its allies, particularly in the 5 elongate, white subapical ultracellular spots of the forewing, which are produced on the under surface (as well as all the other white markings)."

160. Coelites nothis nothis Boisd.

Coelites nothis Boisduval, Gen. Di. Lep. ii, p. 368 (1851).

Coelites nothis nothis Fruhstorfer, in Seitz, Macrolep. ix, p. 328 (1911).

Muak Lek, E. Siam; Khao Sebap, S. E. Siam.

Distribution: Confined to Siam.

Subspecies are found in Burma and Tonkin.

161. Mycalesis mnasicles perna Fruhst.

Mycalesis mnasicles Hewitson, Exot. Butt. iii, pl. 5, figs. 32, 33 (1864).

Mycalesis mnasicles perna Fruhstorfer, in Seitz, Macrolep. ix, p. 340, fig. 91 e (1911).

Me Song forest, Prae, N. Siam.

Distribution: Shan States; Burma; Siam; Malay Peninsula.

The typical form occurs in Borneo and Sumatra.

162. Mycalesis perseus perseus (Fabr.).

Papilio perseus Fabricius, Syst. Ent. p. 488 (1775).

Mycalesis perseus perseus Fruhstorfer, in Seitz, Macrolep. ix, p. 343, fig. 92 a (1911).

Widely spread over Siam.

Distribution: India to Tonkin, Hainan and Formosa, south to Siam and Annam.

The species extends to Australia.

163. Mycalesis mineus (Linn.).

Papilio mineus Linnaeus, Syst. Nat. i, p. 768 (1767).

Distribution: India to Formosa and the Philippines, south to the Malay Peninsula and Sumatra.

* Riley, *in litt.*, 13. 12. 23.

(a) **Mycalesis mineus mineus** (Linn.).

Mycalesis mineus mineus Fruhstorfer, in Seitz, Macrolep. ix, p. 344 (1911).

Widely spread over Siam.

Distribution : Kulu to Burma ; Tonkin ; Siam ; Annam.

(b) **Mycalesis mineus macromalayana** Fruhst.

Mycalesis mineus macromalayana Fruhstorfer, op. cit., p. 344.

Khao Ram, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula ; Singapore, Sumatra.

164. **Mycalesis perseoides** (Moore).

Calysisme perseoides Moore, Lep. Ind. p. 179 (1890-92); Fruhstorfer (*Mycalesis*), in Seitz, Macrolep. ix, p. 346, 92 a (1911).

Widely spread over Siam.

Distribution : S. India ; Burma ; Tonkin ; Siam ; Annam.

165. **Mycalesis visala neovisala** Fruhst.

Mycalesis visala Moore, Cat. Lep. Mus. E. I. C., p. 230 (1857).

Mycalesis visala neovisala Fruhstorfer, in Seitz, Macrolep. ix, p. 346, fig. 91 f (1911).

1 ♂, 1 ♀ : Chiengrai, N. Siam, 24.4.21.

Distribution : Burma ; Tonkin ; Siam.

The typical form occurs from Sikkim to Assam, and a subspecies is found in the Andamans.

166. **Mycalesis lepcha** (Moore).

Semanta lepcha Moore, Trans. Ent. Soc. Lond. p. 167 (1880).

Mycalesis lepcha lepcha Evans, Journ. B. N. H. S. xxix, p. 530 (1923).

Mycalesis mamerta lepcha Fruhstorfer, in Seitz, Macrolep. ix, p. 437 (1911).

1 ♂ : Huey Tak Sao ; 1 ♂ : Upper Me Klong (1,500 ft.), W. Siam (*S'ockley*).

Distribution : Kulu to Kumaon ; Siam.

Fruhstorfer (*l. c.*) treats *lepcha* as a subspecies of *mamerta* Cram., but both Riley* and Evans (*l. c.*) regard it as a separate species. Evans recognizes subspecies of it from the Palnis, from Orissa, and from Assam.

* Riley, *in litt.*, 13, 12, 23.

167. **Mycalesis mamerta annamitica** Fruhst.*Papilio mamerta* Cramer, Pap. Exot. iv, p. 75 (1780).*Mycalesis mamerta annamitica* Fruhstorfer, in Seitz, Macrolep. ix, p. 347 (1911).

Doi Chang (4,500 ft.); Khun Than (3,500 ft.); Me Song forest, Prae—all in N. Siam. Also obtained at Ban Na Sao in French Laos.

Distribution: Karens to Burma; Siam; S. Annam; French Laos.

The typical form occurs from Assam to the Shan States

168. **Mycalesis malsara f. rufis Moore.***Mycalesis malsara* Moore, Cat. Lep. Mus. E. I. C., p. 231 (1857).*Mycalesis mamerta malsara f. rufis* Fruhstorfer, in Seitz, Macrolep. ix, p. 347 (1910).

1 ♀: Me Wong, W. Siam, 7.4.20 (Stockley).

Distribution: Sikkim to Burma; Siam.Fruhstorfer (*l. c.*) treats *malsara* as a subspecies of *mamerta* Cram.; Riley* regards it as a good species.169. **Mycalesis gotama charaka** Moore.*Mycalesis gotama* Moore, Cat. Lep. Mus. E. I. C., p. 232 (1857).*Mycalesis gotama charaka* Fruhstorfer, in Seitz, Macrolep. ix, p. 349 (!911).

6 ♂♂: Me Song forest, Prae, N. Siam, April 1916.

Distribution: Assam to Burma; Tonkin; Siam; Annam.

The typical form occurs in China, and subspecies are found in Japan.

170. **Mycalesis fuscum fuscum** (Feld.).*Dasyomma fuscum* Felder, Wien. ent. Monats, iv, p. 401 (1860).*Mycalesis fuscum fuscum* Fruhstorfer, in Seitz, Macrolep. ix, p. 305 (1911).

1 ♂: Bangnara, P. Siam, 3. 9. 18.

Distribution: S. Burma; Siam; Malay Peninsula; Sumatra.

Subspecies are found in Borneo, Nias and Java.

171. **Mycalesis orseis nautilus** Butl.*Mycalesis orseis* Hewitson, Exot. Butt. iii, p. 89 (1862).*Mycalesis orseis nautilus* Fruhstorfer, in Seitz, Macrolep. ix, p. 352 (1911); Evans, Journ. Bomb. N. H. S. xxix (2), p. 527 (1923).*Riley, *in litt.*, 13. 12. 23.

Khao Ram and Khao Luang, Nakon Sritamarat (*Pendlebury*);
Khao Kalakiri, Pattani (*Kerr*)—P. Siam.

Distribution: Naga Hills to Burma; P. Siam.

Fruhstorfer (*l. c.*) refers typical *orseis* to Sumatra and
recognizes subspecies from Borneo, Nias, and the Celebes.

172. *Mycalesis anaxias aemate* Fruhst.

Mycalesis anaxias Hewitson, Exot. Butt. iii, p. 86 (1862).

Mycalesis anaxias aemate Fruhstorfer, in Seitz, Macrolep. ix, p. 353
(1911).

Me Song forest, Prae, N. Siam and Sriracha forest, S. E. Siam
(*Godfrey*); Umpang, W. Siam (*Stockley*); Khao Kalakiri, c. 900 m.,
Pattani, P. Siam (*Kerr*).

Distribution: Sikkim to Siam.

The species occurs from Sikkim to the Malay Peninsula, and
in the Andaman and Nicobar Islands.

173. *Mycalesis adamsoni* Wats.

Mycalesis adamsoni Watson, Journ. Bomb. N. H. S. x, p. 640
(1896); Fruhstorfer, in Seitz, Macrolep. ix, p. 353 (1911).

7 ♂♂, 3 ♀♀ : Me Song forest, Prae, April 1916 and 1918.

Distribution: N. Burma; Tonkin; N. Siam.

174. *Mycalesis siamica* Ril. & Godf.

Mycalesis siamica Riley and Godfrey, Journ. N. H. S. Siam, iv, p.
170, pl. 4, figs. 2, 3 (1921).

3 ♀♀ : Me Song forest, Prae, N. Siam, April 1916; 10 ♂♂,

4 ♀♀ : April 1918.

Distribution: Known only from Siam.

175. *Mycalesis anaxioides* Marsh.

Mycalesis anaxioides Marshall, in M. & de Nicév., Butt. Ind. i, p.
107 (1883); Fruhstorfer, in Seitz, Macrolep. ix, p. 354 (1911).

6 ♂♂, 1 ♀ : Me Song forest, Prae, N. Siam, April 1916 and
1918.

Distribution: Karen to S. Burma; N. Siam.

176. *Mycalesis francisca sanatana* Moore.

Papilio francisca Cramer, Pap. Exot. pl. 326, figs. E, F (1782).

Mycalesis francisca sanatana Fruhstorfer, in Seitz, Macrolep. ix, p.
354 (1911).

3 ♂ ♂ : Doi Chang (4,500 ft.), N. Siam, 13. 4. 21.

Distribution : Kulu to Burma ; N. Siam.

The species ranges from India to China and Japan, south to Siam and Annam.

177. **Mycalesis maianeas maianeas** Hew.

Mycalesis maianeas Hewitson, Exot. Butt. iii, p. 87 (1862).

Mycalesis maianeas maianeas Fruhstorfer, in Seitz, Macrolep. ix, p. 355 (1910).

Khao Ram and Khao Luang, Nakon Sritamarat (*Pendlebury*) ; Khao Luang (*Kerr*)—P. Siam.

Distribution : P. Siam ; Malay Peninsula ; Borneo.

A subspecies is found in Sumatra and Bangka.

178. **Mycalesis mystes tunicula** Fruhst.

Mycalesis mystes de Nicéville, Journ. B. N. H. S. vi, p. 343 (1891).

Mycalesis mystes tunicula Fruhstorfer, in Seitz, Macrolep. ix, p. 355, fig. 93 a (1911).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution : Confined to Siam.

Fruhstorfer (*l. c.*) regards *mystes* as a rare species of which he recognizes two subspecies: *mystes* de Nicév. from Burma, and *tunicula* Fruhst. from Siam. Riley* is of the opinion that both *mystes* and *tunicula* are subspecies of *M. sangaica* Butl. from S. China.

179. **Mycalesis oroatis** Hew.

Mycalesis oroatis Hewitson, Exot. Butt. iii (1862).

Distribution : Burma to the Malay Peninsula, Sumatra and Java.

(a) **Mycalesis oroatis surkha** Marsh.

Mycalesis oroatis surkha Fruhstorfer, in Seitz, Macrolep. ix, p. 358 (1911).

1 ♂ : Eka Klo, W. Siam, 3. 4. 20 (*Stockley*).

Distribution : Dawnas to S. Burma ; Siam.

The typical form occurs in Java.

(b) **Mycalesis oroatis ustulata** Dist.

*Riley, *in litt.*, 13. 12. 23.

Mycalesis oroatis ustulata Fruhstorfer, op. cit., p. 358, pl. 93 a.

Khao Luang (2,000 ft.) Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: P. Siam; Malay Peninsula; Sumatra.

180. *Mandarinia regalis baronesa* Fruhst.

Mandarinia regalis Leech, Trans. Ent. Soc. Lond., p. 102 (1889).

Mandarinia regalis baronesa Fruhstorfer, in Seitz, Macrolep. ix, p. 359, fig. 93 c (1911).

2 ♂♂: Chiengmai district, N. Siam, Jan. 1921 (*Tennent*).

Disiribution: Tonkin; Shan States; N. Siam.

The typical form occurs in China.

181. *Orsotriaena medus medus* (Fabr.).

Papilio medus Fabricius, Syst. Ent. p. 488 (1775).

Orsotriaena medus medus Fruhstorfer, in Seitz, Macrolep. ix, p. 358 (1911).

Widely spread over Siam.

Distribution: Sikkim to the Greater Sunda Islands.

The species extends to Australia.

182. *Ragadia crisilda critolaus* de Nicév.

Ragadia critolaus de Nicéville, Journ. B. N. H. S vii, p. 322 (1893).

Ragadia crisilda critolaus Fruhstorfer, in Seitz, Macrolep. ix, p. 360 (1911).

2 ♂♂: Me Wong (2,500 ft.), W. Siam, April 1920 (*Stockley*).

Distribution: Dawnas to S. Burma; Siam.

The typical form occurs in Assam, and subspecies are found in W. China and S. Burma.

183. *Ragadia critias* Ril. & Godf.

Ragadia critias Riley and Godfrey, Journ. N. H. S. Siam, iv, p. 171, pl. iv, fig. 4, ♂ (1921).

1 ♂, 1 ♀: Nam Pat, 36 miles E. of Utaradit, N. E. Siam, Jan. 1920.

Distribution: Known only from Siam.

184. *Ragadia crisia siponta* Fruhst.

Ragadia crisia Hübner, Zutr. Smlg. Exot. Schmett. figs. 675, 676 (1832).

Ragadia crisia siponta Fruhstorfer, in Seitz, Macrolep. ix, p. 361 (1911).

Khao Ram, Nakon Sritamarat (*Pendlebury*); Klawng Hin Kao, Patalung (*Kerr*)—P. Siam.

Distribution: P. Siam; Malay Peninsula; Natuna Islands.

The typical form occurs in Java and subspecies are found in Borneo and Sumatra.

185. *Melanitis leda ismene* (Cram.).

Papilio leda Linnaeus, Syst. Nat. i (2), p. 773 (1767).

Melanitis leda ismene Fruhstorfer, in Seitz, Macrolep. ix, p. 362 (1911).

Widely spread over Siam.

Distribution: Ceylon; India; Burma; China; Siam; Malay Peninsula; Borneo; Sumatra.

Other subspecies occur from Java and the Celebes to Australia and Madagascar.

186. *Melanitis phedima ganapati* Fruhst.

Papilio phedima Cramer, Pap. Exot. iv, pl. 292 (1782).

Melanitis phedima ganapati Fruhstorfer, in Seitz, Macrolep. ix, p. 363 (1911).

Me Song forest, Prae, N. Siam (*Godfrey*); Kambengpet, W. Siam (*Stockley*).

Distribution: Karen Hills to S. Burma; Tonkin; Siam; Annam.

The species ranges from India to China, Formosa and the Philippines, south to the Malay Peninsula, Sumatra and Nias.

187. *Melanitis zitennius zitennius* (Herbst).

Papilio zitennius Herbst, Nat. Syst. Schmett. viii, p. 5 (1796).

Melanitis zitennius zitennius Fruhstorfer, in Seitz, Macrolep. ix, p. 366 (1911).

Me Song forest Prae, N. Siam and Sriracha forest, S. E. Siam (*Godfrey*); Klong Suan Mak, W. Siam (*Stockley*).

Distribution: Kumaon to Burma; Siam; Malay Peninsula.

The species ranges from India and Tonkin south to the Greater Sunda Islands, Lombok and Sumbawa.

ELYMNIINAE.

188. *Elymnias dara daedalion* de Nicév.

Elymnias dara Distant, Ann. Mag. Nat. Hist. 5 (xix), p. 50 (1887).

Elymnias dara daedalion Fruhstorfer, in Seitz, Macrolep. ix, p. 373 (1911).

Me Song forest, Prae, N. Siam.

Distribution: Dawnas to S. Burma; Siam.

Other subspecies are found in Borneo, Sumatra, Java and Palawan.

189. *Elymnias patna patna* (Westw.).

Melanitis patna Westwood, in Dbdly., Westw. & Hew. Gen. Di. Lep., p. 405, pl. 68, fig. 2 (1851).

Elymnias patna patna Fruhstorfer, in Seitz, Macrolep. ix, p. 374 (1911).

1 ♀ : Chiengrai, N. Siam.

Distribution: Kumaon to N. Burma; N. Siam.

The species is also known from Tonkin and Malacca.

190. *Elymnias hypermnestra* (Linn.).

Papilio hypermnestra Linnaeus, Amoen. Acad. vi, p. 407 (1764).

Distribution: India to Tonkin, Hainan and Formosa, south to the Malay Peninsula and Java.

The following forms of *hypermnestra* are known from Siam:—

- (a) *Elymnias hypermnestra violetta* f. *violetta* Fruhst.

Elymnias hypermnestra violetta f. *violetta* Fruhstorfer, in Seitz, Macrolep. ix, p. 375, pl. 87 a (1911).

Distribution: Confined to Siam, and very rare.

- (b) *Elymnias hypermnestra violetta* f. *epixantha* Fruhst.

Elymnias hypermnestra violetta f. *epixantha* Fruhstorfer, op. cit., p. 375.

Me Song forest, Prae, N. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam (Godfrey); Klong Klung, W. Siam (Stockley).

Distribution: Siam; (?) Cambodia.

- (c) *Elymnias hypermnestra meridionalis* Fruhst.

Elymnias hypermnestra meridionalis Fruhstorfer, op. cit., p. 375.

Nong Yai Boo, Sriracha forest, S. E. Siam.

Distribution : Saigon ; Siam ; Annam.

(d) **Elymnias hypermnestra discrepans** Dist.

Elymnias nigrescens discrepans Fruhstorfer, op. cit. p. 378.

4 ♀ ♀ : Bangnara, P. Siam, Sep. 1923.

Distribution : P. Siam ; Malay Peninsula ; Singapore.

191. **Elymnias nesaea** (Linn.).

Papilio nesaea Linnaeus, Mus. Ulric. p. 302 (1764).

Distribution : Sikkim to the Greater Sunda Islands.

The following subspecies have been identified from Siam :—

(a) **Elymnias nesaea apelles** Fruhst.

Elymnias nesaea apelles Fruhstorfer, in Seitz, Macrolep. ix, p. 379, fig. 87 c (1911).

Bangkok, C. Siam.

Distribution : This very distinct subspecies is confined to Siam and, as far as I know, has been obtained only in Bangkok, where it is not uncommon.

(b) **Elymnias nesaea lioneli** Fruhst.

Elymnias nesaea lioneli Fruhstorfer, op. cit., p. 379.

Bangnara, P. Siam (*Godfrey*) ; Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula.

(c) **Elymnias nesaea timandra** Wall.

Elymnias nesaea timandra Fruhstorfer, op. cit., p. 379.

Chiengmai, N. Siam (*Tennent*).

Distribution : Sikkim to N. Burma ; N. Siam.

192. **Elymnias malelas ivena** Fruhst.

Melanitis malelas Hewitson, Exot. Butt. iii, pl. 1, figs. 6, 7 (1863).

Elymnias malelas ivena Fruhstorfer, in Seitz, Macrolep. ix, p. 381 (1911).

Pak Jong, E. Siam ; Bangkok, C. Siam.

Distribution : Tonkin ; Siam.

The species ranges from Sikkim to Tonkin, south to the Malay Peninsula.

193. *Elymnias penanga chelensis* de Nicév.

Melanitis penanga Westwood, in Dblday., Westw & Hew. Gen. Di. Lep., p. 405 (1851).

Elymnias chelensis de Nicéville, Journ Bomb. N. H. Soc., p. 200 (1890).

Elymnias penanga chelensis Fruhstorfer, in Seitz, Macrolep. ix, p. 384. fig. 88 C (1911).

1 ♂: Terutao, P. Siam, 20. 1. 28 (*Kerr*).

Distribution: Assam to Burma; P. Siam.

The species ranges from Assam to the Malay Peninsula, Borneo and Sumatra.

194. *Elymnias vasudeva oberthüri* Fruhst.

Elymnias vasudeva Moore, Cat. Lep. Mus. E. I. C. i, p. 238 (1857).

Elymnias vasudeva oberthüri Fruhstorfer, in Seitz, Macrolep. ix, p. 392 (1911).

Renong, P. Siam (*Doherty*).

Distribution: Fruhstorfer (*l. c.*) restricts this subspecies to Siam. The typical form occurs in Sikkim and subspecies are found in Assam and Burma.

AMATHUSIINAE.

195. *Faunis arcesilaus arcesilaus* (Fabr.).

Papilio arcesilaus Fabricius, Mant. Ins. ii, p. 28 (1787).

Faunis arcesilaus arcesilaus Fruhstorfer, in Seitz, Macrolep. ix, p. 404 (1911).

Me Song forest, Prae, N. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*); Khao Kalakiri, Pattani, P. Siam (*Kerr*); Me Wong, W. Siam (*Stockley*).

Distribution: Sikkim to the Malay Peninsula.

Subspecies are found in Borneo, Sumatra and Java.

196. *Faunis eumeus incerta* (Staud.).

Papilio eumeus Drury, Ill. Exot. Ent. i, p. 4 (1773).

Faunis eumeus incerta Fruhstorfer, in Seitz, Macrolep. ix, p. 406, fig. 100 a (1911).

2 ♂♂: Me Song forest, Prae, N. Siam, April 1919.

Distribution: S. Burma; Siam; Annam.

The typical form occurs in S. China and Hainan.

197. **Faunis faunula faunula** (Westw.).

Melanocyma faunula Westwood, Gen. Di. Lep., p. 234 (1851).

Faunis faunula faunula Fruhstorfer, in Seitz, Macrolep. ix, p. 406, fig. 100 a (1911).

Koh Chang (*Godfrey*) ; Khao Sebap (*Kerr*)—S. E. Siam.

Distribution : Siam ; Malay Peninsula ; Cambodia.

A subspecies is found in N. Burma.

198. **Aemonia lena** Atk.

Aemonia lena Atkinson, Proc. Zool. Soc. Lond., p. 215 (1871) ; Fruhstorfer, in Seitz, Macrolep. ix, p. 407, fig. 102 b (1911).

1 ♂ : Me Lamoung (3,000 ft.), W. Siam, 14. 4. 20 (*Stockley*) ;

2 ♂ ♂ : Doi Chang (4,500 ft.), N. Siam, 12. 4. 21 (*Godfrey*).

Distribution : Shan States ; S. Yunnan ; Siam.

199. **Xanthotaenia busiris busiris** Westw.

Xanthotaenia busiris Westwood, Trans. Ent. Soc. Lond. p. 187 (1856).

Xanthotaenia busiris busiris Fruhstorfer, in Seitz, Macrolep. ix, p. 407 (1911).

Bangnara (*Godfrey*) ; Khao Kalakari, Pattani ; Bang-to, Panga ; Klong Kampuam and Kraburi, Renong (*Kerr*) ; Khao Ram, Nakon Sritamarat (*Pendlebury*)—all in P. Siam.

Distribution : Dawnas to S. Burma ; P. Siam ; Malay Peninsula.

Subspecies are found in Borneo, Sumatra, and Nias.

200. **Stichophthalma louisa siamensis** Roths.

Stichophthalma louisa Wood-Mason, Proc. As. Soc. Beng. p. 163 (1871) ; Fruhstorfer, in Seitz, Macrolep. ix, p. 425 (1911).

Stichophthalma louisa siamensis Rothschild, Nov. Zool. xxiii, p. 308 (1916).

Distribution : Rothschild (*l. c.*) restricts this subspecies to Siam, but does not state from what part of the country it came, or by whom it was obtained.

The typical form occurs from the Karen Hills to S. Burma and subspecies are found in Tonkin and N. Burma.

201. **Stichophthalma camadeva camadevooides** de Nicév.

Thaumantis camadeva Westwood, Cab. Or. Ent., p. 9, pl. 4, figs. 1, 2.

Stichophthalma camadeva camadevooides Fruhstorfer, in Seitz, Macrolep. ix, p. 426 (1911).

2 ♀ ♀: Doi Sutep, N. Siam, March 1921 (*Tennent*).

Distribution: N. Burma; N. Siam.

The typical form is found in Sikkim and subspecies are known from the Naga Hills, and Assam.

202. *Stichophthalma cambodia editha* Ril. & Godf.

Stichophthalma cambodia Hewitson, Exot. Butt. iii (1862); Fruhstorfer, in Seitz, Macrolep. ix, p. 425 (1911).

Stichophthalma cambodia editha Riley and Godfrey, Journ. N. H. S. Siam, iv, p. 173, pl. 4, fig. 5 (1921).

Hup Bon, Sriracha forest and Khao Sebap, S. E. Siam.

Distribution: Known only from Siam.

S. cambodia was described by Hewitson from a single ♀ (from Cambodia) which is in the British Museum and which is still unique.

203. *Stichophthalma godfreyi* Roths.

Stichophthalma godfreyi Rothschild, Ann. Mag. Nat. Hist. xxii, p. 474 (1916).

Kanburi, W. Siam.

Distribution: Siam; Tenasserim.

This species was described from a single ♀ obtained on May 8th. 1914 in dense evergreen forest about 40 miles north-west of Kanburi, and some 4 miles from the Tenasserim boundary.

Other specimens have since been obtained by Ollenbach* in the Tavoy district, Tenasserim.

204. *Amathusia phidippus* (Johans.).

Papilio phidippus Johanssen, Amoen. Acad. vi, p. 402 (1764).

Distribution: S. India to the Malay Peninsula, Borneo, Sumatra, Java, and the Celebes.

(a) ***Amathusia phidippus adustatus* Fruhst.**

Amathusia phidippus adustatus Fruhstorfer, in Seitz, Macrolep. ix, p. 427 (1911).

Pak Jong, E. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam.

* Ollenbach, Journ. Bomb. N. H. S. xxvi, p. 867, (1920); xxvii, p. 887 (1921).

Distribution: Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

(b) **Amathusia phidippus chersias** Fruhst.

Amathusia phidippus chersias Fruhstorfer, op. cit., p. 427.

Wat Kiriwong, Nakon Sritamarat, P. Siam (*Pendlebury*; *Kerr*).

Distribution: P. Siam; Malay Peninsula.

205. Amathuxidia amythaon (Doubl.).

Amathusia amythaon Doubleday, A. N. N. H. xix, p. 175 (1847).

Distribution: Sikkim to the Malay Peninsula; Borneo; Sumatra; Java; Philippines.

(a) **Amathuxidia amythaon amythaon** (Doubl.).

Amathuxidia amythaon amythaon Fruhstorfer, in Seitz, *Macrolep.* ix, p. 432 (1911).

Me Song forest, Prae (*Godfrey*); Doi Sutep (*Tennent*)—N. Siam.

Distribution: Sikkim to N. Siam.

(b) **Amathuxidia amythaon dilucida** (Honr.).

Amathusia dilucida Honrath, Berl. Ent. Zeitschr. xxviii, p. 206 (1884); Distant, Rhop. Malayana, p. 423, Tab. xxxviii, fig. 7 ♂ (1882-1886).

1 ♂: Khao Bang-to, Pang-nga (c. 300 m.), P. Siam, 21. 2. 29 (*Kerr*).

Distribution: P. Siam; Malay Peninsula; Perak.

206. Zeuxidia amethystus Butl.

Zeuxidia amethystus Butler, Proc. Zool. Soc. Lond., p. 485 (1865).

Distribution: Burma to the Malay Peninsula, Sumatra, Borneo, Palawan and Mindano.

(a) **Zeuxidia amethystus masoni** Moore.

Zeuxidia amethystus masoni Fruhstorfer, in Seitz, *Macrolep.* ix, p. 435 (1911).

Me Song forest, Prae, N. Siam (*Godfrey*); Hot Spring, Me Lamoung, W. Siam (*Stockley*).

Distribution: Karen Hills to S. Burma; Siam.

(b) **Zeuxidia amethystus amethystus** Butl.

Zeuxidia amethystus amethystus Fruhstorfer, op. cit., p. 435.

Khao Ram and Khac Luang, Nakon Sritamarat (*Pendlebury*);
Khao Kalakiri, Pattani (*Kerr*)—P. Siam.

Distribution : P. Siam ; Malay Peninsula ; Sumatra.

207. *Zeuxidia aurelius aurelius* (Cram.).

Papilio aurelius Cramer, Pap. Exot. ii, p. 110 (1777).

Zeuxida aurelius aurelius Fruhstorfer, in Seitz, Macrolep. ix, p. 437
(1911).

Khao Ram, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula ; Sumatra.

A subspecies is found in Borneo.

208. *Zeuxidia doubledayi chersonesia* Fruhst.

Zeuxidia doubledayi Westwood, Gen. Di. Lep. ii, p. 329 (1848).

Zeuxidia doubledayi chersonesia Fruhstorfer, in Seitz, Macrolep. ix, p.
436 (1911).

Bangkok, C. Siam (*Godfrey*) ; Khao Bang-to, Pang-nga (c.
300 m.), P. Siam (*Kerr*).

Distribution : S. Burma to Perak.

The typical form occurs in Borneo and subspecies are found
in Sumatra and Banka.

209. *Thaumantis diores diores* Doubt.

Thaumantis diores Doubleday, Ann. Mag. N. H. xvi, p. 234 (1845)

Thaumantis diores diores Fruhstorfer, in Seitz, Macrolep. ix, p. 438,
fig. 104 a (1911).

Me Song forest, Prae, and Doi Chang (4,500 ft.), N. Siam
Sriracha and Khao Sebap, S. E. Siam (*Godfrey*) ; Me Wong (2,500 ft.)
W. Siam (*Stockley*).

Distribution : Sikkim to Burma ; Siam.

A subspecies is found in Hainan.

210. *Thaumantis lucipor* Westw.

Thaumantis lucipor Westwood, Gen. Diurn. Lep. ii, p. 337 (1851)
Fruhstorfer, in Seitz, Macrolep. ix, p. 438, fig. 104 a (1911).

Khao Ram, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : S. Burma ; Siam ; Malay Peninsula ; Sumatra ;
Borneo.

211. *Thaumantis odana pishuna* Fruhst.

Thaumantis odana Godart, Enc. Méth. Zool. ix, p. 445 (1823).

Thaumantis odana pishuna Fruhstorfer, in Seitz, Macrolep. ix, p. 440, fig. 105 b (1911).

Khao Ram and Khao Luang, Nakon Sritamarat (*Pendlebury*); Khao Kalakiri, Pattani (*Kerr*)—P. Siam.

Distribution: P. Siam; Malay Peninsula.

The typical form occurs in Java and subspecies are found in Sumatra and Nias.

212. *Thauria lathyi siamensis* Roths.

Thauria lathyi Fruhstorfer, Iris xv, p. 177 (1902).

Thauria aliris lathyi Fruhstorfer, in Seitz, Macrolep. ix, p. 441, fig. 102 b (1911).

Thauria lathyi siamensis Rothschild, Ann. Mag. N. H. xvii, No. 102, p. 474 (1916).

Hup Bon and Nong Khor, Sriracha forest, S. E. Siam.

Distribution: Rothschild (*l. c.*) treats *lathyi* and *aliris* as separate species, and restricts *lathyi siamensis* to Siam.

Typical *lathyi* occurs in Tonkin.

213. *Thauria aliris intermedia* Crowley.

Thauria aliris Westwood, Trans. Ent. Soc. Lond (2) iv, p. 76 (1858).

Thauria aliris intermedia Fruhstorfer, in Seitz, Macrolep. ix, p. 441 (1911).

1 ♂, 1 ♀: Me Wong (2,500 ft.), W. Siam, 13. 4. 20 (*Stockley*).

Distribution: N. Burma; Siam.

Th. a. aliris Westw. occurs in Borneo, and *Th. a. pseudaliris* Butl. in S. Burma and Perak.

214. *Discophora tullia zal* Westw.

Papilio tullia Cramer, Pap. Exot. i (1775).

Discophora tullia zal Fruhstorfer, in Seitz, Macrolep. ix, p. 443 (1911).

Me Song forest, Prae, N. Siam and Bangkok, C. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*); Klong Suan Mak, W. Siam (*Stockley*).

Distribution: Sikkim to Siam.

The species ranges from India to China and the Philippines, south to the Malay Peninsula, Sumatra, Borneo and Java.

215. **Discophora timora timora** Doubt.

Discophora timora Doubleday, Gen. Di. Lep. pl. 54, fig. 2 (1851).

Discophora continentalis continentalis Fruhstorfer, in Seitz, Macrolep. ix, p. 445 (1911).

Me Song forest, Prae, N. Siam; Bang Kwang, C. Siam.

Distribution: Sikkim to Burma; Siam.

Riley* regards *continentalis continentalis* as an absolute synonym of *timora timora*.

216. **Discophora deo deo** de Nicév.

Discophora deo deo de Nicéville, Journ. Bomb. N. H. S. xii, p. 137 (1998).

Discophora deo deo Fruhstorfer, in Seitz, Macrolep. ix, p. 444 (1911).

Me Song forest, Prae, N. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*).

Distribution: Assam to Siam.

Fruhstorfer (*l. c.*) recognizes a subspecies from Tonkin.

217. **Enispe euthymius euthymius f. tesselata** Moore.

Adolias euthymius Doubleday, Ann. Mag. N. H. xvi, p. 179 (1845).

Enispe euthymius euthymius f. tesselata Fruhstorfer, in Seitz, Macrolep. ix, p. 448 (1911).

3 ♂♂: Me Song forest, Prae, N. Siam, 2. 4. 18 (*Godfrey*);

1 ♂: Khao Luang (1,100 m.), P. Siam 30. 4. 28 (*Kerr*).

Distribution: Sikkim to Siam.

Subspecies are found in Borneo and Sumatra.

NYMPHALIDAE.

218. **Ergolis ariadne** (Johans.).

Papilio ariadne Johanssen, Amoen. Acad vi, p. 407 (1764).

Distribution: India and China to Celebes and Flores.

(a) **Ergolis ariadne ariadne** (Johans.).

Ergolis ariadne ariadne Fruhstorfer, in Seitz, Macrolep. ix, p. 456 (1911).

Tai Sai River, Nakon Sritamarat, P. Siam (Pendlebury); Pataling, P. Siam (*Kerr*).

Riley, *in litt.*, 30. 8. 24.

Distribution: P. Siam; Malay Peninsula; Borneo; Sumatra; Java.

(b) **Ergolis ariadne pallidior Fruhst.**

Ergolis ariadne pallidior Fruhstorfer, op. cit., p. 456.

Me Song forest, Prae, N. Siam; Pak Song, E. Siam.

Distribution: Mussoorie to Burma; Siam.

219. Ergolis merione pharis Fruhst.

Papilio merione Cramer, Pap. Exot. ii, p. 76 (1777).

Ergolis merione pharis Fruhstorfer, in Seitz, Macrolep. ix, p. 456 (1911)

Widely spread over Siam.

Distribution: S. Burma to the Karen; Siam; Annam.

The species is distributed from the Himalayas to Sumatra, Java and the Philippines.

The type of *pharis* came from Siam.

220. Ergolis specularia arca Fruhst.

Ergolis specularia Fruhstorfer, Berl. Ent. Zeit. 44, p. 91 (1900).

Ergolis specularia arca Fruhstorfer, in Seitz, Macrolep. ix, p. 458 (1911).

Pak Song, E. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam (*Godfrey*); Klong Pong Kapo, W. Siam (*Stockley*).

Distribution: Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

The typical form occurs in S. E. Borneo and Java.

221. Ergolis isaeus isaeus Wall.

Ergolis isaeus Wallace, Trans. Ent. Soc. Lond. p. 333 (1869).

Ergolis isaeus isaeus Fruhstorfer, in Seitz, Macrolep. ix, p. 459 (1911).

1 ♂: Me Song forest, Prae, N. Siam, April 1918.

Distribution: Burma; Siam; Malay Peninsula; W. Borneo; Sumatra.

Subspecies occur in Nias and Java.

This record is of interest. Fruhstorfer (*l. c.*) refers typical *isaeus* to the Malay Peninsula, Sumatra and W. Borneo and, in so doing, says:—"according to de Nicéville once found in Burma, but

this locality is not confirmed by Bingham and is doubtful, as de Nicéville was not able to distinguish the four Indian-Macrolayan species." The occurrence of *isaeus* in N. Siam would seem to indicate that de Nicéville was perfectly correct in both his identification and record.

222. *Laringa horsfieldi glaucescens* de Nicév.

Eurytela horsfieldi Boisduval, Faun. Ent. Mad. p. 54 (1833).

Laringa horsfieldi glaucescens, Fruhstorfer, in Seitz, Macrolep. ix, p. 459 (1912).

3 ♂♂ : Me Song forest, Prae, N. Siam, April 1916 (*Godfrey*) ;
1 ♂ : Chiengmai, N. Siam, December 1921 (*Tennent*).

Distribution : N. Shan States to Dawnas ; Siam.

The typical form occurs in Java and subspecies are known from the Andamans and Sumatra.

223. *Laringa castelnaui castelnaui* (Feld.).

Eurytela castelnaui Felder, Wien. ent. Monats. p. 401 (1860).

Laringa castelnaui castelnaui Fruhstorfer, in Seitz, Macrolep. ix, p. 406 (1912).

Khao Ram, Nakon Sritamarat (*Pendlebury*) ; Nang Yawn, Takuapa (*Kerr*)—P. Siam.

Distribution : S. Burma ; Siam ; Malay Peninsula.

Subspecies occur in Borneo, Sumatra and Java.

224. *Pseudergolis wedah* (Koll.).

Ariadne wedah Kollar, in Hügel's Kaschmir, iv (2), p. 437 (1844).

Pseudergolis wedah Fruhstorfer, in Seitz, Macrolep. ix, p. 461 (1912).

1 ♂ : Chieng Dao (1,500 m.), N. Siam, 7. 11. 22 (*Godfrey*) ; 1 ♂ : Chiengmai, N. Siam, Jan. 1921 (*Tennent*).

Distribution : Kashmir to S. China ; Siam.

225. *Penthema darlisa melema* Ril. & Godf.

Penthema darlisa Moore, Proc. Zool. Soc. Lond., p. 829 (1878) ; Fruhstorfer, in Seitz, Macrolep. ix, p. 464 (1912).

Penthema darlisa melema Riley and Godfrey, Journ. N. H. Soc. Siam, iv, p. 175, pl. v, fig. 1 (1921).

Me Song forest, Prae, N. Siam.

Distribution: This subspecies is confined to Siam.

The typical form occurs in Burma and a subspecies is known from S. Annam.

226. *Penthema binghami mimetica* Lathy.

Penthema binghami Wood-Mason, Journ. As. Soc. Beng. p. 87 (1881).

Penthema darlisa mimetica Fruhstorfer, in Seitz, Macrolep. ix, p. 464 (1912).

Penthema binghami mimetica Riley and Godfrey, Journ. Nat. Hist. Soc. Siam, iv, p. 175, pl. v, fig. 2 (1921).

Me Song forest, Prae, N. Siam; Pak Jong, E. Siam; Hup Bon, S. E. Siam.

Distribution: Siam.

Replaced in Burma by *P. b. binghami* W-M.

227. *Cupha erymanthis lotis* (Sulz.).

Papilio erymanthis Drury, Ill. Ex. Ent. i, pl. 15, figs. 3, 4 (1770).

Cupha erymanthis lotis Fruhstorfer, in Seitz, Macrolep. ix, p. 465 (1912).

Widely spread over Siam.

Distribution: Mussoorie to the Malay Peninsula; Borneo; Sumatra.

The typical form occurs in China and subspecies are known from Java and Palawan.

228. *Atella phalantha phalantha* (Drury).

Papilio phalantha Drury, Ill. Exot. Ent. i, pl. 21, figs. 1, 2 (1770).

Atella phalantha phalantha Fruhstorfer, in Seitz, Macrolep. ix, p. 471 (1912).

Widely spread over Siam.

Distribution: India; Burma; Tonkin; Siam; Annam.

Subspecies are found in Hainan and the Philippines.

229. *Atella alcippe alcippoides* Moore.

Papilio alcippe Cramer, Pap. Exot. iv, pl. 389, figs. G, H (1782).

Atella alcippe alcippoides Fruhstorfer, in Seitz, Macrolep. ix, p. 471 (1912).

Me Song forest, Prae, N. Siam; Sriracha, S. E. Siam (*Godfrey*); Nakon Sritamarat, P. Siam (*Pendlebury*); Kambengpet, W. Siam (*Stockley*).

Distribution: Sikkim to the Malay Peninsula; Borneo; Sumatra.

Other subspecies occur in Ceylon; Java, Palawan and the Moluccas.

230. *Issoria sinha sinha* (Koll.).

Terinos sinha Kollar, in Hügel's Kaschmir, iv (2), p. 438 (1844).

Issoria sinha sinha Fruhstorfer, in Seitz, Macrolep. ix, p. 473 (1912).

Widely spread over Siam.

Distribution: Sikkim to Siam.

The species ranges from India to New Guinea and the Solomon Islands.

231. *Cynthia erota* (Fabr.).

Papilio erota Fabricius, Ent. Syst. iii (1), p. 76 (1793).

Distribution: India to the Celebes and Philippines.

(a) ***Cynthia erota erota* (Fabr.).**

Cynthia erota erota Fruhstorfer, in Seitz, Macrolep. ix, p. 476, fig. 109 a (1912).

Common in all forest areas in Siam.

Distribution: Sikkim to Siam.

(b) ***Cynthia erota erotella* Butl.**

Cynthia erota erotella Fruhstorfer, in Seitz, Macrolep. ix, p. 477 (1912).

Khao Ram, Nakon Sritamarat (*Pendlebury*); Wat Kiriwong, Nakon Sritamarat (*Kerr*)—P. Siam.

Distribution: P. Siam; Malay Peninsula; Borneo; Sumatra; Java.

232. *Ducapa fasciata* (Feld.).

Atella fasciata Felder, Wien. ent. Monats. iv, p. 236 (1860).

Distribution: S. Burma to the Malay Peninsula, Borneo, Sumatra, Java, Palawan, and the Philippines.

(a) ***Ducapa fasciata fasciata* (Feld.).**

Cirrochroa fasciata fasciata Fruhstorfer, in Seitz, Macrolep. ix, p. 484 (1912).

Muak Lek, E. Siam and Bangkok, C. Siam (*Godfrey*) ; Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : S. Burma to the Malay Peninsula ; Sumatra.

- (b) **Ducapa fasciata fasciata f. flavobrunnea** (*Grose-Smith*).

Ducapa fasciata fasciata f. flavobrunnea Fruhstorfer, op. cit. p. 485.

This well-marked form has been obtained in Siam in the following localities :—Pak Jong, E. Siam (*Godfrey*) ; Me Wong, W. Siam (*Stockley*) ; Bangkok, C. Siam (*Aagaard*).

According to Fruhstorfer, it also occurs in Tenasserim.

233. Cirrochroa tyche (*Feld.*)

Atella tyche Feider, Wien. ent. Monats. v, p. 301 (1861).

Distribution : Sikkim to Sumatra, Borneo, Java, Palawan and the Philippines.

- (a) **Cirrochroa tyche mithila** Moore.

Cirrochroa tyche mithila Fruhstorfer, in Seitz, Macrolep. ix, p. 486 (1912).

Widely spread over Siam proper.

Distribution : Sikkim to Siam ; Tonkin.

- (b) **Cirrochroa tyche rotundata** Butl.

Cirrochroa tyche rotundata Fruhstorfer, op. cit., p. 487.

Khao Luang and Wat Kiriwong, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula.

234. Cirrochroa aoris olivacea de Nicév.

Cirrochroa aoris Doubleday, Gen. Di. Lep. i, p. 157 (1848).

Cirrochroa aoris olivacea Fruhstorfer, in Seitz, Macrolep. ix, p. 488 (1912).

3 ♂ ♂ : Me Song forest, Prae, N. Siam, April 1918.

Distribution : N. Burma to N. Siam.

The typical form occurs from Sikkim to Assam.

235. Cirrochroa surya siamensis Fruhst.

Cirrochroa surya Moore, Proc. Zool. Soc. Lond., p. 827 (1878).

Cirrochroa surya siamensis Fruhstorfer, in Seitz, Macrolep. ix, p. 489, fig. 107 g (1912).

Bangkok, C. Siam.

Distribution: Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

The typical form occurs from S. Burma to the Dawnas.

236. *Cirrochroa emalea martini* Fruhst.

Cirrochroa emalea Guérin, Del. Souv. Ind. p. 72 (1843).

Cirrochroa emalea martini Fruhstorfer, in Seitz, Macrolep. ix, p. 489 (1912).

Tai Sai River, Khao Ram and Khao Luang, P. Siam (*Pendlebury*).

Distribution: Mergui; P. Siam; Malay Peninsula; Sumatra. Other subspecies are found in Borneo, Java and Nias.

237. *Cirrochroa orissa orissa* Feld.

Cirrochroa orissa Felder, Wien. ent. Monats. iv, p. 399 (1860).

Cirrochroa orissa orissa Fruhstorfer, in Seitz, Macrolep. ix, p. 491, fig. 1079 (1912).

Bangnara (*Godfrey*); Khao Ram and Khao Luang, Nakon Sritamarat (*Pendlebury*); Ban Ton, Satul (*Kerr*)—all in P. Siam.

Distribution: S. Burma to the Malay Peninsula; Sumatra. A subspecies occurs in N. Borneo and Sarawak.

238. *Cirrochroa chione* Ril. & Godf.

Cirrochroa chione Riley and Godfrey, Journ. Nat. Hist. Soc. Siam, iv, pl. 5, fig. 3 (1921).

2 ♂♂: Khao Sebap, Chantabun, S. E. Siam, March 1916.

Distribution: Known only from Siam.

239. *Terinos terpander* Hew.

Terinos terpander Hewitson, Proc. Zool. Soc. Lond., p. 90 (1862).

Distribution: Siam; Malay Peninsula; Borneo; Sumatra; Nias; Java.

(a) ***Terinos terpander intermedia* Godf.**

Terinos terpander intermedia Godfrey, Journ. N. H. Soc. Siam ii, p. 126 (1916); Riley and Godfrey, iv, pl. 7, figs. 2, 3, ♂, ♀ (1921).

Distribution: This subspecies is known only from Siam.

(b) ***Terinos terpander robertsia* Butl.**

Terinos terpander robertsia Fruhstorfer, in Seitz, Macrolep. ix, p. 493, fig. 108 c (1912).

Khao Ram and Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*) ; Khao Kalakiri, P. Siam (*Kerr*).

Distribution : P. Siam ; Malay Peninsula.

240. *Terinos clarissa* Boisd.

Terinos clarissa Boisduval, Spéc. Gén. Lép. i, Taf. 9. (1836).

Distribution : S. Burma to the Malay Peninsula ; Sumatra ; Borneo ; Java ; Palawan ; Philippines.

(a) *Terinos clarissa falcata* Fruhst.

Terinos clarissa falcata Fruhstorfer, in Seitz, Macrolep. ix, p. 495 (1912).

Hinlap, E. Siam (*Fruhstorfer*) ; Muak Lek, E. Siam and Pechaburi River, W. Siam (*Godfrey*).

Distribution : This subspecies is confined to E. and W. Siam.

(b) *Terinos clarissa clarissa* Boisd.

Terinos clarissa clarissa Fruhstorfer, op. cit., p. 494.

Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula.

241. *Cethosia biblis perakana* Fruhst.

Papilio biblis Drury, Ill. Exot. Ent. i, pl. 4, fig. 2 (1770).

Cethosia biblis perakana Fruhstorfer, in Seitz, Macrolep. ix, p. 499, pl. 110 c (1912).

Widely spread over Siam.

Distribution : Siam ; Malay Peninsula.

The species extends from India to the Moluccas and in a northerly direction from the Philippines to Hainan and Hongkong.

242. *Cethosia cyane euanthes* Fruhst.

Papilio cyane Drury, Ill. Exot. Ent. i, pl. 4, fig. 1 (1770).

Cethosia cyane euanthes Fruhstorfer, in Seitz, Macrolep. ix, p. 503 (1912).

Muak Lek, E. Siam ; Sriracha forest, S. E. Siam ; Bangkok, C. Siam (*Godfrey*) ; Me Wong, W. Siam (*Stockley*).

Distribution : Tonkin ; Siam ; Annam.

The typical form occurs from Mussoorie to Burma.

243. *Cethosia methypsea methypsea* Butl.

Cethosia methypsea Butler, Trans. Linn. Soc. Lond., p. 443 (1877).

Cethosia methypsea methypsea Fruhstorfer, in Seitz, Macrolep. ix, p. 506 (1912).

Sriracha, S. E. Siam (*Godfrey*); Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*); Satul, Patalung and Pulo Adang, P. Siam (*Kerr*).

Distribution: Siam; Malay Peninsula.

A subspecies occurs in Sumatra.

244. *Argynnis hyperbius hyperbius* (Johans.).

Papilio hyperbius Johanssen, Amoen. Acad. vi, p. 408 (1764); Fruhstorfer (*Argynnis*), in Seitz, Macrolep. ix, p. 515 (1912).

1 ♂, 2 ♀ ♀: Doi Sutep (5,000 ft.), N. Siam, March 1921 (*Tennent*); 1 ♂: Doi Chang (4,500 ft.), N. Siam, April 1921 (*Godfrey*).

Distribution: Chitral to N. Burma; China; Tonkin; Siam.

According to Fruhstorfer, the species ranges from Abyssinia to Australia, and from S. Japan to the Malay Peninsula.

245. *Precis iphita iphita* (Cram.).

Papilio iphita Cramer, Pap. Exot. iii, pl. 209, figs. C. D (1779).

Distribution: C. China; Formosa; Sikkim to the Malay Peninsula.

(a) ***Precis iphita iphita* (Cram.).**

Precis iphita iphita Fruhstorfer, in Seitz, Macrolep. ix, p. 517 (1912).

Widely spread over Siam proper.

Distribution: Sikkim to Tonkin and Hainan; Siam.

(b) ***Precis iphita horsfieldi* Moore.**

Precis iphita horsfieldi Fruhstorfer, op. cit., p. 516.

Nakon Sritamarat, P. Siam (*Pendlebury*; *Kerr*).

Distribution: P. Siam; Malay Peninsula; Borneo; Sumatra; Java.

246. *Precis atlites atlites* (Johans.).

Papilio atlites Johanssen, Centur. Ins., p. 24 (1763).

Precis atlites atlites Fruhstorfer, in Seitz, Macrolep. ix, p. 519, fig. 117 a (1912).

Widely spread over Siam.

Distribution: India; China; Siam; Annam; Malaya.

Fruhstorfer (*l. c.*) recognizes a subspecies from the Celebes.

247. **Precis almana almana** (Linn.).

Precis almana Linnaeus, Syst. Nat. x, p. 472 (1758).

Precis almana almana Fruhstorfer, in Seitz, Macrolep. ix, p. 519 (1912).

Widely spread over Siam.

Distribution: Ceylon; India; China; Tonkin; Hainan; Siam; Annam.

The species extends to the Malayan subregion.

248. **Precis lemonias** (Linn.).

Papilio lemonias Linnaeus, Syst. Nat. x, p. 473 (1758); Fruhstorfer (*Precis*) in Seitz, Macrolep. ix, p. 520 (1912).

Widely spread over Siam.

Distribution: Throughout the Indo-Australian region.

249. **Precis orithya** (Linn.).

Papilio orithya Linnaeus, Syst. Nat. x, p. 473 (1758).

Distribution: India; China; Hainan; Siam; Malaya; Australia.

(a) **Precis orithya ocyale** (Hübner).

Precis orithya ocyale Fruhstorfer, in Seitz, Macrolep. ix, p. 522 (1912).

Me Song forest, Prae, N. Siam; Muak Lek, E. Siam; Sriracha, S. E. Siam (*Godfrey*); Kambengpet, W. Siam (*Stockley*).

Distribution: Sikkim to Siam and Annam.

(b) **Precis orithya wallacei** (Distant).

Junonia wallacei Distant, Rhop. Malay., p. 95, Tab. xi, figs. 3, 4.

Precis orithya wallacei Fruhstorfer, op. cit., p. 523.

Padang Besar, P. Siam (*Pendlebury*).

Distribution: P. Siam; Malay Peninsula.

250. **Precis hierta hierta** (Fabricius).

Papilio hierta Fabricius, Ent. Syst. Suppl. p. 424 (1798); Fruhstorfer (*Precis*), in Seitz, Macrolep. ix, p. 524 (1912); Evans (*Precis*), Journ. Bomb. N. H. S. xxx (1), p. 84 (1924).

Widely spread over Siam.

Distribution: Ceylon; India; Burma; China; Hainan; Siam.

Evans (*l. c.*) recognizes a subspecies, *P. h. oenone* Cr., from Sikkim and Burma.

251. *Vanessa indica indica* (Herbst).

Papilio atalanta indica Herbst, Natursyst. Schmett. vii, p. 171 (1794).

Pyrameis indica indica Fruhstorfer, in Seitz, Macrolep. ix, p. 525, (1912); Evans (*Vanessa*), Journ. Bomb. N.H.S. xxx (1), p. 85 (1924).

1 ♂ : Khun Tan (4,000 ft.), N. Siam, March 1916 (*Godfrey*);
1 ♂ : Doi Sutep (4,500 ft.), N. Siam, April 1921 (*Tennent*).

Distribution : Kashmir to N. Burma; China; Hainan; Siam.

The species is distributed from India and China to the Philippines and Celebes.

252. *Vanessa canace canace* (Johans.).

Papilio canace Johanssen, Centur. Ins. p. 23 (1763).

Vanessa canace canace Fruhstorfer, in Seitz, Macrolep. ix, p. 528 (1912).

1 ♂ : Chiengrai, N. Siam, April 1921.

Distribution : Sikkim to N. Burma; Siam.

The species ranges from India, China and Japan to the Malay Peninsula and Archipelago.

253. *Symbrenthia hippoclus draca* Moore.

Symbrenthia hippoclus de Nicéville, Butt. Ind. ii, p. 240 (1886).

Symbrenthia hippoclus daruca Fruhstorfer, in Seitz, Macrolep. ix, p. 529 (1912).

Me Song forest, Prae, N. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*); Me Lamoung (2,000 ft.), W. Siam (*Stockley*).

Distribution : Sikkim to Siam and Annam.

The species ranges from India to China, Tonkin and Hainan, south through Malaya to New Guinea.

254. *Symbrenthia hypselis sinis* de Nicév.

Vanessa hypselis Godart, Encycl. Méth. ix, Suppl., p. 818 (1823).

Symbrenthia hypselis sinis Fruhstorfer, in Seitz, Macrolep. ix, p. 532 (1912).

Me Song forest, Prae, N. Siam (*Godfrey*); Me Lamoung (2,000 ft.), W. Siam. (*Stockley*).

Distribution : Fruhstorfer (*l. c.*) refers this subspecies to the Karen Hills, Tenasserim and the Malay Peninsula.

The species ranges from India to Sumatra, Borneo, Nias, Java and Palawan.

255. *Symbrenthia hypatia chersonesia* Fruhst.

Laogona hypatia Wallace, Trans. Ent. Soc. Lond., p. 343 (1869).

Symbrenthia hypatia chersonesia Fruhstorfer, in Seitz, Macrolep. ix, p. 534 (1912).

1 ♀ : Kao Kalakiri, Pattani, P. Siam, 1. 4. 28 (*Kerr*).

Distribution : P. Siam ; Malay Peninsula ; Sumatra.

The typical form occurs in Java and a subspecies is found in Borneo.

256. *Rhinopalpa polynice eudoxia* (Guér.).

Papilio polynice Cramer, Pap. Exot. iii, pl. 195, figs. D, E (1780).

Rhinopalpa polynice eudoxia Fruhstorfer, in Seitz, Macrolep. ix, p. 538 (1912).

1 ♂ : Wat Kiriwong, Nakon Sritamarat, P. Siam, 3. 5. 28 (*Kerr*).

Distribution : P. Siam ; Malay Peninsula.

The species ranges from Assam to Sumatra, Java, the Philippines and the Celebes.

257. *Yoma sabina vasuki* Doh.

Yoma sabina Cramer, Pap. Exot. iv, Taf. 289 (1782).

Yoma vasuki Doherty, Journ. As. Soc. Beng., p. 259 (1886.).

Yoma sabina vasuki Fruhstorfer, in Seitz, Macrolep. ix, p. 540. fig. 113 b (1912).

Me Song forest, Prae, N. Siam and Sriracha forest, S. E. Siam (*Godfrey*) ; Chiengmai, N. Siam (*Tennient*) ; Huey Tuk Sao, W. Siam (*Stockley*).

Distribution : Burma ; Siam.

The species ranges from Burma to the Moluccas, south to Australia.

258. *Hypolimnas misippus* (Linn.).

Papilio misippus Linnaeus, Mus. Ulr. p. 264 (1764) ; Fruhstorfer (*Hypolimnas*), in Seitz, Macrolep. ix, p. 547 (1912).

Bankok C. Siam.

Distribution : Throughout the Indo-Australian region.

259. **Hypolimnas bolina bolina** (Linn.).

Papilio bolina Linnaeus, Syst. Nat. x, p. 479 (1758).

Hypolimnas bolina bolina Fruhstorfer, in Seitz, Macrolep. ix, p. 547 (1912).

Widely spread over Siam.

Distribution : India to the Malay Peninsula; Sumatra; Borneo; Java; Philippines.

Fruhstorfer divides up this species into a large number of different races and forms ranging from India to New Guinea and Australia.

260. **Doleschallia bisaltide siamensis** Fruhst.

Papilio bisaltide Cramer, Pap. Exot. ii, pl. 102 (1779).

Doleschallia bisaltide siamensis Fruhstorfer, in Seitz, Macrolep. ix, p. 557 (1912).

Me Song forest, Prae, N. Siam; Pak Jong, E. Siam; Sriracha, S. E. Siam.

Distribution : Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

The species ranges from India through Malaya to the Bismarck Archipelago.

261. **Kallima inachus** (Boisd.).

Paphia inachus Boisduval, in Crochard's ed. Cuv. Règ. Anim. Ins. ii, pl. 139, fig. 3 (1836).

Distribution : N. India to China, southwards to the Malay Peninsula and the Greater Sunda Islands.

(a) **Kallima inachus siamensis** Fruhst.

Kallima inachus siamensis Fruhstorfer, in Seitz, Macrolep. ix, p. 566 (1912).

Me Song forest, Prae, N. Siam; Muak Lek and Pak Jong, E. Siam; Sriracha forest, S. E. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*).

Distribution : Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

(b) **Kallima inachus limborgi** Moore.

Kallima limborgii Moore, Proc. Zool. Soc. Lond., p. 828 (1878).

2 ♂♂ : Wat Kiriwong, Nakon Sritamarat, P. Siam, 3. 5. 28 (*Kerr*).

Fruhstorfer treats *limborgi* as a separate species; Riley* regards it as a subspecies of *inachus*.

262. *Cyrestis periander periander* (Fabr.).

Papilio periander Fabricius, Mant. Ins. ii, p. 9 (1787).

Cyrestis periander periander et siamensis Fruhstorfer, in Seitz, Macrolep. ix, p. 573 (1912).

Me Song forest, Prae, N. Siam ; Muak Lek and Pak Jong, E. Siam ; Pechaburi River, W. Siam (*Godfrey*) ; Pulo Adang, P. Siam (*Kerr*).

Distribution : Shan States ; Siam ; Malay Peninsula.

The species ranges from Burma to Sumatra and Java, but has not yet been found in Nias or Borneo.

C. p. siamensis Fruhst., a very large and dark form described by its author from two specimens one of which is in the British Museum, cannot be referred to Siam. From an entry in Register 56.113 in the British Museum, it may possibly have come from Pulo Aor, an island off the south-east coast of the Malay Peninsula.

263. *Cyrestis cocles cocles* (Fabr.).

Papilio cocles Fabricius, Mant. Ins. ii, p. 7 (1787).

Cyrestis cocles cocles Fruhstorfer, in Seitz, Macrolep. ix, p. 574, fig. 122 b (1912).

Me Song forest, Prae, and Chiengrai, N. Siam ; Muak Lek, E. Siam ; Pechaburi River, W. Siam (*Godfrey*) ; Kaw Tao, P. Siam (*Kerr*).

Distribution : Sikkim to Burma ; Hainan ; Siam.

This species, which was originally described from Siam, ranges from Sikkim to Borneo.

264. *Cyrestis nivea nivalis* Feld.

Amathusia nivea Zinken-Sommer, Nova Acta Acad., p. 138, pl. 14, fig. 1 (1831).

Cyrestis nivea nivalis Fruhstorfer, in Seitz, Macrolep. ix, p. 576 (1912).

*Riley, *in litt.*, 24. 10. 29.

Bangnara, P. Siam (*Godfrey*) ; Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : S. Burma ; Siam ; Malay Peninsula ; Borneo ; Sumatra.

The typical form occurs in Java and subspecies are known from Hainan and the Philippines.

265. *Cyrestis thyodamas thyodamas* Boisd.

Cyrestis thyodamas Boisduval, Cuv. Règ. Anim. Ins. ii, pl. 138, fig. 4 (1836).

Cyrestis thyodamas thyodamas Fruhstorfer, in Seitz, Macrolep. ix, p. 581 (1912).

Common in all forest areas in Siam.

Distribution : Sikkim to Burma ; Siam ; Yunnan ; Annam ; Tonkin.

Subspecies are known from China, Japan, the lesser Sunda Islands, and New Guinea.

266. *Chersonesia risa risa* (Doubt).

Cyrestis risa Doubleday, Gen. Di. Lep. ii, p. 262 (1850).

Cyrestis risa risa Fruhstorfer, in Seitz, Macrolep. ix, p. 592, fig. 121 f (1912).

Me Song forest, Prae, N. Siam ; Pak Jong, E. Siam ; Sriracha forest, S. E. Siam (*Godfrey*), Eka Klo, W. Siam (*Stockley*).

Distribution : Kumaon to Burma ; Tonkin ; Hainan ; Siam ; Annam.

Subspecies are known from Borneo and Sumatra.

267. *Chersonesia rahria rahrioides* Moore.

Cyrestis rahria Moore, Cat. Lep. Mus. E. I. C., p. 147 (1857).

Chersonesia rahrioides Moore, Lep. Ind. iv, p. 56 (1899-1900) ; Fruhstorfer, in Seitz, Macrolep. ix, p. 593 (1912).

Chersonesia rahria rahrioides Evans, Journ. Bomb. N. H. S. xxx (1), p. 82 (1924).

Pak Jong, E. Siam.

Distribution : Assam to Burma ; Tonkin ; Siam.

The species ranges from the Himalayas to Sumatra, Nias. Borneo and Java.

268. Chersonesia peraka Dist.

Chersonesia peraka Distant, Ann. Mag. N. H., p. 199 (1884); Fruhstorfer, in Seitz, Macrolep. ix, p. 594, fig. 121 f (1912).

Khao Luang, Nakon Sritamarat, P. Siam (*Penilebury*). Also recorded by Fruhstorfer from "Lower Siam"

Distribution: S. Burma to the Malay Peninsula, Sumatra, Nias, Borneo and Java.

269. Rahinda hordonia (Stoll).

Papilio hordonia Stoll, Pap. Exot. Suppl. pl. 33, figs. 4, 4d (1791).

Rahinda hordonia hordonia Fruhstorfer in Seitz, Macrolep. ix, p. 597 (1912).

Me Song forest, Prae, N. Siam; Muak Lek, E. Siam; Sriracha, S. E. Siam.

Distribution: Burma; Siam; Malay Peninsula.

The species ranges from India to Hainan and Formosa; south to Borneo, Sumatra, Nias, Java, Bali, and Sumbawa.

270. Rahinda paraka (Butl.).

Neptis paraka Butler, Trans. Linn. Soc. ser. 2, Zool. i, p. 452, pl. 68, fig. 2 (1879).

Rahinda paraka assamica Fruhstorfer, in Seitz, Macrolep. ix, p. 598 (1912).

Neptis dindinga assamica Evans, Journ. Bomb. N. H. S. xxx, p. 81 (1924).

Pak Jong and Muak Lek, E. Siam; Bangnara, P. Siam.

Distribution: Assam to the Malay Peninsula; Borneo; Sumatra; Java; Banka; Palawan.

Other subspecies are found in Nias and Mentawai. Fruhstorfer (*l. c.*) gives *R. p. assamica* Moore as the Siamese race, but specimens obtained by me in Siam have been identified at the British Museum as typical *paraka*.

Evans (*l. c.*) treats *assamica* as a very rare subspecies of *N. dindinga* Butl.

271. Rahinda aurelia Staud.

Rahinda aurelia Staudinger, Exot. Schmett. p. 145 (1886); Fruhstorfer, in Seitz, Macrolep. ix, p. 599 (1912).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Assam to the Malay Peninsula; Sumatra; Borneo.

272. **Neptis hylas** (Linn.).

Papilio hylas Linnaeus, Syst. Nat. i, p. 486 (1758).

Distribution: Throughout the Indo-Australian region.

(a) **Neptis hylas hylas** (Linn.).

Neptis hylas hylas Fruhstorfer, in Seitz, Macrolep. ix, p. 601 (1912).

Widely spread over Siam proper.

Distribution: China; Tonkin; Siam; Annam.

(b) **Neptis hylas mamaea** Butl.

Neptis hylas mamaea Fruhstorfer, op. cit., p. 602.

Nakon Sritamarat, P. Siam (*Pendlebury*; *Kerr*); Bangnara, P. Siam (*Godfrey*).

Distribution: P. Siam; Malay Peninsula.

273. **Neptis magadha magadha** Feld.

Neptis magadha Felder, Nov. Reise, Lep. Rhop. p. 427 (1867).

Neptis magadha magadha Fruhstorfer, in Seitz, Macrolep. ix, p. 604 (1912).

1 ♂ : Klong Klung, 1. 3. 24; 1 ♂ : Me Wong, 19. 2. 24; 1 ♂ : Hue Tak So, 24. 2. 24 (*Stockley*)—all in W. Siam.

Distribution: S. Burma; Siam.

The species ranges from Bhutan to the Malay Peninsula, Sumatra, Borneo and Java.

274. **Neptis duryodana nesia** Fruhst.

Neptis duryodana Moore, Proc. Zool. Soc. Lond., p. 10 (1858).

Neptis duryodana nesia Fruhstorfer, in Seitz, Macrolep. ix, p. 604 (1912).

Khao Ram, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: P. Siam; Malay Peninsula; Sumatra.

The typical form occurs in Burma and subspecies are known from Nias and Java.

275. **Neptis nandina** Moore.

Neptis nandina Moore, Cat. Lep. Mus. E. I. C., p. 168 (1857).

Distribution: India to Formosa and the Philippines; south through the Malay Peninsula and Archipelago to Lombok and Flores.

(a) **Neptis nandina gonatina** Fruhst.

Neptis nandina gonatina Fruhstorfer, in Seitz, Macrolep ix, p. 606 (1912).

Muak Lek and Pak Jong, E. Siam.

Distribution: Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

(b) **Neptis nandina susrutina** Fruhst.

Neptis nandina susrutina Fruhstorfer, op. cit., p. 606.

Me Wong, W. Siam (*Stockley*).

Distribution: Burma and W. Siam.

276. Neptis soma tushita Fruhst.

Neptis soma Moore, Proc. Zool. Soc. Lond., p. 9 (1858).

Neptis soma tushita Fruhstorfer, in Seitz, Macrolep. ix, p. 607 (1912).

Muak Lek and Pak Jong, E. Siam ; Sriracha, S. E. Siam.

Distribution: Fruhstorfer (*l. c.*) restricts this subspecies to Siam, and states that the species is " confined to the Continent and nowhere found in the Northern Philippines or in the Macro-malayan Islands, Engano and Perak excepted."

277. Neptis yerburii shania Evans.

Neptis yerburii Butler, Proc. Zool. Soc. Lond., p. 360 (1886); Fruhstorfer, in Seitz, Macrolep. ix, p. 608 (1912).

Neptis yerburii shania Evans, Journ. Bomb. N. H. S. xxx, p. 78 (1924).

1 ♂ : Hot Spring (3,000 ft.), W. Siam, 28.1.24 (*Stockley*).

Distribution: N. Burma ; Siam.

The typical form occurs from Kashmir to Kumaon, and subspecies are found in Sikkim, in China, and in Sumatra.

278. Neptis zaida Westw.

Neptis zaida Westwood, Gen. Di. Lep. p. 272 (1850); Moore, Lep. Ind. iv, p. 20 (1899-1900); Fruhstorfer, in Seitz, Macrolep. ix, p. 609 (1912).

Moore (*l. c.*) records a male of the wet-season form with pure ochraceous markings, labelled "Siam," from the Grosse-Smith collection.

Distribution: Murree to the Shan States ; Siam.

279. *Neptis heliodore* (Fabr.).

Papilio heliodore Fabricius, Ent. Syst. iii, p. 130 (1793).

Distribution: This species, which was described from Siam, ranges from Assam to the Malay Peninsula and Archipelago.

(a) *Neptis heliodore heliodore* (Fabr.).

Neptis heliodore heliodore Fruhstorfer, in Seitz, Macrolep. ix, p. 610 (1912).

Muak Lek, E. Siam (*Fruhstorfer*) ; Me Song forest, Prae, N Siam (*Godfrey*).

Distribution: Burma ; Siam.

(b) *Neptis heliodore dorelia* Butl.

Neptis heliodore dorelia Fruhstorfer, op. cit., p. 611.

Khao Ram, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: P. Siam ; Malay Peninsula, Borneo.

280. *Neptis harita* Moore.

Neptis harita Moore, Proc. Zool. Soc. Lond. p. 571, pl. 66, fig. 8 (1874) ; Evans, Journ. Bomb. N. H. S. xxx (1), p. 78 (1924).

Neptis vikasi harita Fruhstorfer, in Seitz, Macrolep. ix, p. 611 (1912).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Bengal to Burma ; Siam ; Malay Peninsula. Fruhstorfer (*l. c.*) treats *harita* as a subspecies of *vikasi* Horsf., but both Riley* and Evans (*l. c.*) regard it as a separate species.

281. *Neptis vikasi sakala* Fruhst.

Neptis vikasi Horsfield, Cat. Lep. Mus. E. I. C., pl. 5, figs. 2, 2a (1829).

Neptis vikasi sakala Fruhstorfer, in Seitz, Macrolep. ix, p. 611 (1912).

2 ♂♂ : Me Song forest, Prae, N. Siam, April 1918.

Distribution: Tonkin ; Siam.

The species ranges from India to Tonkin, Palawan and the Philippines ; and southwards to Borneo, the Celebes, Sumatra and Java.

* Riley, *in litt.*, *13. 12. 24.

282. *Neptis cartica* meraka Ril. & Godf.

Neptis cartica Moore, Proc. Zool. Soc. Lond. p. 562 (1872); Fruhstorfer, in Seitz, Macrolep. ix, p. 613 (1912).

Neptis cartica meraka Riley and Godfrey, Journ. N. H. S. Siam, iv, p. 177, pl. 5, fig. 4 (1921).

Me Song forest, Prae, N. Siam.

Distribution: Known only from Siam.

The typical form occurs from Sikkim to Assam and a subspecies is known from Burma.

283. *Neptis columella* (Cram.).

Papilio columella Cramer, Pap. Exot. iv, pl. 296, figs. A, B. (1782); Fruhstorfer (*Neptis*), in Seitz, Macrolep. ix, p. 615 (1912).

Distribution: According to Fruhstorfer, this species "is represented by a long series of local and insular forms ranging over the entire Oriental Region as well as a large number of Maero and Micromalayan Islands."

(a) *Neptis columella martabana* f. *martabana* Moore.

Neptis columella martabana f. *martabana* Fruhstorfer, op. cit., p. 615.

Chiengmai, N. Siam (*Tennent*); Muak Lek, E. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam (*Godfrey*); Chumporn, P. Siam (*Aagaard*), Pan Yao, W. Siam (*Stockley*).

Distribution: S. Burma; Siam; Malay Peninsula; Annam.

(b) *Neptis columella martabana* f. *alesia* Fruhst.

Neptis columella martabana f. *alesia*, Fruhstorfer, op. cit., p. 615.

Nam Pat, N. E. Siam (*Godfrey*); Pan Yao, W. Siam (*Stockley*).

Distribution: Siam; Malay Peninsula; S. Annam.

284. *Neptis sankara* quiltä Swinh.

Liminitis sankara Kollar, in Hügel's Kaschmir iv (2) p. 428 (1844).

Neptis sankara quiltä Fruhstorfer, in Seitz, Macrolep. ix, p. 619 (1912).

1 ♂ : Chiengmai, N. Siam, Dec. 1920 (*Tennent*).

Distribution: Sikkim to Burma; Siam.

The typical form occurs from Kashmir to Kumaon and subspecies are found in the Andamans, Yunnan, and Sumatra.

285. **Neptis viraja viraja** Moore.

Neptis viraja Moore, Proc. Zool. Soc. Lond. p. 563 (1872); Frühstorfer, in Seitz, Macrolep. ix, p. 620, fig. 125 c. (1912).

Neptis viraja viraja Evans, J. Bomb. N. H. S. xxx, p. 80 (1924).

1 ♂ : Umpang, W. Siam, 23. 4. 21 (*Stockley*); 1 ♂ : Chiengsen, N. Siam, 20. 4. 21 (*Godfrey*).

Distribution: Kumaon to Burma; Siam.

The race *N. v. kanara* Evans (S. India-Orissa) is not included in Seitz' "Macrolepidoptera."

286. **Neptis miah nolana** Druce.

Neptis miah Moore, Cat. Lep. Mus. E. I. C., p. 164 (1857).

Neptis nolana Druce, P. Z. S., p. 105, ♀ (1874).

Neptis miah nolana Frühstorfer, in Seitz, Macrolep. ix, p. 621, pl. 125 c. (1912).

Pak Jong and Muak Lek, E. Siam.

Distribution: Burma; Tonkin; Hainan; Siam; Annam.

The species ranges from India and China to the Malay Peninsula, Sumatra, Borneo and Java.

N. nolana was originally described from Siam.

287. **Neptis ebusa fuliginosa** Moore.

Neptis ebusa Felder, Wien. ent. Monats. vii, p. 112. (1861).

Neptis fuliginosa fuliginosa Frühstorfer, in Seitz, Macrolep. ix, p. 622 (1912).

Neptis ebusa fuliginosa Evans, Journ. Bomb. N. H. S. xxx (1), p. 79 (1924).

Pak Jong, E. Siam.

Distribution: S. Burma; Siam; Malay Peninsula.

Typical *ebusa* occurs in the Philippines.

Frühstorfer (*l. c.*) treats *fuliginosa* as a separate species, but remarks:—"I consider it highly probable that *fuliginosa* Moore represents only the Indo-Malayan branch of *ebusa*, but I lack sufficient material to decide this question."

288. **Pantoporia pravara indosinica** Frühst.

Athydas pravara Moore, Cat. Lep. Mus. E. I. C., p. 173 (1857).

Pantoporia pravara indosinica Frühstorfer, in Seitz, Macrolep. ix p. 624 (1912).

Muak Lek (*Fruhstorfer*) ; Pak Jong (*Godfrey*)—E. Siam.

Distribution: Tonkin ; Siam.

The species ranges from Assam to the Malay Peninsula, Sumatra, Borneo, Java, and Palawan.

289. *Pantoporia perius* (Linn.).

Papilio perius Linnaeus, Syst. Nat. x, p. 471 (1758) ; Fruhstorfer (*Pantoporia*), in Seitz, Macrolep. ix, p. 626 (1912).

Widely spread over Siam.

Distribution: India to China and Formosa, south to Sumatra and Sumbawa.

290. *Pantoporia larymna siamensis* Fruhst.

Liminitis larymna Doubleday, Gen. Di. Lep. ii, pl. 35, fig. 1 (1850).

Pantoporia larymna siamensis Fruhstorfer, in Seitz, Macrolep. ix, p. 627 (1912).

Me Song forest, Prae, and Chiengrai, N. Siam ; Pak Jong and Muak Lek, E. Siam.

Distribution: Assam to the Malay Peninsula.

Other subspecies are found in Palawan, Borneo, Sumatra, Nias, and Java.

291. *Pantoporia kanwa kanwa* (Moore).

Athyyna kanwa Moore, P. Z. S. Lond., p. 17, pl. 51, fig. 2 (1858).

Pantoporia kanwa kanwa Fruhstorfer, in Seitz, Macrolep. ix, p. 628 (1912).

Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: P. Siam ; Malay Peninsula ; Singapore ; Borneo.

Other subspecies are found in Assam and Burma ; in Sumatra ; and in Nias.

292. *Pantoporia reta moorei* Fruhst.

Pantoporia reta Moore, Proc. Zool. Soc. Lond., p. 12 (1858).

Pantoporia reta moorei Fruhstorfer, in Seitz, Macrolep. ix, p. 629 (1912).

1 ♀ : Bangnara, P. Siam, 24. 9. 23.

Distribution: P. Siam ; Malay Peninsula ; Singapore.

The species ranges from Assam to Borneo, Sumatra, and Nias.

293. **Pantoporia ranga obsolescens** Fruhst.

Athyma ranga Moore, Cat. Lep. Mus. E. I. C., p. 175 (1857).

Pantoporia ranga obsolescens Fruhstorfer, in Seitz, Macrolep. ix, p. 629 (1912).

Me Song forest, Prae, N. Siam ; Pak Jong, E. Siam.

Distribution : Tonkin ; Siam.

The typical form occurs from Sikkim to Burma, and subspecies are known from S. India and from W. China.

294. **Pantoporia abiasa clerica** Butl.

Athyma abiasa Moore, Proc. Zool. Soc. Lond., p. 16 (1858).

Pantoporia abiasa clerica Fruhstorfer, in Seitz, Macrolep. ix, p. 630 (1912).

Khao Luang (5,800 ft.), Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : S. Burma ; P. Siam ; Malay Peninsula ; Sumatra.

The typical form occurs in Java, and subspecies are known from Borneo and Nias.

295. **Pantoporia opalina orientalis** Elw.

Liminitis opalina Kollar, in Hügel's Kaschmir, iv, pt. 2, p. 427 (1844).

Pantoporia opalina orientalis Fruhstorfer, in Seitz, Macrolep. ix, p. 630 fig. 124 c. (1912).

1 ♂ : Chiangmai district, N. Siam (*Tennent*).

Distribution : Sikkim to N. Siam.

The species ranges from Kashmir to China, south to Siam and Annam.

296. **Pantoporia selenophora** (Koll.).

Liminitis selenophora Kollar, in Hügel's Kaschmir, iv (2), p. 426 (1844).

Distribution : India ; China ; Burma ; Malay Peninsula ; Sumatra ; Borneo ; Java.

(a) **Pantoporia selenophora selenophora** (Koll.).

Pantoporia selenophora selenophora Fruhstorfer, in Seitz, Macrolep. ix, p. 631 (1912).

Me Song forest, Prae, N. Siam ; Sriracha forest, S. E. Siam,

Distribution : Mussoorie to Burma ; N. Siam,

b) **Pantoporia selenophora amharina** Fruhst.

Pantoporia selenophora amharina Fruhstorfer, op. cit., p. 631.

Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula.

297. **Pantoporia zeroa** (Moore).

Athyma zeroa Moore, Proc. Zool. Soc. Lond., p. 564 (1872).

Distribution : Kumaon to Burma ; Tonkin ; Hainan ; Siam.

(a) **Pantoporia zeroa galaesus** Fruhst.

Pantoporia zeroa galaesus Fruhstorfer, in Seitz, Macrolep. ix, p.

632, fig. 123 e (1912).

Muak Lek, E. Siam (*Fruhstorfer*) ; Pak Jong, E. Siam (*Godfrey*).

Distribution : Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

(b) **Pantoporia zeroa zeroa** (Moore).

Pantoporia zeroa zeroa Fruhstorfer, op. cit., p. 632.

Me Song forest, Prae, N. Siam (*Godfrey*) ; Chiengmai, N. Siam (*Tennent*).

Distribution : Sikkim to Burma ; N. Siam.

298. **Pantoporia cama cama** (Moore).

Athyma cama Moore, Cat. Lep. Mus. E. I. C. i, p. 174 (1857).

Pantoporia cama cama Fruhstorfer, in Seitz, Macrolep. ix, p. 632 (1912).

1 ♂ : Klong Klung, 1. 3. 24 ; 1 ♂ : Huey Tak Sao, 24. 2. 24

W. Siam (*Stockley*) ; 2 ♂ ♂ (no date) Chiengmai, N. Siam (*Tennent*).

Distribution : Mussoorie to Burma and Siam ; Hainan.

Other subspecies are found in Tonkin and Formosa.

299. **Pantoporia nefte asita** (Moore).

Papilio nefte Cramer, Pap. Exot. iii, pl. 256, figs. E, F (1782).

Athyma asita Moore, Proc. Zool. Soc., p. 13 (1858).

Pantoporia nefte asita Fruhstorfer, in Seitz, Macrolep. ix, p. 633, fig. 123 e (1912).

Me Seng forest, Prae, N. Siam (*Godfrey*) ; Chiengmai, N. Siam (*Tennent*).

Distribution : Shan States to Tavoy ; Tonkin ; Hainan ; Siam.

The species ranges from India to China, and south to Borneo, Sumatra and Java.

300. *Liminitis daraxa daraxa* Moore.

Liminitis daraxa Moore, in Doubleday and Westwood, Gen. Di. Lep. ii, p. 276, pl. 34, fig. 4 (1850); Fruhstorfer, in Seitz, Macrolep. ix, p. 638, fig. 122 f. (1913).

1 ♂ : Chiengmai district, N. Siam, March 1921 (*Tennent*).

Distribution : Sikkim to Burma ; Siam ; S. Annam.

Other subspecies are found in the Malay Peninsula, Borneo, and Sumatra.

301. *Liminitis procris procris* (Cram.).

Papilio procris Cramer, Pap. Exot. ii, pl. 106, figs. E, F. (1777).

Liminitis procris procris Fruhstorfer, in Seitz, Macrolep. ix, p. 640 (1912).

Widely spread over Siam, but nowhere common.

Distribution : Sikkim to Burma ; Siam.

The species ranges from India to China and south to Borneo, Sumatra, Java, Lombok and Flores.

302. *Pandita sinope sinope* Moore.

Pandita sinope Moore, Cat. Lep. Mus. E. I. C. i, p. 102 (1857).

Pandita sinope sinope Fruhstorfer, in Seitz, Macrolep. ix, p. 643, fig. 124 f. (1912).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution : Siam ; Malay Peninsula ; Singapore ; Sumatra ; Java.

A subspecies, *P. s. sinoria* Feld., is found in Borneo, Palawan and the Natunas.

303. *Lebadea martha martha* (Fabr.).

Papilio martha Fabricius, Mant. Ins. ii, p. 56 (1787).

Lebadea martha martha Fruhstorfer, in Seitz, Macrolep. ix, p. 643 (1912).

Widely spread over Siam.

Distribution : S. Burma ; Siam ; Tonkin ; Annam ; Borneo.

Replaced from Manipur to Assam by *L. m. ismene* Doubl.

304. *Lebadea alankara malayana* Fruhst.

Lebadea alankara Horsfield, Cat. Lep. Mus. E. I. C. i, p. 179 (1857).

Lebadea alankara malayana Fruhstorfer, in Seitz, Macrolep. ix, p. 644 (1912).

Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: P. Siam; Malay Peninsula.

The typical form occurs in Java and subspecies are found in Borneo, the Natunas, Palawan, and Sumatra.

305. *Parthenos sylvia* Cram.

Papilio sylvia Cramer, Pap. Exot. i, Taf. 43 (1776).

Distribution: India to China and the Philippines, south to the Greater Sunda Isles., Celebes, New Guinea and the Solomon Islands.

(a) ***Parthenos sylvia gambrisius* (Fabr.).**

Parthenos sylvia gambrisius Fruhstorfer, in Seitz, Macrolep. ix, p. 645 (1912).

Me Song forest, Chiengmai and Chiengrai, N. Siam; Pak Jong and Muak Lek, E. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam.

Distribution: Bengal to Burma; Mergui; Siam; Annam.

(b) ***Parthenos sylvia lilacinus* Butl.**

Parthenos sylvia lilacinus Fruhstorfer, op. cit., p. 645.

Bangnara, P. Siam (*Godfrey*); Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: P. Siam; Malay Peninsula.

306. *Neurosigma doubledayi doubledayi* (Westw.).

Acontia doubledayi Westwood, Cab. Orient. Ent. p. 76, pl. 37, fig. 4 (1848).

Neurosigma doubledayi doubledayi Fruhstorfer, in Seitz, Macrolep. ix, p. 648 (1913).

1 ♂: Doi Sutep, N. Siam, Feb. 1921 (*Tennent*).

Distribution: Sikkim to N. Siam.

Replaced in the Karen Hills by *N. d. nonius* de Nicév.

307. *Euthalia pelea pelea* (Fabr.).

Papilio pelea Fabricius, Mant. Insec. ii, p. 53 (1787).

Tanaëcia pelea pelea Fruhstorfer, in Seitz, Macrolep. ix, p. 650 (1913).

Khao Luang (2,000 ft.), Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : Mergui ; P. Siam ; Malay Peninsula.

Other subspecies are found in Borneo, Sumatra and the Natunas.

308. *Euthalia lepidea* (Butl.).

Adolias lepidea Butler, Ann. Mag. Nat. Hist. 4 (i), p. 71 (1868).

Distribution : India ; Burma ; Tonkin ; Siam ; Malay Peninsula.

(a) ***Euthalia lepidea lepidea* (Butl.).**

Euthalia lepidea lepidea Fruhstorfer, in Seitz, Macrolep. ix, p. 656 (1913).

Me Song forest, Prae, N. Siam (*Godfrey*) ; Klong Suan Mak, W. Siam (*Stockley*).

Distribution : Kumaon to Assam ; Siam ; Malay Peninsula.

(b) ***Euthalia lepidea cognata* Moore.**

Euthalia lepidea cognata Fruhstorfer, op. cit., p. 657.

Muak Lek, E. Siam ; Sriracha forest, S. E. Siam.

Distribution : Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

309. *Euthalia julii* (Boug.).

Nymula julii Bouganville, Voy. "Thetis," ii, p. 345 (1837).

Distribution : N. India to Siam, Annam, the Malay Peninsula, and Sumatra.

(a) ***Euthalia julii odilina* Fruhst.**

Euthalia julii odilina Fruhstorfer, in Seitz, Macrolep. ix, p. 657 (1913).

Me Song forest, Prae, N. Siam ; Muak Lek and Pak Jong, E. Siam ; Sriracha forest, S. E. Siam.

Distribution : Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

(b) ***Euthalia julii julii* (Boug.).**

Euthalia julii julii Fruhstorfer, op. cit., p. 658.

Khao Ram and Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula ; Sumatra.

310. *Euthalia cocytus cocytus* (Fabr.).

Papilio cocytus Fabricius, Mant. Ins. ii, p. 29 (1787).

Euthalia cocytus cocytus Fruhstorfer, in Seitz, Macrolep. ix, p. 658 (1913).

Me Song forest, Prae, N. Siam ; Muak Lek, E. Siam ; Sriracha, S. E. Siam.

Distribution : Typical *cocytus* is confined to Siam, from whence it was described.

E. c. satropaces Hew. occurs from Manipur to S. Burma, and *E. c. ambrysus* Fruhst. in Tonkin.

311. *Euthalia godarti asoka* Feld.

Adolias godarti Gray, Lep. Ins. Nep., p. 14 (1846).

Euthalia godarti asoka Fruhstorfer, in Seitz, Macrolep. ix, p. 659 (1913).

1 ♂ : Bandon, P. Siam, September 1918.

Distribution : Mergui ; P. Siam ; Malay Peninsula.

The species ranges from Mergui to Sumatra, Nias, Borneo, Java, and the Philippines.

312. *Euthalia flora* Butl.

Euthalia flora Butler, Proc. Zool. Soc. Lond., p. 235 (1873) ; Fruhstorfer, in Seitz, Macrolep. ix, p. 659 (1913).

Distribution : From S. Burma to Siam, the Malay Peninsula and Sumatra.

This is a very confusing species and little appears to be known about it at present. It has been recorded from Siam under four different subspecific names :—

(a) ***Euthalia flora flora* Butl.**

Euthalia flora flora Fruhstorfer, Macrolep. ix, p. 659 (1913).

1 ♂ : Bandon, P. Siam, Sep. 1918 ; 1 ♀ : Bangnara, P. Siam, Jan. 1918 (*Godfrey*) ; 1 ♂, 1 ♀ : Hue Sak, Chumporn, P. Siam (*Aagaard*).

The specimens from Bandon and Bangnara were identified at the British Museum.

Distribution : P. Siam ; Malay Peninsula.

(b) ***Euthalia flora maclayi* Dist.**

Euthalia flora maclayi Fruhstorfer, op. cit., p. 660.

1 ♀ : Bandon, P. Siam, Sep. 1918.

This specimen was identified at the British Museum and is in my possession.

Distribution : P. Siam ; Malay Peninsula.

Fruhstorfer (*l. c.*) states that the type of *maclayi* is in the British Museum, but this is not so; neither the type nor any other specimen is to be found there.

(c) **Euthalia flora salangana** Fruhst.

Euthalia flora salangana Fruhstorfer, op. cit., p. 660.

The type of this form came from Salanga Island off the coast of P. Siam. There are two specimens in the British Museum (*Adams Coll.*) from the same locality.

(d) **Euthalia flora andersoni** Moore.

Euthalia flora andersoni Fruhstorfer, op. cit., p. 660.

There are 2 ♂♂, 2 ♀♀ of this form from Renong, P. Siam, in the British Museum (*Adams Coll.*).

Evans* treats *andersoni*, which occurs from S. Burma to Tavoy and P. Siam, as a subspecies of *E. lepidea* Butl.

313. Euthalia cocytina puseda Moore.

Euthalia cocytina Horsfield, Zool. Journ. v, p. 67 (1829).

Euthalia cocytina puseda Fruhstorfer, in Seitz, Macrolep. ix, p. 660, pl. 133 c (1913).

Khao Ram (1,200 ft.), Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula.

The species is found throughout the Macromalayan region, except in Java.

314. Euthalia jahnu jahnides Fruhst.

Adolias jahnu Moore, Cat. Lep. Mus. E. I. C., p. 192 (1857).

Euthalia jahnu jahnides Fruhstorfer, in Seitz, Macrolep. ix, p. 663, pl. 132 c. ♂, b. ♀ (1913).

Me Song forest, Prae, N. Siam ; Muak Lek, E. Siam ; Sriracha forest, S. E. Siam (*Godfrey*) ; Eka Klo, W. Siam (*Stockley*).

Distribution : Tonkin ; Siam.

The typical form occurs from Sikkim to N. Burma, and a subspecies is known from S. Burma.

* Evans, Journ. B. N. H. S. xxix, p. 901 (1924).

315. *Euthalia kesava discispilota* Moore.

Adolias kesava Moore, Trans. Ent. Soc. Lond. v, p. 67, pl. 3, fig. 5 (1859).

Euthalia kesava discispilota Fruhstorfer, in Seitz, Macrolep. ix, p. 664 (1913).

Muak Lek, E. Siam and Sriracha, S. E. Siam (*Godfrey*) ; Chiengmai, N. Siam (*Tennent*) ; Klong Suan Mak, W. Siam (*Stockley*).

Distribution : Shan States ; S. Burma ; Siam.

The typical form occurs from Assam to N. Burma and subspecies are known from Sikkim, Tonkin, Hainan, and Annam.

316. *Euthalia monina grahami* Ril. & Godf.

Papilio monina Fabricius, Mant. Ins. ii, p. 51 (1787) ; Fruhstorfer (*Euthalia*), in Seitz, Macrolep. ix, p. 664 (1913).

Euthalia monina grahami Riley and Godfrey, Journ. N. H. S. Siam iv, p. 178, pl. v, figs. 5, 6 (1921).

Bangnara, P. Siam.

Distribution : P. Siam.

The species ranges from P. Siam through the Malay Peninsula to Borneo, Sumatra, Banka, Java, Bali, Lombok, and the Sulu Isles.

317. *Euthalia garuda apama* Fruhst.

Adolias garuda Moore, Cat. Lep. Mus. E. I. C., p. 186 (1857).

Euthalia garuda apama Fruhstorfer, in Seitz, Macrolep. ix, p. 667 (1913).

Widely distributed over Siam.

Distribution : Cochin-China ; Siam.

The species ranges from Sikkim and Hainan to the Malay Peninsula, Sumatra, Borneo, Java, Palawan and the Sulu Isles.

318. *Euthalia eriphyle* de Nicév.

Euthalia eriphyle de Nicéville, Journ. Bomb. N. H. S. vi, p. 353 (1891).

Distribution : Assam to Burma ; Tonkin ; Siam.

(a) ***Euthalia eriphyle chula* Fruhst.**

Euthalia eriphyle chula Fruhstorfer, in Seitz, Macrolep. ix. p. 670, pl. 129 d (1913).

Muak Lek and Pak Jong, E. Siam ; Sriracha forest, S. E. Siam.

Distribution : Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

(b) **Euthalia eriphyle eriphyle** de Nicév.

Euthalia eriphyle eriphyle Fruhstorfer, op. cit., p. 670.

Me Wong, W. Siam (*Stockley*).

Distribution : Assam to S. Burma ; W. Siam.

Evans (Journ. Bomb. N. H. S. XXIX, p. 903, 1924) treats *eriphyle* as a subspecies of *E. merta* Moore.

319. Euthalia anosia (Moore).

Adolias anosia Moore, Cat. Lep. Mus. E. I. C. i, p. 187 (1857); Fruhsstorfer (*Euthalia*), in Seitz, Macrolep. ix, p. 674 (1913) ; Moulton (*Euthalia*), Sarawak Mus. Journ. ii, 6, p. 226 (1915).

1 ♂ : Me Song forest, Prae, N. Siam, April 1916 (*Godfrey*) ;
1 ♂ : Chiengmai district, N. Siam, Jan. 1921 (*Tennent*).

Distribution : Sikkim ; Assam ; Malay Peninsula ; Sumatra Borneo.

Fruhstorfer (*l. c.*) divides up this species into six different local races, but Moulton (*l. c.*) rejects these on the grounds that the differences given are insufficient and founded on too few specimens. He treats all as one species having the above distribution.

320. Euthalia phemius phemius (Doubl.).

Itanus phemius Doubleday, Gen. Di. Lep. pl. 41, fig. 4 (1848).

Euthalia phemius phemius Fruhstorfer, in Seitz, Macrolep. ix, p. 674, figs. 131 c, d (1913).

Me Song forest, Prae, N. Siam ; Sriracha, S. E. Siam.

Distribution : Sikkim to Burma ; Tonkin ; Siam.

Fruhstorfer (*l. c.*) recognizes races from Hongkong and Perak.

321. Euthalia lubentina (Cram.).

Papilio lubentina Cramer, Pap. Exot. ii, pl. 155, figs. C, D (1779).

Distribution : India ; Ceylon ; China ; Siam ; Malay Peninsula ; Sumatra ; Borneo ; Java ; Philippines.

(a) **Euthalia lubentina indica** Fruhst.

Euthalia lubentina indica Fruhstorfer, in Seitz, Macrolep. ix, p. 676, fig. 13 a (1913).

Bangkok, C. Siam.

Distribution : Sikkim to Burma ; Siam.

(b) **Euthalia lubentina chersonesia** Fruhst.

Euthalia lubentina chersonesia Fruhstorfer, op. cit., p. 676.

Bangnara (*Godfrey*) ; Wang Sai, Songkla (*Kerr*) ; Khao Ram, Nakon Sritamarat (*Pendlebury*)—all in P. Siam.

Distribution : P. Siam ; Malay Peninsula.

322. *Euthalia ludonia siamica* Ril. & Godf.

Euthalia ludonia Staudinger, Deutsche Ent. Zschr. p. 72 (1889).

Euthalia ludonia siamica Riley and Godfrey, Entomol. Iviii, p. 140 (1925).

Euthalia djata ludonia Fruhstorfer, in Seitz, Macrolep. ix, p. 678 (1913).

Pak Jong, E. Siam.

Distribution : Confined to Siam.

Fruhstorfer (*l. c.*) treats *ludonia* as a race of *djata* Distant, and refers it to Palawan.

323. *Euthalia adonia beata* Fruhst.

Papilio adonia Cramer, Pap. Exot. iii, pl. 255, figs. C, D (1782).

Euthalia adonia beata Fruhstorfer, Ent. Zschr. Guben, p. 63 (1905) ; id., in Seitz, Macrolep. ix, p. 679 (1913).

Bangkok, C. Siam.

Distribution : Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

The species ranges from Siam through the Malay Peninsula to Sumatra, Borneo, and Java.

This subspecies was described from a single ♀ sent to Fruhstorfer from Bangkok in 1905 ; no other specimen is known.

324. *Euthalia teuta teuta* (Doubl.).

Adolias teuta Doubleday, Gen. Di. Lep. ii, p. 291, pl. 44, fig. 2 (1850).

Euthalia teuta teuta Fruhstorfer, in Seitz, Macrolep. ix, p. 680 (1913).

Me Song forest Prae, N. Siam ; Sriracha S. E. Siam.

Distribution : Assam ; Burma ; Siam.

Fruhstorfer (*l. c.*) recognizes subspecies from the Andaman Islands ; Luzon and Mindoro ; and Bazilan Island.

325. *Euthalia bellata* Dist.

Euthatia bellata Distant, Rhop. Malay., p. 436, pl. 37, fig. 4 (1886).

Distribution : S. Burma ; Siam ; Malay Peninsula ; Natuna Isles ; Nias ; Sumatra ; Borneo ; Java ; Palawan.

(a) **Euthalia bellata gupta** de Nicév.

Euthalia bellata gupta Fruhstorfer, in Seitz, Macrolep. ix, p. 681
(1913).

1 ♂ : Me Song forest, Prae, N. Siam, Ap. 1918; 1 ♀ : Hup Bon,
Sriracha forest, S. E. Siam, March 1914.

Distribution : Burma ; Siam.

Fruhstorfer (*l.c.*) writes of *gupta* :—“One male discovered by
Bingham in the renowned Thungyeen forest is, together with a pair
in the Nicéville collection of the Calcutta Museum, all we possess of
of this form”.

Evans* treats *gupta* as a race of *E. teuta* Doubl.

(b) **Euthalia bellata goodrichi** Fruhst.

Euthalia bellata goodrichi Fruhstorfer, in Seitz, Macrolep. ix, p. 681
(1913).

Ban Lan Sah Kah, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula.

326. Euthalia recta monilis Moore.

Symphaedra recta de Nicéville, Butt. Ind. ii, p. 188 (1886).

Euthalia recta monilis Fruhstorfer, in Seitz, Macrolep. ix, p. 682
(1913).

Me Song forest, Prae, N. Siam ; Muak Lek, E. Siam ; Sriracha,
S. E. Siam.

Distribution : Fruhstorfer (*l. c.*) restricts this subspecies to
Siam.

The typical form occurs from Assam to Burma.

327. Euthalia evelina (Stoll).

Papilio evelina Stoll, in Cramer's Pap. Exot. Suppl. v, p. 132 (1790).

Distribution : India to S. China, Hainan, and the Philippines ;
south to the Greater Sunda Islands.

(a) **Euthalia evelina annamita** Moore.

Euthalia evelina annamita Fruhstorfer, in Seitz, Macrolep. ix, p. 684
(1913).

Muak Lek, E. Siam ; Sriracha forest, S. E. Siam.

Distribution : Cochin-China ; Siam.

* Evans, Journ. Bomb. N. H. S. xxix, 4, p. 906 (1924).

Specimens taken in the above localities in Siam agree exactly with Moore's types of *annamita* in the British Museum.

(b) **Euthalia evelina vallona** Fruhst.

Euthalia evelina vallona Fruhstorfer, op. cit., p. 685 (1913).

Kanburi, W. Siam (*Fruhstorfer*).

Distribution: Fruhstorfer restricts this subspecies to Siam.

328. Euthalia dunya Doubl.

Euthalia dunya Doubleday, Gen. Di. Lep. ii, p. 291, pl. 44, fig. 3 (1850); Fruhstorfer, in Seitz, Macrolep. ix, p. 687, fig. 130 b (1913); Moulton, Sarawak Mus. Journ. ii, (6) p. 228 (1915).

Khao Luang (2,500 ft.), Nakon Sritamarat, P. Siam (*Pendlebury*); Kao Pawta Luang Keo (c. 800 m.), Ranawng, P. Siam (*Kerr*); Klong Klung, W. Siam (*Stockley*); Pran River, W. Siam (*Ladell*).

Distribution: S. Burma; Siam; Malay Peninsula; Borneo; Sumatra; Java.

A subspecies is found in Nias.

Fruhstorfer (*l. c.*) splits this variable species into five local races but, according to Moulton (*l. c.*), all the differences given for Bornean, Sumatran and Javan forms are covered by a series (*dunya*) from Sarawak. The Nias form is, apparently, quite distinct.

329. Adolias dirtea (Fabr.).

Papilio dirtea Fabricius, Syst. Ent. iii (1), p. 59 (1793).

Distribution: India to Hainan and the Philippines, south to the Malay Peninsula, Borneo, Sumatra, Nias, and Java.

(a) **Adolia dirtea jadeitina** Fruhst.

Adolias dirtea jadeitina Fruhstorfer, in Seitz, Macrolep. ix, p. 687, fig. 127 b (1913).

Me Song forest, Prae, N. Siam; Muak Lek, E. Siam; Sriracha, S. E. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*); Eka Klo, W. Siam (*Stockley*).

Distribution: Manipur to Burma; Siam.

(b) **Adolias dirtea dirtea** (Fabr.).

Adolias dirtea dirtea Fruhstorfer, op. cit., p. 687.

Khao Ram and Khao Luang (*Pendlebury*); Wat Kiriwong (*Kerr*)—all in Nakon Sritamarat, P. Siam.

Distribution: Malay Peninsula; Sumatra; Borneo.

330. *Adolias cyanipardus albopunctata* Crowley.

Symphaedra cyanipardus Butler, Proc. Zool. Soc. Lond., p. 613 (1868).

Adolias cyanipardus albopunctata Fruhstorfer, in Seitz, Macrolep. ix, p. 693 (1913).

Muak Lek, E. Siam (Fruhstorfer).

Distribution: Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

The typical form occurs from Assam to the Karens, and subspecies are found in Borneo, Sumatra, and Bangka.

331. *Apatura parisatis siamensis* Fruhst.

Apatura parisatis Westwood, Gen. Di. Lep. ii, p. 305 (1850).

Apatura parisatis siamensis Fruhstorfer, in Seitz, Macrolep. ix, p. 698, pl. 115 a (1913).

Me Song forest, Prae, N. Siam; Pak Jong and Muak Lek, E. Siam (Godfrey); Nakon Sritamarat, P. Siam (Pendlebury); Pan Yao, W. Siam (Stockley).

Distribution: Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

The species ranges from Hong Hong to Ceylon, the Philippines and the Greater Sunda Islands.

I retain Fruhstorfer's name for Siamese specimens, but am of the opinion that his *siamensis* is merely a seasonal form of typical *parisatis*.

332. *Apatura ambica miranda* Fruhst.

Apatura ambica Kollar, in Hugel's Kaschmir, iv, 2, p. 431 (1844).

Apatura ambica miranda Fruhstorfer, in Seitz, Macrolep. ix, p. 700 (1913).

Me Song forest, Prae, and Chiengrai, N. Siam; Muak Lek, E. Siam; Sriracha, S. E. Siam.

Distribution: Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

The typical form occurs from Kashmir to the Dawnas and in Hainam; and subspecies known from Chitral, Tonkin, and Sumatra.

Evans ("The Identification of Indian Butterflies", p. 89, 1927) treats as synonyms of *ambica ambica* the following races recorded

by Fruhstorfer:—*a. zamaa* Hew., *a. namouna* Doubl., *a. bhavana* Moore, and *a. garlanda* Fruhst.

333. Sephis chandra chandra (Moore).

Castalia chandra Moore, Cat. Lep. Mus. E. I. C., p. 200 (1858); Bingham (*Sephisa*), Faun. Br. Ind. Butt. i, p. 239; Fruhstorfer (*Sephisa*), in Seitz, Macrolep. ix, p. 701 (1913).

Both Bingham and Fruhstorfer record this species from N. Siam.

Distribution: Sikkim to N. Burma; N. Siam.

A subspecies is found in Formosa.

334. Eulaceura osteria osteria (Westw.).

Apatura osteria Westwood, in Dblday., Westw. & Hew. Gen. Di. Lep., p. 305 (1850).

Apatura osteria osteria Fruhstorfer, in Seitz, Macrolep ix, p. 702 (1913).

1 ♂: Kao Pawta Chongdong, c. 400 m., Ranawng (Renong), P. Siam, 20. 1.29 (*Kerr*).

Distribution: P. Siam; Malay Peninsula; Borneo; Sumatra; Java.

Subspecies are known from S. Burma, Hainan, and Nias.

335. Hestina nama nama Doubl.

Diadema nama Doubleday, Ann. Mag. Nat. Hist. xvi, p. 232 (1845).

Hestina nama nama Fruhstorfer, in Seitz, Macrolep. ix, p. 704 (1913).

Chiengrai, N. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*).

Distribution: Mussoorie to Burma; Tonkin; Siam.

Subspecies are found in Perak and Sumatra.

336. Calinaga sudassana Melv.

Calinaga sudassana Melvill, Trans. Ent. Soc. Lond., p. 121 (1893); Bingham, Faun. Br. Ind. i, p. 467 (1905); Fruhstorfer, in Seitz, Macrolep. ix, p. 705 (1914).

Calinaga buddha sudassana Fruhstorfer, in Seitz, Macrolep. ix, p. 462 (1912).

Both Bingham and Fruhstorfer record this species from N. Siam.

Distribution: Shan States to the Karen; N. Siam.

Reduced by Fruhstorfer in 1912 to subspecific rank, *sudassana* was restored by the same authority in 1914 to its original status as a good species.

337. *Calinaga buddha avalokita* Fruhst.

Calinaga buddha Moore, Cat. Lep. Mus. E. I. C. p. 163, pl. 3a (1857).

Calinaga buddha avalokita Fruhstorfer, in Seitz, Macrolep. ix, p. 705 (1914).

Type from N. Siam in the Tring Museum.

Distribution : Burma ; Siam.

The typical form occurs from Murree to Kumaon and subspecies are found in Sikkim, Assam, China, and Formosa.

Fruhstorfer (*l. c.*) writes of *avalokita* :—“The discovery of this form and its occurrence on the border of Siam and Burma, side by side with *C. sudassana* Melv., prove the specific rights of the latter as a separate species.”

338. *Herona marathus marathus* Doubl.

Herona marathus Doubleday, Gen. Di. Lep. p. 294, pl. 41, fig. 3 (1850).

Herona marathus marathus Fruhstorfer, in Seitz, Macrolep. ix, p. 705, fig. 116 b (1914).

Me Song forest, Prae, N. Siam and Pak Jong, E. Siam (*Godfrey*) ; Chiengmai, N. Siam (*Tennent*) ; Me Wong, W. Siam (*Stockley*).

Distribution : Sikkim to the Shan States ; Siam.

Other subspecies are found in S. Burma and the Andamans.

339. *Euripus halitherses* Doubl.

Euripus halitherses Doubleday, Gen. Di. Lep. p. 293, pl. 41, fig. 2 (1850).

Distribution : India ; Burma ; Siam ; Malay Peninsula ; Sumatra ; Borneo ; Java ; Philippines.

(a) ***Euripus halitherses halitherses* Doubl.**

Euripus halitherses halitherses Fruhstorfer, in Seitz, Macrolep. ix, p. 707 (1914).

Chiengrai, N. Siam ; Nam Phat (Sen Tor), N. E. Siam ; Muak Lek, E. Siam.

Also obtained at Ban Na Sao in French Laos.

Distribution : Sikkim to Burma ; Siam ; French Laos.

(b) **Euripus halitherses pfeifferae** Feld.

Euripus halitherses pfeifferae Fruhstorfer, op. cit., p. 707.

Ban Lan Sah Kah, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula.

340. Euripus consimilis eurinus Fruhist.

Diadema consimilis Westwood, Gen. Di. Lep. p. 281 (1850).

Euripus consimilis eurinus Fruhstorfer, in Seitz, Macrolep. ix, p. 710.

1 ♀ : Chiengsen, N. Siam, 24. 4. 21.

Distribution : Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

The typical form occurs from the N. W. Himalayas to Burma and a subspecies is known from S. India.

341. Prothoe franckii angelica Butl.

Prothoe franckii Butler, Ann. Mag. Nat. Hist. (5) XVI, pp. 53, 54 (1885).

Prothoe franckii angelica et *vilma* Fruhstorfer, in Seitz, Macrolep. ix, p. 714 (1914).

Klong Klung, W. Siam (*Stockley*) ; Nakon Sritamarat, P. Siam (*Pendlebury*) ; Kao Pawta Chongdong, c. 800 m., Ranawng (Renong), P. Siam (*Kerr*).

Distribution : Karen Hills to S. Burma ; Siam.

The species ranges from Assam to the Malay Peninsula Sumatra, Borneo, Banka, Nias, Palawan and the Philippines.

Fruhstorfer's *vilma* (described from Muak Lek, E. Siam) is probably a synonym of *angelica*.

342. Prothoe calydonia calydonia Hew.

Nymphalis calydonia Hewitson, Exot. Butt. ii, p. 86, pl. 42, figs. 3, 4 (1855).

Prothoe calydonia calydonia Fruhstorfer, in Seitz, Macrolep. ix, p. 717 (1914).

Khao Ram (1,200 ft.), Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : P. Siam ; Malay Peninsula ; Sumatra ; Borneo.

A subspecies, *P. c. belisama* Crow., occurs from the Karen Hills to S. Burma.

343. *Eriboea athamas samatha* Moore.

Papilio athamas Drury, Ill. Exot. Ent. i, p. 5, pl. 2, fig. 4 (1770).

Eriboea athamas samatha Fruhstorfer, in Seitz, Macrolep. ix, p. 719 (1914).

Widely spread over Siam, but nowhere common.

Distribution: Karen Hills to S. Burma; Siam; Malay Peninsula.

The species ranges from Sikkim to China and the Philippines; south to the Malay Peninsula, Sumatra, Borneo and Java.

344. *Eriboea arja* (Feld.).

Charaxes arja Felder, Nov. Reise, Lep. Rhop. p. 348 (1867); Fruhstorfer (*Eriboea*), in Seitz, Macrolep. ix, p. 720 (1914).

Me Song forest, Prae, and Chiengrai, N. Siam; Muak Lek, E. Siam; Sriracha forest, S. E. Siam (Godfrey); Me Wong, W. Siam (Stockley).

Distribution: Sikkim to Burma; Tonkin; Siam.

345. *Eriboea jalusus* (Feld.).

Charaxes jalusus Felder, Nov. Reise, Lep. Rhop. p. 438, pl. 59, fig. 59 (1867); Fruhstorfer (*Eriboea*), in Seitz, Macrolep. ix, p. 722 (1914).

1 ♂ : Me Wong (1,000 ft.), W. Siam, 22. 2. 24 (Stockley); 1 ♂ : Kao Kalakiri (c. 300 m.), Pattani, P. Siam, 29. 3. 28 (Kerr).

Distribution: Karen Hills to S. Burma; Tonkin; Siam; Malay Peninsula; Sumatra; Borneo.

346. *Eriboea eudamippus nigrobasalis* (Lathy.).

Charaxes eudamippus Doubleday, Ann. Soc. Ent. Fr., p. 218 (1843).

Eriboea eudamippus nigrobasalis Fruhstorfer, in Seitz, Macrolep. ix, p. 722 (1914).

Me Song forest, N. Siam; Muak Lek, E. Siam; Pechaburi River, W. Siam.

Distribution: N. Burma; Siam; Cambodia.

According to Fruhstorfer, the species ranges from Nepal to S. Burma, Siam, and Annam; and from Yunnan to the Liu-Kiu Islands, Formosa and Hainan.

347. *Eriboea nepenthes nepenthes* (Grose-Smith).

Charaxes nepenthes Grose-Smith, Ent. Mon. Mag. xx, p. 58 (1883).

Eriboea nepenthes nepenthes, Fruhstorfer, in Seitz, Macrolep. ix, p. 723 (1914).

1 ♂ : Me Song forest, Prae, N. Siam, April 1918.

Distribution : Known only from Siam (from whence the species was described) and the Shan States.

A subspecies *E. n. fugator* Fruhst. is known from Tonkin.

348. *Eriboea dolon grandis* (Roths. & Jord.).

Charaxes dolon Westwood, Cab. Or. Ent. p. 5, 5 pl. 27, figs. 2, 3 (1848).

Eriboea dolon grandis Fruhstorfer, in Seitz, Macrolep. ix, p. 723 (1914).

1 ♂ : Me Song forest, Prae, N. Siam, April 1918.

Distribution : Shan States to the Karen; N. Siam.

The species ranges from the Himalayas to N. Siam.

349. *Eriboea delphis delphis* (Doubl.).

Charaxes delphis Doubleday, Ann. Soc. Ent. Fr. p. 217, pl. 7 (1843).

Eulepis delphis delphis Fruhstorfer, in Seitz, Macrolep. ix, p. 723 (1914).

Me Song forest, Prae, N. Siam; Muak Lek, E. Siam; Petchaburi River, W. Siam.

Distribution : Assam to Burma; Siam.

The species ranges from Assam to the Malay Peninsula, Borneo, Sumatra, Java, and Palawan.

350. *Charaxes fabius sulphureus* Roths. & Jord.

Papilio fabius Fabricius, Spec. Ins. ii, p. 12 (1781).

Charaxes fabius sulphureus Fruhstorfer, in Seitz, Macrolep. ix, p. 730 (1914).

Me Song forest, Prae, N. Siam; Muak Lek, E. Siam; Sriracha forest, S. E. Siam (*Godfrey*); Me Wong, W. Siam (*Stockley*).

Distribution : Assam to Burma; Siam.

The species ranges from Sikkim to the Malay Peninsula, Sumatra, Borneo, Java, Celebes and the Philippines.

351. *Charaxes polyxena* (Cram.).

Papilio polyxena Cramer, Pap. Exot. i, pl. 54, figs. A, B (1779); Fruhstorfer (*Charaxes*), in Seitz, Macrolep. ix, p. 731 (1914).

Distribution : From India and China to the Malay Peninsula, Sumatra, Nias, Banka, Java, Borneo, and Palawan.

(a) ***Charaxes polyxena hierax f. hierax* Feld.**

Charaxes hierax Felder, Nov. Reise, Lep. Rhop., p. 442 (1867).

Me Song forest, Prae, N. Siam (*Godfrey*) ; Chiengrai, N. Siam (*Tennent*) ; Me Wong, W. Siam (*Stockley*).

Distribution : Sikkim to N. Burma ; Siam.

(b) **Charaxes polyxena hierax f. agna** Moore.

Charaxes agna Moore, Proc. Zool. Soc. Lond., p. 832 (1878).

Pak Jong and Muak Lek, E. Siam ; Sriracha forest, S. E. Siam.

Distribution : Karen Hills to S. Burma ; Siam.

(c) **Charaxes polyxena repetitus** Butl.

Charaxes repetitus Butler, Lep. Exot., p. 16 (1872).

Khao Ram and Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*) ; Khao Kalakiri, Pattani, P. Siam (*Kerr*).

Distribution : P. Siam ; Malay Peninsula ; Borneo ; Sumatra ; Natunas.

352. Charaxes marmax marmax Westw.

Charaxes marmax Westwood, Cab. Or. Ent., p. 43, pl. 21 (1848).

Charaxes marmax marmax Fruhstorfer, in Seitz, Macrolep. ix, p. 736, fig. 131 b (1914).

Me Song forest, Prae, N. Siam (*Godfrey*) ; Chiengmai, N. Siam (*Tennent*).

Distribution : Kumaon to Burma ; Siam.

Other subspecies are found in Tonkin and the Malay Peninsula.

353. Charaxes aristogiton aristogiton Feld.

Charaxes aristogiton Felder, Nov. Reise, Lep. Rhop., p. 445 (1867).

Charaxes aristogiton aristogiton Fruhstorfer, in Seitz, Macrolep. ix, p. 736, pl. 136 d (1914).

4 ♂♂ : Me Song forest, Prae, N. Siam, April 1916.

Distribution : Sikkim to Burma ; Siam.

Ch. a. aristogiton f. adamsoni Moore also occurs in Siam, two males having been obtained in the Me Song forest in 1914.

Fruhstorfer has described a subspecies, *Ch. a. peridoneus*, from Tonkin.

354. Charaxes kahruba (Moore)

Haridra kahruba Moore, Lep. Ind. ii, p. 235, pl. 171, fig. 1 (1893-96) ; Fruhstorfer (*Charaxes*), in Seitz, Macrolep. ix, p. 736, figs. 136 b, c (1914).

1 ♂ : Chiengmai district, N. Siam (*Tennent*).

Distribution : Kumaon to Burma ; N. Siam.

355. Charaxes durnfordi durnfordi Dist.

Charaxes durnfordi Distant, Entomol. xvii, p. 191 (1884); Fruhstorfer, in Seitz, Macrolep. ix, p. 738 (1914).

Khao Luang (2,000 ft.), Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: S. Burma; P. Siam; Malay Peninsula.

Other subspecies are known from Borneo, Sumatra and Java.

356. Pareba vesta sordice Fruhst.

Papilio vesta Fabricius, Mant. Ins. ii, p. 14 (1787).

Pareba vesta sordice Fruhstorfer, in Seitz, Macrolep. ix, p. 741 (1915).

2 ♂ ♂, 2 ♀ ♀: Doi Sutep, N. Siam, Jan. 1921 (*Tennent*).

Distribution: Burma; Siam.

The species ranges from India to China and Tonkin, south to Sumatra, Java and Bali.

ERYCINIDAE.

LIBYTHEINAE.

357. Libythea narina Godt.

Libythea narina Godart, Encyc. Néth. ix, p. 171 (1819).

Distribution: India to the Philippines, south to New Guinea.

(a) **Libythea narina rohini** Marshall.

Libythea narina rohini Fruhstorfer, in Seitz, Macrolep. ix, p. 768, fig. 139 f (1914).

Me Song forest, N. Siam; Pak Jong, E. Siam; Sriracha forest, S. E. Siam; Pechaburi River, W. Siam.

Distribution: Assam to Burma; Siam; Malay Peninsula.

(b) **Libythea narina rohini f. libera** de Nicév.

Libythea rohini var. *libera* Bingham, Faun. Br. Ind. Butt. i, p. 477. Pechaburi River, W. Siam.

Distribution: Occurs with *rohini* in Assam, Burma and Siam.

358. Libythea lepita lepita Moore.

Libythea lepita Moore, Cat. Lep. Mus. E. I. C. p. 240 (1857).

Libythea lepita lepita Fruhstorfer, in Seitz, Macrolep. ix, p. 769, fig. 139 e (1914).

2 ♂ ♂ : Me Lamoung (2,000 ft.), W. Siam, 15. 4. 20 (*Stockley*).

Distribution: Kashmir to N. Burma; Siam.

The species ranges from India to China and Japan, south to Siam.

359. *Libythea myrrha sanguinalis* Fruhst.

Libythea myrrha Godart, Encyc. Méth. ix, p. 171 (1819).

Libythea myrrha sanguinalis Fruhstorfer, in Seitz, Macrolep. ix, p. 769, fig. 139 e (1914).

Me Song forest, Prae and Chiengrai, N. Siam.

Distribution : Sikkim to Siam.

The species ranges from India to China and Tonkin, south to the Malay Archipelago.

360. *Libythea geoffroyi alompra* Moore.

Libythea geoffroyi Godart, Encycl. Méth. ix, p. 813 (1823).

Distribution : Burma to the Solomon Islands.

(a) *Libythea geoffroyi alompra* Moore.

Libythea geoffroyi alompra Fruhstorfer, in Seitz, Macrolep. ix, p. 770, fig. 139 f (1914).

Pak Jong and Muak Lek, E. Siam; Sriracha, S. E. Siam; Pechaburi River, W. Siam.

Distribution : S. Burma; Siam; French Laos.

(b) *Libythea geoffroyi alompra f. hauxwelli* Moore.

Libythea hauxwelli Bingham, Faun. Br. Ind. Butt. i, p. 478.

Pechaburi River, W. Siam.

Distribution : Occurs with *alompra* in Tenasserim and Siam.

RIODININAE.

361. *Zemeros flegyas allica* (Fabr.).

Papilio flegyas Cramer, Pap. Exot. iii, pl. 280, figs. E, F (1780).

Zemeros flegyas allica Fruhstorfer, in Seitz, Macrolep. ix, p. 773 (1914).

Common in all forest areas in Siam.

Distribution : Confined to Siam.

The species ranges from Mussoorie to the Malay Peninsula and Archipelago.

362. *Dodona egeon* (Doubt.).

Taxila egeon Doubleday, Gen. Di. Lep., p. 422 (1851); Fruhstorfer (*Dodona*), in Seitz, Macrolep. ix, p. 776 (1914).

1 ♂ : Doi Chang (4,500 ft.), N. Siam, 14. 4. 2.

Distribution : Nepal to N. Siam.

363. *Dodona henrici binghami* Moore.

Dodona henrici Holland, Trans. Amer. Ent. Soc. xiv, p. 119 (1887).

Dodona henrici binghami Fruhstorfer, in Seitz, Macrolep. ix, p. 778, fig. 141 b (1914).

1 ♂ : Pang Yao, W. Siam, 20. 2. 20 (Stockley).

Distribution : Burma; Siam.

The typical form occurs in Hainan.

364. *Dodona deodata deodata* Hew.

Dodona deodata Hewitson, Exot. Butt. iii, pl. 1. figs. 1, 2 (1866).

1 ♂ : Me Song forest, Prae, N. Siam, April 1918.

Distribution : Shan States to S. Burma; Siam.

Other subspecies are found in Assam and Annam.

365. *Abisara fylla fylla* (Doubl.).

Taxila fylla Doubleday, List. Lep. B. M. pt. 2, p. 2 (1847).

Abisara fylla fylla et *fyllaria* Fruhstorfer, in Seitz, Macrolep. ix, p. 780 (1914).

Doi Chang (4,500 ft.) and Me Song forest, Prae (*Godfrey*); Chiengmai (*Tennent*)—all in N. Siam. Also 1 ♂ of the d. s. f. *fyllaria* Fruhst., Me Lamoung, W. Siam (Stockley).

Distribution : Mussoorie to Siam.

Subspecies are found in China and Tonkin.

366. *Abisara echerius* (Stoll).

Papilio echerius Stoll, Pap. Exot. v. pl. 35, figs. 1 a, 1 b (1790).

Distribution : N. India to China, Tonkin, Hainan and the Philippines; south to Annam and Siam.

(a) Abisara echerius siamensis Fruhst.

Abisara echerius siamensis Fruhstorfer, in Seitz, Macrolep ix, p. 781, pl. 138 c, ♂, 140 d, ♀ (1914).

Me Song forest, N. Siam and Sriracha forest, S. E. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*).

Distribution: Fruhstorfer (*b. c.*) restricts this subspecies to Siam.

(b) Abisara echerius angulata f. abnormis Moore.

Abisara echerius angulata f. abnormis Fruhstorfer, op. cit., p. 782, pl. 140 b.

1 ♂ : Me Lamoung, W. Siam, 15. 4. 20 (Stockley).

Distribution : Burma; W. Siam.

367. **Abisara kausambi kausambi** Feld.

Abisara kausambi Felder, Wien. Ent. Monats. iv, p. 397 (1860).

Abisara kausambi kausambi Fruhstorfer, in Seitz, Macrolep. ix, p. 782 (1914).

1 ♀ : Adang, 10. 1. 28; 1 ♀ : Nakon Sritamarat, 25. 4. 28; 1 ♂ : Ban Prakawp, Songkla, 18. 7. 28 (*Kerr*)—all in P. Siam.

Distribution : P. Siam ; Malay Peninsula.

The species ranges from India to the Malay Peninsula, Sumatra, Nias, Borneo, and Java.

368. **Abisara neophron neophron** (Hew.)

Sospita neophron Hewitson, Exot. Butt. ii, pl. 1, fig. 3 (1861).

Abisara neophron neophron Fruhstorfer, in Seitz, Macrolep. ix, p. 785 (1914).

Me Song forest, Prae, N. Siam ; Nam Phat (Sen Tor), N. E. Siam ; Khao Sabap, Chantabun, S. E. Siam.

Distribution : Assam to N. Burma ; Siam.

The species ranges from India to China and Tonkin, south to the Malay Peninsula.

369. **Abisara savitri savitri** Feld.

Abisara savitri Felder, Wien. Ent. Monats., p. 397 (1860).

Abisara savitri savitri Fruhstorfer, in Seitz, Macrolep. ix, p. 785 (1914).

Khao Ram, Nakon Sritamarat (*Pendelbury*) ; Klawng Ton, Satul (*Kerr*)—P. Siam.

Distribution : Mergui ; P. Siam ; Malay Peninsula.

Subspecies are found in Sumatra, Borneo and the Natuna Islands.

370. **Laxita damajanti damajanti** (Feld.).

Abisara damajanti Felder, Wien. Ent. Monats. iv, p. 397 (1860).

Laxita damajanti damajanti Fruhstorfer, in Seitz, Macrolep. ix, p. 789 (1914).

1 ♂ : Kao Luang, Nakon Sritamarat, 29. 4. 28; 1 ♂ : Kao Kalakiri, Pattani, 31. 3. 28—P. Siam (*Kerr*).

Distribution : P. Siam ; Malay Peninsula.

Subspecies are found in Sumatra and Borneo.

371. **Laxita telesia** (Hew.).

Taxila telesia Hewitson, Exot. Butt. ii, pl. 1, fig. 5 (1861).

Distribution: Siam; Malay Peninsula; Sumatra; Borneo.

(a) **Laxita telesia boulleti** Fruhst.

Laxita boulleti Fruhstorfer, in Seitz, Macrolep. ix, p. 790, pl. 141 a (1914).

Laxita telesia boulleti Riley and Godfrey, Journ. N. H. S. Siam, iv, p. 178, pl. 5, fig. 7 (1921).

1 ♂ : Me Song forest, Prae, N. Siam, April 1918.

Described by Fruhstorfer, from a single female obtained by the Harmand Mission in Siam. Although *boulleti* is very distinct from typical *telesia*, Riley and Godfrey (*l. c.*) do not consider it worthy of more than subspecific rank.

Distribution: Confined to Siam.

(b) **Laxita telesia lyclene** Fruhst.

Laxita telesia lyclene Fruhstorfer, op. cit., p. 790.

Khao Ram, Nakon Sritamarat (*Pendlebury*); Khao Kalakiri, Pattani and Tambon Kao Panom, Krabi (*Kerr*)—all in P. Siam.

Distribution: P. Siam; Malay Peninsula.

372. **Laxita orphna lyncestis** (de Nicév.).

Taxila orphna Boisduval, Spec. Gen. i, pl. 21, fig. 4 (1836).

Laxita orphna lyncestis Fruhstorfer, in Seitz, Macrolep. ix, p. 790 (1914).

1 ♂ : Kao Keo Range, Satul, 13. 4. 28; 4 ♂♂ : Kao Soi Dao (c. 400 m.), Patalung, 29. 4. 30 (*Kerr*)—P. Siam.

Distribution: P. Siam; Malay Peninsula.

The typical form occurs in Borneo and a subspecies is known from Sumatra.

373. **Taxila thuisto** Hew.

Taxila thuisto Hewitson, Exot. Butt. ii, pl. 1; figs. 5, 6 (1861); Fruhstorfer, in Seitz, Macrolep. ix, p. 791 (1914).

Distribution: Burma to the Malay Peninsula, Sumatra, and Borneo.

(a) **Taxila thuisto ephorus** Fruhst.

Taxila thuisto ephorus Fruhstorfer, op. cit., p. 791.

Hinlap, E. Siam (*Fruhstorfer*); Me Song forest, N. Siam; Muak Lek, E. Siam; Sriracha forest, S. E. Siam (*Godfrey*).

Distribution : Confined to Siam.

- (b) *Taxila thuisto thuisto* Hew.

Taxila thuisto thuisto Fruhstorfer, op. cit., p. 792.

1 ♂ : Ban Ton, Satul, P. Siam, 15. 3. 28 (*Kerr*).

Distribution : P. Siam ; Malay Peninsula ; Singapore.

374. *Taxila haquinus* (Fabr.).

Papilio haquinus Fabricius, Ent. Syst. iii, p. 55 (1793) ; Fruhstorfer (*Taxila*), in Seitz, Macrolep. ix, p. 792 (1914).

Distribution : Fruhstorfer (*l. c.*) gives the range of this species as :—“North Borneo to Siam and in the whole Macromalayana, as well as in Palawan and some satellite islands between Borneo and Palawan”.

- (a) *Taxila haquinus fasciata* Moore.

Taxila haquinus fasciata Fruhstorfer, op. cit., p. 792.

Me Song forest, Prae, N. Siam (*Godfrey*) ; Ban Samui, W. Siam (*Stockley*).

Distribution : Assam to Burma ; N. and W. Siam.

- (b) *Taxila haquinus berthae* Fruhst.

Taxila haquinus berthae Fruhstorfer, op. cit., p. 792, pl. 138 g.

Muak Lek, E. Siam ; Sriracha forest, S. E. Siam.

Distribution : Confined to Siam.

375. *Stiboges nymphidia nymphidia* Butl.

Stiboges nymphidia Butler, Proc. Zool. Soc. p. 309 (1876).

Stiboges nymphidia nymphidia Fruhstorfer, in Seitz, Macrolep. ix, p. 796 (1914).

Hup Bon, Sriracha forest, S. E. Siam (*Godfrey*) ; Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*) ; Surat, Satul, and Krabi, P. Siam (*Kerr*).

Distribution : Bhutan to Burma ; Siam ; Malay Peninsula.

Subspecies are found in W. China and Tonkin, in Sumatra, and in Java.

LYCAENIDAE.

PORITIINAE.

376. *Poritia sumatrae* (Feld.).

Pseudodipsas sumatrae Felder, Novara Reise, Lep. ii, p. 259 (1865) ; Seitz (*Poritia*), Macrolep. ix, p. 1008, pl. 154 h. (1927).

1 ♂ : Chawng, Trang, 14. 4. 28; 1 ♂ : Wang Sai, Songkla, 20. 7. 28 (*Kerr*)—P. Siam.

Distribution: This very rare species is known only from Sumatra and P. Siam.

377. *Poritia hewitsoni tavoyana* Doh.

Poritia hewitsoni Moore, Proc. Zool. Soc. p. 775, pl. 41, fig. 10 (1865).

Poritia hewitsoni tavoyana Seitz, Macrolep. ix, p. 1008 (1926).

Me Song forest, Prae, N. Siam (*Godfrey*); Pan Yao, W. Siam (*Stockley*).

Distribution: Burma; Siam; Annam.

The typical form occurs from Kumaon to Assam.

378. *Poritia erycinoides phraatica* Hew.

Pseudodipsas erycinoides Felder, Nov. Reise, Lep. ii, p. 259 (1865).

Poritia erycinoides phraatica Seitz, Macrolep. ix, p. 1008 (1926).

1 ♂ : Me Song forest, Prae, N. Siam, April 1918 (*Godfrey*); 1 ♂ : (*in Brit. Mus.*) Raheng, W. Siam (*Doherty*).

Distribution: Burma; Siam; Malacca.

The species ranges from Assam to Sumatra, Borneo, Java and Mindano.

379. *Simiskina pharyge* (Hew.).

Poritia pharyge Hewitson, Trans. Ent. Soc. Lond., p. 345 (1874); Seitz, Macrolep. ix, p. 1009, fig. 154 i (1927).

Renong, P. Siam (*Doherty*).

Distribution: Mergui; Siam; Perak; Penang; Sumatra; Borneo; Java.

380. *Simiskina pasira dohertyi* Evans.

Poritia pasira Moulton, Journ. Straits Branch R. A. Soc. No. 60, p. 122, fig. 2 (1911).

Simiskina pasira dohertyi Evans, Journ. Bomb. N. H. S. xxx (2), p. 335, pl. 26, fig. H. 3.4 (1925).

1 ♂ : Chiengmai district, N. Siam, Jan. 1921 (*Tennent*).

Distribution: S. Burma to Tavoy; Siam.

The typical form occurs in Borneo.

The species *pasira* is not included in Seitz' "Macrolepidoptera."

381. **Simiskina phalia** (Hew.).

Poritia phalia Hewitson, Trans. Ent. Soc. Lond., p. 345 (1874); Seitz, Macrolep. ix, p. 1009, fig. 154 l (1927).

Me Song forest, Prae, N. Siam; Sriracha forest, S. E. Siam.

Distribution: Burma to the Malay Peninsula, Sumatra and Borneo.

382. **Simiskina pavonica veturia** (Fruhst.).

Simiskina paronica de Nicéville, Journ. Bomb. N. H. S, p. 28 (1895).

Poritia pavonica veturia Fruhstorfer, Berl. Ent. Zeit. p. 201 (1911); Seitz, Macrolep ix, p. 1009, fig. 54 h (1927).

Hin Lap, S. E. Siam (*Fruhstorfer*).

Distribution: Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

The typical form occurs in Sumatra and a subspecies is found in Borneo.

383. **Deramas livens jasoda** (de Nicév.).

Zarona jasoda de Nicéville, Journ. As. Soc. Beng. p. 280 (1888).

Zarona zanella Bingham, Faun. Br. Ind. Butt. ii, p. 471 (1907).

Deramas livens jasoda Seitz, Macrolep. ix, p. 1010, fig. 155 a (1927).

Siam-Tenasserim frontier (*Bingham*).

Distribution: Burma; Siam.

The species extends to Sumatra, Borneo and Java.

Bingham (*l. c.*) writes of *zanella* :—"I have very little doubt that this form is only the ♀ of *jasoda*".

384. **Deramas bradamante** (Doherty MS.).

Zarona bradamante de Nicéville, Butt. Ind. iii, p. 36, foot note (1890).

Distribution: "Lower Siam."

MILETINAE*.

385. **Miletus croton mallus** (Fruhst.).

Gerydus croton Doherty, Journ. As. Soc. Beng. p. 439 (1889).

Gerydus croton mallus Fruhstorfer, in Seitz, Macrolep. ix, p. 816, fig. 141 f (1915).

* I am indebted to Captain Riley for the following note :— "*Miletus* Hübner (1818) originally contained two species, *polycletus* and *symethus*. Scudder (Historical Sketch, p. 219, 1875) specified *polycletus* as the type, and the generic name has been used in that sense ever since.

5 ♂ ♂, 10 ♀ ♀ : Me Song forest, Prae, N. Siam, April 1916 and 1918.

Distribution : Siam ; Tonkin ; Annam.

The typical form occurs from Bhamo to S. Burma.

386. *Miletus boisduvali irroratus* (Druce).

Miletus boisduvali Moore, Cat. Lep. Mus. E. I. C. p. 19 (1857).

Gerydus boisduvali irroratus Fruhstorfer, in Seitz, Macrolep. ix, p. 817 (1915).

Chiengrai and Me Song forest, Prae, N. Siam ; Pak Jong, E. Siam ; Bangkok, C. Siam ; Pechaburi River, W. Siam.

Distribution : Confined to Siam.

The species ranges from Sikkim to New Guinea.

387. *Miletus biggsii* (Dist.).

Gerydus biggsii Distant, Rhop. Malay., p. 206 (1844) ; Fruhstorfer, in Seitz, Macrolep. ix, p. 819, fig. 141 g. (1915).

Nakon Sritamarat, P. Siam (*Pendlebury*).

388. *Miletus ancon siamensis* (Godf.).

Gerydus ancon Doherty, Journ. As. Soc. Beng. p. 438 (1889).

Gerydus ancon siamensis Godfrey, Journ. N. H. S. Siam, ii, p. 134 (1916) ; Riley and Godfrey, op. cit., p. 179, pl. 5, figs. 8, 9 (1921).

Me Song forest, Prae, N. Siam ; Muak Lek, E. Siam.

Distribution : Confined to Siam.

The typical form occurs in Burma and a subspecies is found in Borneo.

389. *Allotinus drumila grisea* Ril. & Godf.

Miletus drumila Moore, Proc. Zool. Soc. Lond. p. 77 (1865) ; Fruhstorfer (*Allotinus*), in Seitz, Macrolep. ix, p. 815 (1915).

Allotinus drumila grisea Riley and Godfrey, Journ. N. H. S. Siam, iv, p. 180, pl. 6, fig. 3, ♀ (1921).

1 ♀ : Me Song forest, Prae, N. Siam, April 1918.

Distribution : Confined to Siam.

The typical form occurs from Sikkim to Assam.

Unfortunately he overlooked the quite definite type fixation by Doubleday and Hewitson (Gen. Diurn. Lep. p. 502, 1852) who specify *symethus* as the type. *Miletus* (Hübner 1818) therefore replaces *Gerydus* (Boisduval, 1835) ; *symethus* being the geno-type in both cases. Under Article 5 of the International Rules the name of the subfamily becomes *Miletinae*."

390. **Allotinus posidion** Fruhst.

Allotinus posidion Fruhstorfer, Zeits. Wiss. Insbiol. ix, p. 368 (1913).

Distribution : Burma to the Malay Peninsula, Sumatra, Borneo, Java and the Philippines.

(a) **Allotinus posidion rekkia** Ril. & Godf.

Allotinus posidion rekkia Riley and Godfrey, Journ. N. H. S. Siam, iv, p. 180, pl. 6, figs. 1, 2 (1921).

Pak Jong and Muak Lek, E. Siam ; Sriracha, S. E. Siam.

Distribution : Confined to E. and S. E. Siam.

(b) **Allotinus posidion atacinus** Fruhst.

Allotinus posidion atacinus Fruhstorfer, in Seitz, Macrolep. ix, p. 811 (1915).

3 ♀ ♀ : Klong Klung, W. Siam, 1. 3. 24 (Stockley).

Distribution : Burma ; W. Siam.

391. **Logania massalia massalia** Doh.

Logania massalia Doherty, J. A. S. B., p. 37 (1891).

Logania massalia massalia Fruhstorfer, in Seitz, Macrolep. ix, p. 807 (1915).

1 ♂ : Sadao, Songkla, P. Siam, 22. 7. 28 (Kerr).

Distribution : Assam to Burma ; P. Siam.

Fruhstorfer (*l. c.*) recognizes subspecies from Borneo, Sumatra and Java.

L. massalia is a very rare species.

LYCAENINAE.

392. **Taraka hamada mendesia** Fruhst.

Miletus hamada Druce, Cist. Ent. i, p. 361 (1875).

Taraka hamada mendesia Fruhstorfer, in Seitz, Macrolep. ix, p. 884, fig. 154 f. (1915).

Me Song forest, Prae, N. Siam.

Distribution : Sikkim to Borneo.

The typical form occurs in Japan and subspecies are found in China and Java.

393. **Spalgis epius epius** (Westw.).

Lucia epius Westwood, Gen. Di. Lep. ii, p. 502 (1852).

Spalgis epius epius Fruhstorfer, in Seitz, Macrolep. ix, p. 881 (1915).

1 ♂ : Klong Pong Kapo, W. Siam, 2. 4. 20 (Stockley).

Distribution: N. India to Siam.

The species ranges from India to Hainan, Formosa, and the Philippines; south to Borneo, Sumatra and Nias.

394. *Pithecopus hylax hylax* (Fabr.).

Papilio hylax Fabricius, Syst. Ent. p. 526 (1775).

Pithecopus hylax hylax Fruhstorfer, in Seitz, Macrolep. ix, p. 879 (1915).

Me Song forest, Prae and Chiengrai, N. Siam; Pak Jong, E. Siam.

Distribution: Sikkim to Burma; Siam.

The species ranges from Sikkim to the Malay Peninsula and Archipelago.

395. *Neopithecops zalmora zalmora* (Butl.).

Pithecopus zalmora Butler, Cat. Fabr. Lep. B. M. p. 161 (1869).

Pithecopus zalmora zalmora Fruhstorfer, in Seitz, Macrolep. ix, p. 880 (1915).

Me Song forest, Prae, and Chiengrai, N. Siam; Muak Lek, E. Siam (*Godfrey*); Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: Kumaon to the Malay Peninsula.

The species ranges from N. India to the Malay Archipelago.

396. *Megisba malaya sikkima* Moore

Lycaena malaya Horsfield, Cat. Lep. Mus. E. I. C. p. 78 (1828).

Megisba malaya sikkima Fruhstorfer in Seitz, Marerolep. ix, p. 857 (1915).

Chiengrai, N. Siam; Pak Jong, E. Siam.

Distribution: Kumaon to Burma; Siam.

The species ranges from India to Formosa and the Philippines south to New Guinea.

397. *Talicada metana* Ril. & Godf.

Talicada metana Riley and Godfrey, Journ. N. H. S. Siam, iv, p. 181, pl. 6, fig. 4 (1921).

Me Song forest, Prae, N. Siam.

Distribution: Confined to Siam.

398. **Castalius rosimon rosimon** (Fabr.).

Papilio rosimon Fabricius, Syst. Ent. p. 523 (1775).

Castalius rosimon rosimon Fruhstorfer, in Seitz, Macrolep. ix, p. 885 (1915).

Widely spread over Siam.

Distribution : Ceylon ; India ; Burma ; Siam ; Annam.

The species ranges from India to the Lesser Sunda Islands.

399. **Castalius caleta decidia** (Hew.).

Lycaena caleta Hewitson, Exot. Butt. v, Lyc., pl. 1, fig. 1 (1876).

Castalius caleta decidia Fruhstorfer, in Seitz, Macrolep. ix, p. 891, figs. 153 a, b (1915).

1 ♂, 1 ♀ : Chieng Sen, N. Siam, 24. 4. 21.

Distribution : Sikkim to Burma ; Siam.

The typical form occurs in the Celebes and subspecies are found in Ceylon, and the Philippines.

400. **Castalius ethion ethion** (Doubt. & Hew.).

Lycaena ethion Doubleday and Hewitson, Gen. Di. Lep. ii, p. 490, pl. 76, fig. 3 (1852).

Castalius ethion ethion Fruhstorfer, in Seitz, Macrolep. ix, p. 886 (1915).

Me Song forest, Prae, and Chiengrai, N. Siam ; Sriracha, S. E. Siam.

Distribution : Assain to Burma ; Andaman Islands ; Siam.

The species ranges from India to the Philippines.

401. **Castalius elna noliteia** Fruhst.

Lycaena elna Hewitson, Exot. Butt. v, pl. 1, fig. 8 (1876).

Castalius elna noliteia Fruhst., in Seitz, Macrolep. ix, p. 889 (1923).

Me Song forest, Prae, and Chiengrai, N. Siam ; Pak Jong and Muak Lek, E. Siam ; Sriracha, S. E. Siam.

Distribution : Sikkim to Burma ; Andaman Islands ; Siam.

The species is distributed from India to Java and the Celebes.

402. **Castalius roxus roxana** (Godt.).

Polyommatus roxus Godart, Ency. Meth. ix, p. 659 (1823).

Castalius roxus roxana Fruhstorfer, in Seitz, Macrolep. ix, p. 888 (1915).

Me Song forest, Prae, N. Siam; Pak Jong, E. Siam.

Distribution: Assam to N. Burma; Siam.

According to Fruhstorfer, the species is distributed:—"from Indo-China to Siam and Tonkin, in the whole of Macro- and Micromalayana, in Celebes and all the Philippines."

403. *Euchrysops cnejus* (Fabr.).

Hesperia cnejus Fabricius, Ent. Syst. Suppl. p. 430 (1798); Seitz

(*Catochrysops*), Macrolep. ix, p. 922 (1924); Evans (*Euchrysops*), Journ. Bomb. N. H. S. xxx (2), p. 339, pl. 27, fig. 16. 1 (1925).

Chiengrai, N. Siam; Muak Lek, E. Siam; Bangkok, C. Siam.

Distribution: India to Hainan, south to Australia and the South Sea Islands.

404. *Euchrysops pandava pandava* (Horsf.).

Lycaena pandava Horsfield, Cat. Lep. Mus. E. I. C., p. 84 (1829).

Euchrysops pandava pandava Evans, Journ. Bomb. N. H. S. xxx, p. 340 (1925); Seitz (*Catochrysops*), Macrolep. ix, p. 922, fig. 135 k (1924).

Chiengrai, N. Siam; Pak Jong, E. Siam; Bangkok, C. Siam.

Distribution: India to Hainan, south to Borneo, Java and the Natuna Islands.

Evans has described a subspecies from Ceylon.

405. *Everes parhasius rileyi* Godf.

Hesperia parhasius Fabricius, Ent. Syst. iii (1), p. 289 (1793); Evans (*Everes*), Journ. Bomb. N. H. S., xxx (2), p. 340 (1925).

Everes rileyi Godfrey, Journ. N. H. S. Siam, ii, p. 135 (1916); Riley and Godfrey, op. cit., iv, pl. 7, figs. 4, 5 (1921).

Me Song forest, Prae, N. Siam; Pak Jong, E. Siam; Bangnara, P. Siam;

Distribution: Confined to Siam.

Evans (*l. c.*) recognizes as good races: *E. p. parhasius* Fabr. (S. India, Ceylon); *E. p. assamica* Tyt. (Assam-Burma.); and *E. p. pila* Evans (Andamans and Nicobars).

Seitz (Macrolep. ix, p. 923) treats *parhasius* as a subspecies of *argiades* Pallas.

406. *Bothrinia chennelli* (de Nicév.).

Cyaniris chennelli de Nicéville, Journ. As. Soc. Beng. p. 72, pl. 1, fig. 10 (1883); Seitz (*Bothrinia*), Macrolep. ix, p. 877 (1924).

3 ♂♂, 12 ♀♀ : Doi Chang (4,500 ft.) and Phar Mieng, Chiengrai, N. Siam, April 1921.

Distribution : Assam to Burma ; Siam.

407. *Lycaenopsis haraldus* (Fabr.).

Papilio haraldus Fabricius, Mant. Ins. ii, p. 82, n. 744 (1787) ; Fruhstorfer (*Lycaenopsis*), in Seitz, Macrolep. ix, p. 874 (1923).

La-un, Ranawng (Renong), 30. 12. 28 ; Nang Yawn, Takuapa, 8. 2. 29 ; Nawng Chik, Krabi, 19. 3. 30 ; Tambon Kao Panom, Krabi, 30. 3. 30 (*Kerr*)—all in P. Siam.

Distribution : P. Siam ; Malay Peninsula ; Borneo ; Java.

Fruhstorfer (*l. c.*) divides up this rare and variable species into three races : *ananga* Feld. (Malay Peninsula), *cornuta* Druce (Borneo), and *haraldus* Fabr. (Java). The validity of these races is, however, questionable.

408. *Lycaenopsis transpектa* Moore.

Lycaenopsis transpектa Moore, Proc. Zool. Soc. p. 139 (1879) ; Fruhstorfer, in Seitz, Macrolep. ix, p. 867, figs. 152 e, f (1924).

Pak Jong, E. Siam.

Distribution : Sikkim to Burma ; Siam.

409. *Lycaenopsis puspa gisca* Fruhst.

Polyommatus puspa Horsfield, Cat. Lep. Mus. E. I. C., p. 67 (1827).

Lycaenopsis puspa gisca Fruhstorfer, in Seitz, Macr. ix, p. 869 (1924).

Cyaniris puspa imperatrix Butler, Stet. Ent. Zeit. lxxi, p. 285. (1910) ; Fruhstorfer (*Lycaenopsis*), in Seitz, Macrolep. ix, p. 869 (1924).

Doi Chang (4,500 ft.) and Me Song forest, Prae, N. Siam ; Pak Jong and Muak Lek, E. Siam ; Sriracha, S. E. Siam.

Distribution : Sikkim to Burma ; Siam.

According to Fruhstorfer, the species ranges from India to the Philippines and from the Andamans to the Key Islands.

Fruhstorfer, following Butler, gives *imperatrix* as the Siamese race, but this is a mistake. Siamese specimens are referable to *gisca* and not to *imperatrix*, which has no trace of white on the forewing above.

410. *Lycaenopsis limbata placida* Moore.

Polyommatus limbatus Moore, Proc. Zool. Soc. p. 139 (1879).

Lycaenopsis limbatus placida Fruhstorfer, in Seitz, Macrolep. ix, p. 872 (1924).

3 ♂♂ : Chiengrai, N. Siam, April 1921.

Distribution : Sikkim to Burma; Siam.

The species ranges from India to Tonkin, Formosa and the Philippines; south to Borneo, Sumatra and Java.

411. *Lycaenopsis cardia dilecta* Moore.

Lycaena cardia Felder, Sitzungsber, Wiss. Wien. xl, p. 459 (1860).

Lycaenopsis cardia dialectus Fruhstorfer, in Seitz, Macrolep. ix, p. 865 (1924).

4 ♂♂, 2 ♀♀ : Doi Chang (4,500 ft.), April 1921 (*Godfrey*);

1 ♂, 1 ♀ : Chiengmai, December 1921 (*Tennent*)—N. Siam.

Distribution : Kashmir to Burma; Siam.

The species ranges from India to Hainan, China, Formosa and the Philippines; south to the Malay Peninsula and Archipelago.

412. *Lycaenopsis argiola jynteana* (de Nicév.).

Papilio argiolus Linnaeus, Syst. Nat. ed. x, p. 483 (1758).

Lycaenopsis argiolus jynteana Evans, Journ. Bomb. N. H. S. xxx. p. 344 (1925).

Lycaenopsis limbatus jynteana Fruhstorfer, in Seitz, Macrolep. ix, p. 873 (1922).

1 ♂, 1 ♀ : Doi Chang (4,500 ft.), N. Siam, April 1921 (*Godfrey*);
1 ♂ : Hot Spring, Me Lamoung (2,000 ft.), W. Siam, April 1920 (*Stockley*).

Distribution : Sikkim to Burma; Siam.

Fruhstorfer (*l. c.*) treats *jynteana* as a subspecies of *limbatus* Moore, but expresses doubt as to whether it really belongs to that species.

413. *Chilades laius laius* (Cram.).

Papilio laius Cramer, Pap. Exot. iv, p. 62 (1780); Fruhstorfer (*Chilades*), in Seitz, Macrolep. ix, p. 926 (1924); Evans (*Chilades*), Journ. Bomb. N. H. S., xxx (2), p. 344 (1925).

Distribution : India to Hongkong, Formosa and the Philippines; south to Siam.

Evans (*l. c.*) recognizes a subspecies from Tavoy.

414. *Zizera trochilus* (Frey.).

Lycaena trochilus Freyer, Neu. Beit. Schmett. v, p. 98 (1844); Seitz (*Chilades*), Macrolep. ix, p. 926 (1924).

2 ♂♂ : Muak Lek, E. Siam, 30. 9. 23; 1 ♀ : Chiengrai, 10. 5. 21.

Distribution: According to Seitz, this species extends:—“from S. E. Europe through Egypt, Syria, Anterior Asia to India, to the east across the Malayan Archipelago as far as the Philippines and to the south as far as Australia.”

415. *Zizera maha* (Koll.).

Lycaena maha Kollar, in Hugel's Kaschmir, iv, p. 422 (1848); Evans (*Zizera*), Journ. Bomb. N. H. S. xxx (2), p. 345 (1925); Joicey and Talbot (*Zizera*), Bulletin Hill Mus., iii (3), p. 157 (1929).

8 ♂♂, 7 ♀♀ : Chiengrai, N. Siam, April 1921.

Distribution: N. India to Burma; Hainan; Siam.

Seitz makes no mention of this species.

416. *Zizera gaika* (Trim.).

Lycaena gaika Trimen, Trans. Ent. Soc. Lond. iii (1), p. 403 (1862); Seitz (*Zizera*), Macrolep. ix, p. 925 (1924).

Widely spread over Siam.

Distribution: India to Sumatra and Java.

417. *Zizera otis* (Fabr.).

Papilio otis Fabricius, Mant. Ins. ii, p. 73 (1787); Seitz (*Zizera*), Macrolep. ix, p. 925, fig. 153 c (1924).

Widely spread over Siam.

Distribution: India to China and Formosa, south to Borneo and Java.

418. *Syntarucus plinius* (Fabr.).

Hesperia plinius Fabricius, Ent. Syst. iii, p. 284 (1793); Fruhstorfer (*Tarucus*), in Seitz, Macrolep. ix, p. 893 (1923); Evans (*Syntarucus*), Journ. Bomb. N. H. S., xxx (2), p. 350 (1925); Joicey and Talbot (*Syntarucus*), Bulletin Hill Mus. iii (3), p. 154 (1929).

Me Song forest, Prae, and Chiengrai, N. Siam; Bangkok, C. Siam.

Distribution: India to China, Hainan and Formosa; south to Borneo and Java.

Joicey and Talbot (*l. c.*) point out that this genus is distinguished from *Tarucus* by the hairy eyes and the close approximation of veins 11 and 12.

419. *Catochrysops strabo* (Fabr.).

Hesperia strabo Fabricius, Ent. Syst. iii (1), p. 287 (1793); Seitz (*Catochrysops*), Macrolep. ix, p. 921, fig. 153 k (1924).

Chiengrai, N. Siam; Pak Jong, E. Siam; Sriracha, S. E. Siam; Bangnara, P. Siam.

Distribution: India to China, Hainan, and the Philippines; south through Malaya to Australia.

420. *Catochrysops lithargyria* (Moore).

Lampides lithargyria Moore, Ann. Mag. N. H. (4) xx, p. 340 (1877).

Catochrysops strabo ab. *lithargyria* Seitz, Macrolep. ix, p. 921 (1924).

1 ♂ : Sriracha, S. E. Siam, 27. 9. 14. (*Godfrey*); 1 ♂ Nakon Chaisee, W. Siam (*in Br. Mus.*).

Distribution: Ceylon; Assam-Burma; Andamans; Siam.

Seitz (*l. c.*) treats *lithargyria* as an aberration of *strabo*, but both Riley and Evans regard it as a good species.

421. *Lampides boeticus* (Linn.).

Papilio boeticus Linnaeus, Syst. Nat. xii, 1, p. 789 (1767); Evans (*Lampides*), Journ. Bomb. N. H. S. xxx, p. 351, pl. 27, fig. 29. 1 (1925); Fruhstorfer (*Polyommatus*), in Seitz, Macrolep. ix, p. 895 (1923).

Chiengrai, N. Siam; Bangnara, P. Siam; Bangkok, C. Siam.

Distribution: Europe, Africa, Asia through Malaya to Australia and the Hawaiian Islands.

Fruhstorfer (*l. c.*) wrongly places this species in the genus *Polyommatus*.

422. *Nacaduba angusta* (Druce).

Cupido angusta Druce, Proc. Zool. Soc. p. 349, pl. 32, fig. 9 (1873).

Distribution: Burma to the Malay Peninsula, Sumatra, Nias, Borneo, Java, the Celebes, and the Philippines.

(a) ***Nacaduba angusta kerriana* Dist.**

Nacaduba angusta kerriana Seitz, Macrolep. ix, p. 914, fig. 152 h (1924).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution : Burma ; Siam.

(b) **Nacaduba angusta f. albida** Ril. & Godf.

Nacaduba angusta f. albida Riley and Godfrey, Entomol. lviii, p. 141, pl. 3, fig. 2 (1925).

Phar Mieng, near Chiengrai, N. Siam.

Distribution : Burma ; N. Siam.

Riley and Godfrey (*l. c.*) are of the opinion that, when more material has been obtained, *albida* will prove to be a good subspecies.

423. Nacaduba pactolus continentalis Fruhst.

Nacaduba pactolus Felder, Sitzungsber. Ak. Wiss. Wien. xl p. 458 (1860).

Nacaduba pactolus continentalis Fruhstorfer, Zool. Med. Leid. ii, p. 114 (1916) ; Seitz, Macrolep. ix, p. 914 (1924).

2 ♂♂, 1 ♀ : Me Song forest, Prae, N. Siam, April 1916 ; 6 ♂♂ : Phar Mieng, near Chiengrai, N. Siam, April 1921.

Distribution : Sikkim to Burma ; Siam.

The species is distributed from India to Hainan and Formosa south to the Solomon Islands.

424. Nacaduba vajuva Fruhst.

Nacaduba vajuva Evans, Journ. Bomb. N. H. S. xxx, p. 611 (1925).

Nacaduba pavana vajuva Fruhstorfer, Zool. Med. Lied. ii, p. 108 (1916) ; Seitz, Macrolep. ix, p. 913 (1924).

Chiengrai, N. Siam, April 1921.

Distribution : Sikkim to Burma ; Siam.

According to genitalia *vajuva* is a good species.

425. Nacaduba pavana (Horsf.).

Lycaena pavana Horsfield, Cat. Lep. Mus. E. I. C., p. 77 (1828) ; Evans (*Nacaduba*), Journ. Bomb. N. H. S. xxx, p. 611 (1925).

2 ♂♂ : Me Song forest, Prae, N. Siam, 14. 4. 16.

Distribution : S. Burma ; Andamans ; Siam ; Sumatra ; Borneo ; Java.

426. Nacaduba viola merguiana (Moore.).

Lampides viola Moore, Ann. Mag. N. H. (4) xx, p. 340 (1877).

Lycaenesthes merguiana Moore, Journ. As. Soc. Beng., p. 23 (1884).

Nacaduba viola merguiana Seitz, Macrolep. ix, p. 917 (1924).

4 ♂ ♂ : Chiengrai, N. Siam, April 1921.

Distribution : Sikkim to Siam.

The species extends from the Himalayas to Australia.

427. *Nacaduba atrata euplea* Fruhst.

Lycaena atrata Horsfield, Cat. Lep. Mus. E. I. C., p. 78 (1828).

Nacaduba atrata euplea Evans, Journ. Bomb. N. H. S. xxx, p. 612 (1925).

Nacaduba perusia euplea Seitz, Macrolep. ix, p. 919 (1924).

2 ♂ ♂ : Phar Mieng, near Chiengrai, N. Siam, April 1921.

Distribution : Sikkim to Burma; Andamans; Nicobars; Siam.

The typical form occurs in Java, Bali and Lombok.

428. *Nacaduba akaba gythion* Fruhst.

Nacaduba akaba Druce, Proc. Zool. Soc. Lond, p. 350 (1873).

Nacaduba akaba gythion Evans, Journ. Bomb. N. H. S. xxx, p. 612 (1925).

Nacaduba atrata gythion Seitz, Macrolep. ix, p. 919 (1924).

1 ♀ : Bangkok, C. Siam.

Distribution : Ceylon; S. India; Sikkim to Burma; Siam.

The typical form occurs in Borneo.

Gythion is a subspecies of *akaba* and not of *atrata* which, Riley* informs me, has been misidentified by nearly all authors.

429. *Nacaduba berenice aphyia* Fruhst.

Lycaena berenice H.-Sch, Stett. Ent. xxxiii, p. 74 (1869).

Nacaduba berenice aphyia Fruhstorfer, Leid. Zool. Med. ii, p. 127 (1916); Seitz, Macrolep. ix, p. 918 (1924).

Me Song forest, Prae, and Chiengrai, N. Siam; Pak Jong, E. Siam.

Distribution : Siam; Malay Peninsula.

The species extends from Burma to New Guinea.

* Riley, *in litt.*, 29. 9. 26.

430. *Nacaduba nora* (Feld.).

Lycaena nora Felder, Sitzungsber. Ak. Wiss. Wien. xl, p. 458 (1860); Seitz, Macrolep. ix, p. 915 (1924).

Widely spread over Siam.

Distribution : India to Formosa, Hainan and the Philippines, south to Australia.

431. *Nacaduba dubiosa sivoka* Evans.

Nacaduba dubiosa Semper, Journ. Mus. Godeff. xiv, p. 159 (1879).

Nacaduba dubiosa sivoka Evans, Journ. Bomb. N. H. S. xxx, p. 613 (1925).

Me Song forest, Prae, N. Siam; Bangkok, C. Siam.

Distribution : Sikkim to Siam.

Both Riley and Evans regard *dubiosa* as a good species; Seitz (Macrolep. ix, p. 916) does not.

432. *Nacaduba donina* (Snell.).

Lycaena donina Snellen, Tidj. v. Ent. xlvi, p. 262 (1901).

Nacaduba nora donina Seitz, Macrolep. ix, p. 916 (1924).

1 ♂ : Muak Lek, E. Siam, 30. 7. 23 (Godfrey); 1 ♂ : Chiengmai, N. Siam. (Tennent).

Distribution : Siam; Borneo; Java; Celebes.

Seitz (*l. c.*) treats *donina* as a subspecies of *nora*, but Riley* regards it as a good species.

433. *Jamides bochus bochus* (Cram.).

Papilio bochus Cramer, Pap. Exot. iv, p. 210 (1782).

Jamides bochus bochus Seitz, Macrolep. ix, p. 901 (1924).

Me Song forest, Prae and Chiengrai, N. Siam; Sriracha, S. E. Siam.

Distribution : Ceylon; India; Burma; Andamans; Siam; Malay Peninsula.

The species ranges from India through Malaya to Australia.

434. *Jamides cleodus pura* (Moore).

Lampides cleodus Felder, Nov. Reise Lep. ii, p. 272 (1865).

* Riley, *in litt.*, 30. 8. 24.

Jamides cleodus pura Evans, Journ. Bomb. N. H. S. xxx, p. 614 (1925).

Lampides celeno pura Seitz, Macrolep. ix, p. 904 (1924).

1 ♂ : Upper Me Klong (1,500 ft.), W. Siam, 14. 1. 24 (Stockley).

Distribution : S. China ; Assam to Siam.

The species ranges from India to Formosa and the Philippines, south to Sumatra and Nias.

435. *Jamides celeno celeno* (Cram.).

Papilio celeno Cramer, Pap. Exot. i, pl. 31, figs. C, D (1775).

Lampides celeno celeno Seitz, Macrolep. ix, p. 904 (1924).

Widely spread over Siam.

Distribution : India ; Burma ; Siam ; Malaya.

The species ranges from India to Hainan and Formosa, south to New Guinea.

436. *Jamides elpis eurysaces* (Fruhst.).

Polyommatus elpis Godart, Encyc. Méth. p. 654 (1823).

Lampides elpis eurysaces Seitz, Macrolep. ix, p. 908 (1924).

Chiengmai, N. Siam (*Tennent*) ; Muak Lek, E. Siam (*Godfrey*) ; Me Wong and Khambengpet, W. Siam (*Stockley*).

Distribution : Sikkim to Burma ; S. India ; Andamans ; Hainan ; Siam.

The species ranges from India to Hainan, Formosa and the Philippines; south to New Guinea.

437. *Jamides coerulea* (Druce).

Cupido coerulea Druce, Proc. Zool. Soc. Lond. p. 349 (1873).

Jamides coerulea Evans, Journ. Bomb. N. H. S. xxx, p. 615 (1925).

Kankena lampides coerulea Seitz, Macrolep. ix, p. 907, figs. 151 a, 153 a (1924).

1 ♂ : Me Wong (2,000 ft.), W. Siam, 12. 4. 20 (Stockley).

Distribution : Assam to the Malay Peninsula, Borneo and Sumatra.

438. **Azarus urios** Ril. & Godf.

Azarus urios Riley and Godfrey, Journ. Nat. Hist. Soc. Siam, iv, p. 182, pl. 6, fig. 5 (1921).

Me Song forest, Prae, N. Siam.

Distribution : Siam ; Burma.

439. **Lycaenesthes emolus emolus** (Godt.).

Polyommatus emolus Godart, Encyl. Méth. ix, p. 256 (1823).

Lycaenesthes emolus emolus Seitz, Macrolep. ix, p. 897 (1924).

Me Song forest, Prae and Chiengrai, N. Siam ; Sriracha forest, S. E. Siam ; Bangkok, C. Siam.

Distribution : S. India ; Sikkim to Burma ; Tonkin ; Siam.

The species ranges from India through Malaya to New Guinea and Australia.

440. **Lycaenesthes lycaenina lycambes** Hew.

Lycaenesthes lycaenina Felder, Verh. zool-bot. Ges. Wien., xviii, p. 281 (1868).

Lycaenesthes lycambes Hewitson, Ill. Di. Lep., p. 220 (1862-1878).

Lycaenesthes lycaenina lycambes Seitz, Macrolep. ix, p. 898 (1924).

Chiengrai, N. Siam ; Pak Jong, E. Siam ; Bangkok, C. Siam.

Distribution : Sikkim to Siam.

The species ranges from India to Hainan ; south to New Guinea and Australia.

441. **Niphanda cymbia cymbia** de Nicév.

Niphanda cymbia de Nicéville, Journ. As. Soc. Beng. p. 76 (1883).

Niphanda cymbia cymbia Evans, Journ. Bomb. N. H. S. xxxi, p. 716 (1926).

Niphanda fusca marcia Seitz, Macrolep. ix, p. 901 (1927).

Me Song forest, Prae, and Chiengrai, N. Siam.

Distribution : Sikkim to N. Burma ; N. Siam.

A subspecies, *N. c. tessellata* Moore, occurs from S. Burma to Java.

Work done by Toxopeus proves that *fusca* and *cymbia* are separate species. *Marcia* Fawcett is a synonym of *cymbia* and not, as stated by Seitz, "an excellent race".

442. *Heliophorus epicles chinensis* (Fruhst).

Ilerda epicles Godart, Encyl. Méth. ix, p. 646 (1823).

Ilerda epicles chinensis Seitz, Macrolep. ix, p. 933 (1924).

Me Song forest and Chiengrai, N. Siam.

Distribution: Tonkin; Siam; Annam.

The species ranges from N. India to Hainan and Formosa, south to Java and Sumatra.

THECLINAE.

CURETIS-Group.

443. *Curetis saronis indosinica* Fruhst.

Curetis saronis Moore, Proc. Zool. Soc. Lond., p. 587 (1877); Evans, Journ. Bomb. N. H. S. xxx, p. 623 (1925).

Curetis insularis indosinica Seitz, Macrolep. ix, p. 935 (1924).

Me Song forest, Prae, N. Siam; Muak Lek and Pak Jong, E. Siam; Sriracha, S. E. Siam.

Distribution: Siam; S. Annam.

Evans (*l. c.*) recognizes as good races: *C. s. saronis* Moore (Andamans); *C. s. gloriosa* Moore (Cachar-Burma); and *C. s. nicobaria* Swin. (Nicobars).

444. *Curetis sperthis* (Feld.).

Anops sperthis Felder, Novara Reise, Lep. p. 222 (1865); Evans (*Curetis*), Journ. Bomb. N. H. S. xxx (3), p. 623 (1925).

Curetis insularis sperthis Seitz, Macrolep. ix, p. 935 (1924).

Pak Jong, E. Siam; Sriracha, S. E. Siam.

Distribution: Hainan; Mergui; Siam; Malay Peninsula; Borneo; Java.

Seitz (*l. c.*) treats *sperthis* as a subspecies of *insularis* Horsf., but both Riley and Evans regard it as a good species.

445. *Curetis bulis malayica* (Feld.).

Anops bulis Doubleday and Hewitson, Gen. Di. Lep. ii, p. 473 (1852).

Anops malayica Felder, Nov. Reise, Lep. ii, p. 221, pl. 28, fig. 18 (1865).

Curetis sanatana malayica Seitz, Macrolep. ix, p. 936 (1927).

Sriracha, S. E. Siam and Bangnara, P. Siam (*Godfrey*) ; Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*) ; Tepa, Songkla and Khao Kalakiri, P. Siam (*Kerr*).

Distribution : Karen Hills to S. Burma; Siam; Malay Peninsula.
The typical form occurs from Mussoorie to N. Burma.

446. *Curetis acuta fortunatus* Fruhst.

Curetis acuta Moore, Ann. Mag. N. H. p. 50 (1877) ;
Curetis bulis fortunatus Seitz, Macrolep. ix, p. 935 (1924).
Me Song forest, Prae, N. Siam and Sen Tor (Nam Pat), N. E. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*).

Distribution : Tonkin; Yunnan; Siam; Annam.
Seitz treats *acuta* as a subspecies of *bulis*.

447. *Curetis insularis pseudoinsularis* Fruhst.

Phaedra insularis Horsfield, Cat. Lep. E. I. C. p. 125 (1829).
Curetis insularis pseudoinsularis, Seitz, Macrolep. ix, p. 935 (1924).
1 ♂ : Kao Kalakiri (c. 1,000 m.), Pattani, P. Siam, 4. 4. 28
(*Kerr*).

Distribution : P. Siam; Malacca; (?) Sumatra.
The typical form occurs in Java.

AMBLYPODIA-Group.

448. *Iraota timoleon pandara* Fruhst.

Papilio timoleon Stoll, in Suppl. Cramer, Pap. Exot. v, pl. 32, figs. 4, 4 d (1790).

Iraota timoleon pandara Seitz, Macrolep. ix, p. 944 (1926).
1 ♂ : Bangkok, C. Siam, 20. 1. 21 (*Godfrey*); 1 ♀ : Me Lamoung (2,000 ft.), W. Siam (*Stockley*).

Distribution : Seitz (*l. c.*) restricts this subspecies to Siam.

The species ranges from India to China and the Philippines, South to Malaya.

449. *Horsfieldia *anita anita* (Hew.).**

Amblypodia anita Hewitson, Cat. Lyc. B. M., p. 14 (1862).

* *Horsfieldia* Riley, Entomologist, lv, p. 26 (1922).

Horsfieldia anita anita Riley, Entomol. IV, p. 27 (1922).

Horsfieldia narada anita Seitz, Macrolep. ix, p. 945 (1926).

Pak Jong, E. Siam; Sriracha, S. E. Siam (*Godfrey*); Tepa, Songkla, P. Siam (*Kerr*).

Distribution: Typical *anita*, which was described from Siam, extends from the Nilgiris to Khandala and across to Bengal, Assam, Burma and Siam.

Subspecies are found in Ceylon, Java, the Andamans, and Hainan.

Seitz (*l. c.*) incorrectly places *anita* under *narada* Horsf.

450. *Thaduka multicaudata multicaudata* Moore.

Thaduka multicaudata Moore, Proc. Zool. Soc. Lond. p. 836, pl. 52, fig. 7 (1878); Seitz, Macrolep. ix, p. 947 (1926).

Thaduka multicaudata multicaudata Evans, Journ. Bomb. N. H. S. xxx (3), p. 624 (1925).

Me Song forest, Prae, N. Siam.

Distribution: S. Burma; Siam.

Evans has described a subspecies, *Th. m. kanara*, from S. India.

451. *Mahathala ameria zistra* Fruhst.

Amblypodia ameria Hewitson, Cat. Lyc. B. M., p. 14 (1862).

Mahathala ameria zistra Seitz, Macrolep. ix, p. 947, fig. 150 b (1926).

Chiengrai, N. Siam; Pak Jong, E. Siam; Sriracha, S. E. Siam.

Distribution: Seitz (*l. c.*) restricts this subspecies to Siam.

The species ranges from India to China, Hainan and Formosa; south to the Malay Peninsula, Sumatra and Java.

452. *Mahathala atkinsoni* (Doh.).

Arhopala atkinsoni Doherty, Journ. As. Soc. Beng. 58 (2), p. 423; Seitz (*Mahathala*), Macrolep. ix, p. 946 (1926).

2 ♂♂ : Chiengmai, Dec. 1921 (*Tennent*); 1 ♂ : Chiengrai, April 1921 (*Godfrey*); N. Siam.

Distribution: A rare species hitherto recorded only from Assam and S. Burma.

Genus AMBLYPODIA*

ANTHELUS-Subgroup.

453. **Amblypodia subfasciata** (Moore).

Nilasera subfasciata Moore, Proc. Zool. Soc. Lond. p. 532 (1883); Seitz (*Amblypodia*), Macrolep. ix, p. 956 (1926).

1 ♂ Bangnara, P. Siam, 29. 7. 23.

Distribution : Burma ; Siam ; Malacca.

454. **Amblypodia anthelus anthelus** Doubl. & Hew.

Amblypodia anthelus Doubleday and Hewitson, Gen. Di. Lep. ii, p. 478, pl. 74, fig. 6 (1852).

Amblypodia anthelus anthelus Evans, Journ. Bomb. N. H. S. xxx, p. 625 (1925).

Amblypodia anunda anthelus Seitz, Macrolep. ix, p. 955 (1926).

1 ♂ Me Song forest, Prae, N. Siam, 20. 4. 16 (Godfrey) ; 1 ♂ : Me Lamoung (2,000 ft.), W. Siam, 17. 4. 20 (Stockley).

Distribution : N. Burma ; Siam ; Malay Peninsula.

Evans has described a subspecies, *A. a. anthea*, which he states occurs from Tavoy to S. Burma.

455. **Amblypodia anarte** Hew.

Amblypodia anarte Hewitson, Cat. Lyc. B. M. p. 5, No. 20 (1862) ; Evans, Journ. Bomb. N. H. S. xxx (3), p. 625 (1925).

Amblypodia agnis anarte Seitz, Macrolep. ix, p. 956 (1926).

Nong Khai Ploi, Sriracha forest, S. E. Siam.

Distribution : Assam to Siam ; Malacca ; Borneo.

Seitz (*l. c.*) treats *anarte* as a subspecies of *agnis* Feld., but both Riley and Evans regard it as a good species.

456. **Amblypodia camdeo sphendale** (Fruhst.).

Amblypodia camdeo Moore, Cat. Lep. Mus. E. I. C. i, p. 41 (1857).

Amblypodia camdeo sphendale Seitz, Macrolep. ix, p. 952 (1926).

* Riley (Entomologist, lv, p. 26, 1922) points out that the generic name *Amblypodia* (Horsfield, 1829) must supersede *Arhopala* (Boisduval, 1832).

1 ♂ : Me Tharn, Me Song forest, Prae, N. Siam, April 1918.

Distribution : Siam ; Annam.

The typical form occurs from Sikkim to N. Burma, and a subspecies *A. c. varro* Fruhst, occurs from the Karen Hills to S. Burma.

457. *Amblypodia dispar* (Ril. & Godf.).

Arhopala dispar Riley and Godfrey, Journ. N. H. S. Siam, iv, p. 183, pl. 6, fig. 5 (1921) ; Evans (*Amblypodia*), Journ. B. N. H. S. xxx, p. 626 (1925).

2 ♂ ♂ : Me Song forest, Prae, N. Siam, April 1918.

Distribution : This species, originally described from Siam, is recorded by Evans from India (Assam-Dawnas).

458. *Amblypodia opalina* (Moore).

Nilasera opalina Moore, Proc. Zool. Soc. p. 531, pl. 49, fig. 1, (1883) ; Riley and Godfrey, Journ. N. H. S. Siam, iv, p. 184, pl. 6, fig. 7, (1921) ; Seitz (*Amblypodia*), Macrolep. ix, p. 952 (1926).

4 ♂ ♂ , 1 ♀ : Me Song forest, Prae, N. Siam, April 1918.

Distribution : Assam to Tavoy ; Siam.

459. *Amblypodia agnis* (Feld.).

Arhopala agnis Felder, Reise Novara, Lep. ii, p. 228 (1865) ; Seitz (*Amblypodia*), Macrolep. ix, p. 955 (1926).

1 ♂ : Klawng Ton, c. 400 m., Satul, P. Siam, 11. 3. 28 (*Kerr*).

Distribution : Burma to Sumatra, Nias, Java, and Mindano.

460. *Amblypodia atosia* Hew.

Amblypodia atosia Hewitson, Ill. Di. Lep. Lyc. p. 9, pl. 2, figs. 8, 9 (1863) ; Seitz, Macrolep. ix, p. 957 (1926) ;

Amblypodia atosia atosia Evans, Journ. B. N. H. S. xxx, p. 627 (1925).

Nong Yai Boo and Hup Bon, Sriracha forest, S. E. Siam (*Godfrey*) ; Ban Natawn, Nakon Sritamarat, P. Siam (*Kerr*).

Distribution : Burma to Sumatra, Borneo and Palawan.

Evans regards *E. a. aricia* Stgr. (Pegu Yoma to Karens) as a good subspecies.

461. ***Amblypodia metamuta* Hew.**

Amblypodia metamuta Hewitson, Ill. Di. Lep. Lyc. p. 13, n. 59 (1863); Seitz, Macrolep. ix, p. 958, fig. 150 B. g. (1926).

Sriracha forest, S. E. Siam (*Godfrey*); Klawng Ton, Satul, P. Siam (*Kerr*).

Distribution: Burma to Borneo, Sumatra, and Billiton.

462. ***Amblypodia moolaiana* (Moore).**

Narathura moolaiana Moore, Proc. Zool. Soc. Lond. p. 835 (1878); Seitz (*Amblypodia*), Macrolep. ix, p. 957, fig. 148 e (1926).

1 ♀ : Me Song forest, Prae, N. Siam, April 1918 (*in Br. Mus.*).

Distribution: Burma; Siam; Malay Peninsula.

ATRAX-Subgroup.

463. ***Amblypodia albopunctata* Hew.**

Amblypodia albopunctata Hewitson, Ill. Di. Lep. p. 14 (1869); Seitz, Macrolep. ix, p. 953, pl. 150 b (1926).

Me Song forest, Prae, N. Siam (*Godfrey*); Eka Klo, W. Siam (*Stockley*); Nang Yawn, Takuapa and Kâpor, Ranawng (Renong), P. Siam (*Kerr*).

Distribution: Burma; Siam.

464. ***Amblypodia mirabella* Doh.**

Amplypodia mirabella Doherty, Journ. As. Soc. Beng., p. 420 (1889).

Amblypodia alitaeus mirabella Seitz, Macrolep. ix, p. 953 (1926); Evans, Journ. Bomb. N. H. S. xxx (3), p. 628 (1925).

Sriracha forest, S. E. Siam (*Godfrey*); Klong Pong Kapo and Thaungyin River, W. Siam (*Stockley*).

Distribution: Tavoy to S. Burma; Siam.

Both Seitz and Evans treat *mirabella* as a subspecies of *alitaeus*, but Riley* regards it as a good species.

465. ***Amblypodia atrax* Hew.**

Amblypodia atrax Hewitson, Cat. Lyc. B. M., p. 13 (1862); Seitz, Macrolep. ix, p. 965, fig. 150 g (1926).

Me Song forest, N. Siam; Sriracha forest, S. E. Siam (*Godfrey*); Pulo Adang, P. Siam (*Kerr*).

Distribution: Shan States to S. Burma; Siam; Malay Peninsula.

* Riley, *in. litt.*, 13. 10. 27.

466. *Amblypodia ariana* Evans.

Amblypodia ariana Evans, Journ. Bomb. N. H. S. xxx, p. 628 (1925).

Me Wong (700 ft.), W. Siam, 16. 2. 24 (*Stockley*).

Distribution: Assam; Burma; Siam; Annam.

467. *Amblypodia rafflesii* (de Nicév.).

Arhopala rafflesii de Nicéville, Butt. Ind. iii, p. 248, fig. 136 (1890); Evans, Journ. Bomb. N. H. S. xxx (3), p. 628 (1925).

Amblypodia pseudomuta rafflesii Seitz, Macrolep. ix, p. 957 (1926).

1 ♂: Nong Yai Boo, Sriracha forest, S. E. Siam, 13. 4. 14 (*Godfrey*); 1 ♂: Ban Udai, Satul, P. Siam, 2. 1. 28 (*Kerr*).

Distribution: S. Burma to Mergui; Siam; Malay Peninsula.

Seitz (*l. c.*) treats *rafflesii* as a subspecies of *pseudomuta* Staud., but says it may be a distinct species.

SILHETENSIS-Subgroup.

468. *Amblypodia silhetensis silhetensis* Hew.

Amblypodia silhetensis Hewitson, Cat. Lyc. B. M. p. 7, pl. 4, figs. 27, 28 (1862); Seitz, Macrolep. ix, p. 955, fig. 148 d (1926).

Amblypodia silhetensis silhetensis Evans, Journ. Bomb. N. H. S. xxx, p. 628 (1925).

Me Song forest, Prae, N. Siam.

Distribution: Sikkim to N. Burma; N. Siam.

Evans recognizes a subspecies, *A. s. arama* de N., from Burma.

469. *Amblypodia adatha adatha* Hew.

Amblypodia adatha Hewitson, Cat. Lyc. Brit. Mus. p. 7 (1862); Evans, Journ. Bomb. N. H. S. xxx (3), p. 629 (1925); Seitz, Macrolep. ix, p. 955 (1926).

1 ♂: Kao Kalakiri (c. 200 m.), Pattani, P. Siam; 3. 4. 28 (*Kerr*).

Distribution: P. Siam to the Malay Peninsula, Malacca, Singapore, the Moluccas, and the Philippines.

Evans has described a subspecies, *A. a. regia*, from Mergui.

PHAENOPS-Subgroup.

470. *Amblypodia agrata* (de Nicev.).

Arhopala agrata de Nicéville, Butt. Ind. p. 251, fig. 137 (1890); Seitz (*Amblypodia*), Macrolep. ix, p. 963, fig. 150 B. g. (1926).

1 ♀ Me Song forest, Prae, N. Siam, April 1916; 1 ♀ April 1918.

Distribution: Assam to the Malay Peninsula.

471. *Amblypodia hypomuta* Hew.

Amblypodia hypomuta Hewitson, Cat. Lyc. B. M., p. 64 (1862); Seitz, Macrolep. ix, p. 958, fig. 148 g (1926).

Khao Luang (2,000 ft.), Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: P. Siam; Malay Peninsula; Malacca; Borneo.

472. *Amblypodia pseudomuta* Staud.

Amblypodia pseudomuta Staudinger, Iris, p. 125 (1889); Seitz, Macrolep. ix, p. 957, fig. 148 e (1926).

1 ♂ Bandon, P. Siam, Sep. 1918.

Distribution: P. Siam; Malacca; Sumatra; Borneo.

473. *Amblypodia inornata* Feld.

Amblypodia inornata Felder, Wien. Ent. Monats, iv, p. 396, n. 6 (1860); Seitz, Macrolep. ix, p. 957, fig. 148 f (1926).

Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*; *Kerr*).

Distribution: P. Siam; Malay Peninsula; (?) Philippines.

474. *Amblypodia aroa* Hew.

Amblypodia aroa Hewitson, Ill. Di. Lep. Lyc., p. 2 (1862); Seitz, Macrolep. ix, p. 956, fig. 148 e (1926).

Amblypodia phaenops aroa Evans, Journ. Bomb. N. H. S. xxx, p. 630 (1925).

Nong Yai Boo, Sriracha forest, S. E. Siam.

Distribution: Seitz (*l. c.*) gives the range of this species as:— “Burmah across Malacca, Sumatra and Borneo to New Guinea, and in Batjan.”

Evans (*l. c.*) treats *aroa* as a subspecies of *phaenops* Feld., which Seitz restricts to the Philippine Islands.

475. *Amblypodia alea alea* Hew.

Amblypodia alea Hewitson, Cat. Lyc. B. M., p. 12 (1862); Seitz, Macrolep. ix, p. 965, fig. 150 B. d (1926).

Amblypodia alea alea Evans Journ. Bomb. N. H. S. xxx, p. 630 (1925).

2 ♂♂ : Me Song forest, Prae, N. Siam, 3. 4. 18 (*Godfrey*); 1 ♂, 1 ♀ Chiengmai, N. Siam (*Tennent*).

Distribution : Burma; Siam.

Evans recognizes a subspecies from the Andamans.

476. *Amblypodia agaba agaba* Hew.

Amblypodia agaba Hewitson, Cat. Lyc. B. M. p. 8 (1862); Seitz, Macrolep. ix, p. 956, pl. 150 c (1926).

Amblypodia agaba agaba Evans, Journ. B. N. H. S. xxx, p. 630 (1925).

Chiengmai, N. Siam (*Tennent*); Hup Bon, Sriracha forest, S. E. Siam (*Godfrey*); Klong Pong Kapo, W. Siam (*Stockley*).

Distribution : Burma; Cochina-China; Siam.

Evans has described a subspecies, *A. a. aurelia*, extending from Manipur to N. India.

477. *Amblypodia hewitsoni* (B.-Baker).

Arhopala hewitsoni Bethune-Baker, Trans. Zool. Soc. p. 132 (1903); Seitz (*Amblypodia*), Macrolep. ix, p. 965, fig. 150 g (1926).

Me Song forest, Prae, N. Siam (*Godfrey*); Um Pang, W. Siam (*Stockley*).

Distribution : N. India to Burma and Siam.

478. *Amblypodia alemon* (de Nicév.).

Arhopala alemon de Nicéville, Journ. Bomb. N. H. S. p. 271 (? 1891); Seitz (*Amblypodia*), Macrolep. ix, p. 965 (1926).

Me Wong, W. Siam (*Stockley*); Chiengmai, N. Siam (*Tennent*).

Distribution : India; Burma; Siam.

CENTAURUS-Subgroup.

479. *Amblypodia centaurus centaurus* (Fabr.).

Papilio centaurus Fabricius, Syst. Ent. p. 520, n. 329 (1775).

Amblypodia centaurus centaurus Seitz, Macrolep. ix, p. 950, fig. 150 a (1926).

Widely spread over Siam.

Distribution : Burma; Siam; Malay Peninsula.

The species ranges from India, through Malaya, to Australia.

480. **Amblypodia amantes amatrix** (de Nicév.).

Arhopala amantes Hewitson, Cat. Lyc. B. M. p. 42 (1862); Seitz (*Amblypodia*), Macrolep. ix, p. 951 (1926).

Arhopala amatrix de Nicéville, Journ. Bomb. N. H. S. p. 370 (1891).

Amblypodia amantes amatrix Evans, Journ. Bomb. N. H. S. xxx, p. 632 (1925).

Me Song forest, Prae, N. Siam.

Distribution : India ; Burma ; Siam.

The species ranges from N. India to the Malay Peninsula and Archipelago.

Seitz makes no mention of *amatrix*.

481. **Amblypodia bazalus** Hew.

Amblypodia bazalus Hewitson, Cat. Lyc. B. M. p. 8 (1862); Seitz, Macrolep. ix, p. 962 (1926).

1 ♀ : Klong Klung, W. Siam, 29. 2. 24 (*Stockley*).

Distribution : Japan ; Sikkim to Siam ; Borneo ; Sumatra.

482. **Amblypodia eumolphus** (Cram.).

Papilio eumolphus Cramer, Pap. Exot. iv, p. 19, pl. 199 (1780).

Distribution : India, through Malaya, to the Philippines and New Guinea.

(a) **Amblypodia eumolphus eumolphus** (Cram.).

Amblypodia eumolphus eumolphus Seitz, Macrolep. ix, p. 959, fig. 149 d (1926).

Me Song forest, Prae, N. Siam (*Godfrey*) ; Chiengmai, N. Siam (*Tennent*).

Distribution : Sikkim to Burma ; N. Siam.

(b) **Amblypodia eumolphus maxwelli** (Dist.).

Narathura maxwelli Distant, Rhop. Malayana, p. 263 (1882-1886).

Amblypodia eumolphus maxwelli Seitz, Macrolep. ix, p. 959 (1926).

Nakon Sritamarat, P. Siam (*Pendlebury* ; *Kerr*).

Distribution : Mergui ; P. Siam ; Malay Peninsula.

RAMA-Subgroup.

483. **Amblypodia rama ramosa** Evans.

Thecla rama Kollar, in Hugel's Kaschmir, iv (2), p. 412 (1848) ; Seitz (*Amblypodia*), Macrolep. ix, p. 964, fig. 150 B. d (1926).

Amblypodia rama ramosa Evans, Journ. Bomb. N. H. S. xxx, p. 634 (1925).

Me Song forest, Prae, N. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*).

Distribution: Assam to Burma; Siam.

The typical form occurs from Kashmir to Sikkim.

484. *Amblypodia paramuta* (de Nicév.).

Panchala paramuta de Nicéville, Journ. As. Soc. Beng. p. 81 (1883); Seitz (*Amblypodia*), Macrolep. ix, p. 964, fig. 150 B. b (1926).

1 ♂ : Me Lamoung (2,000 ft.), W. Siam, 19. 4. 20 (*Stockley*).

Distribution: Sikkim to Siam.

AGELASTUS-Subgroup.

485. *Amblypodia asopia* Hew. (= *tounguva* Sm.).

Amblypodia asopia Hewitson, Ill. Di. Lep. Lyc. p. 146 (1869); Seitz, Macrolep ix, p. 965, pl. 150 g (1926).

Chiengmai, N. Siam (*Tennent*); Sen Tor (Nam Pat), N. E. Siam (*Godfrey*).

Distribution: Assam to Siam and Annam.

486. *Amblypodia antinuta* (Feld.).

Arhopala antimuta Felder, Nov. Reise, Lep. ii, p. 233 (1865); Seitz (*Amblypodia*), Macrolep. ix, p. 958, fig. 150 c (1926).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Burma to the Malay Peninsula; (?) Borneo.

487. *Amblypodia perimuta* perimuta Moore.

Amblypodia perimuta Moore, Cat. Lep. Mus. E. I. C., p. 42 (1857); Seitz, Macrolep. ix, p. 965, fig. 148 g (1926).

Amplypodia perimuta perimuta Evans, Journ. Bomb. N. H. S. xxx, p. 635 (1925).

Me Song forest, Prae, N. Siam.

Distribution: Sikkim to N. Burma; N. Siam.

Evan's (*l. c.*) recognizes a subspecies from S. Burma and Mergui.

488. *Amblypodia duessa* (Doh.).

Arhopala duessa Doherty, Journ. As. Soc. Beng., p. 41 (1889).

Amblypodia epimete duessa Seitz, Macrolep. ix, p. 965 (1926).

1 ♀ : Me Song forest, Prae, N. Siam, April 1918.

Distribution : Dawna Hills to S. Burma ; N. Siam.

Seitz treats *duessa* as a subspecies of *epimete* Staud., which he refers to Palawan.

GANESA-Subgroup.

489. **Amblypodia parangesa parangesa** de Nicév.

Amblypodia parangesa de Nicéville, Journ. As. Soc. Beng., p. 63 (1882).

Amblypodia parangesa parangesa Seitz, Macrolep. ix, p. 966, fig. 147 f (1926).

1 ♀ : Me Song forest, Prae, N. Siam, April 1918.

Distribution : Kumaon to Bhutan ; Siam.

A subspecies, *A. p. zephyretta* Doh. is found in Assam.

490. **Amblypodia ammonides** Doh.

Amblypodia ammonides Doherty, Journ. As. Soc. Beng. 60 (2), p. 34.

Amblypodia ammon ammonides Seitz, Macrolep. ix, p. 966 (1926).

1 ♂ : Wang Sai, Songkla, P. Siam, 20. 7. 28 (*Kerr*).

Distribution : Burma ; Siam.

Seitz (*l. c.*) treats *ammonides* as a subspecies of *ammon* Hew. and refers it to Tenasserim.

491. **Amblypodia birmana birmana** (Moore).

Panchala birmana Moore, Proc. Zool. Soc. Lond., p. 531 (1883).

Amblypodia birmana birmana Seitz, Macrolep. ix, p. 966 (1926).

2 ♂ ♂ , 1 ♀ : Me Song forest, Prae, N. Siam, April 1918.

Distribution : Assam to Siam.

Seitz (*l. c.*) recognizes a subspecies from Hongkong.

492. **Amblypodia aberrans** (de Nicév.).

Acesina aberrans de Nicéville, Journ. As. Soc. Beng. p. 279 (1888) ; Evans (*Amblypodia*), Journ. Bomb. N. H. S. xxx, p. 637 (1925).

Amblypodia birmana aberrans Seitz, Macrolep. ix, p. 966, fig. 150 B. d (1926).

1 ♂ : Me Song forest, Prae, N. Siam (*Godfrey*) ; 1 ♀ : Chiengmai, N. Siam (*Tennent*).

Distribution : Assam to Siam.

493. *Amblypodia abseus indicus* Ril.

Amblypodia abseus Hewitson, Cat. Lyc. B. M. n. 40 (1862); Seitz, Macrolep. ix, p. 962 (1926).

Amblypodia abseus indicus Riley, Spolia Zeylan. xii, pt. 46, p. 326 (1923).

Distribution: Sikkim to Siam; Annam.

The typical form occurs from Burma to Malacca, and a subspecies is found in Ceylon.

APIDANUS-Subgroup.

494. *Amblypodia diardi diardi* Hew.

Amblypodia diardi Hewitson, Cat. Lyc. B. M. p. 9. n. 43 (1862).

Amblypodia diardi diardi Seitz, Macrolep. ix, p. 961, fig. 150 B. g (1926).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Sikkim to Burma; Siam.

The species ranges from India, through Malaya, to the Philippines.

495. *Amblypodia fulgida fulgida* Hew.

Amblypodia fulgida Hewitson, Ill. Diurn. Lep. Lyc. p. ii (1863).

Amblypodia fulgida fulgida Seitz, Macrolep. ix, p. 960, fig. 150 B. g (1926).

1 ♂ : Chiengmai, N. Siam, Jan. 1921 (*Tennent*).

Distribution: Sikkim to the Shan States; Siam.

The species extends through Malaya to the Philippines.

496. *Amblypodia apidanus apidanus* (Cram.).

Papilio apidanus Cramer, Pap. Exot. ii, pl. 137, figs. f. g. (1777).

Amblypodia apidanus apidanus Seitz, Macrolep. ix, p. 960, fig. 149 e (1926).

Prae, N. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam.

Distribution: Burma; Siam.

The species ranges from N. India, through Malaya, to the Philippines.

497. *Amblypodia fulla ignara* (Ril. & Godf.).

Amblypodia (? *Narathura*) *fulla* var. *andamanica* Wood-Mason and de Nicéville, Proc. As. Soc. Beng. p. 143 (1881).

Arhopala andamanica ignara Riley and Godfrey, Journ. Nat. Hist. Soc. Siam, iv, p. 185, pl. 6, fig. 6, ♂ (1921).

Amblypodia fulla ignara Evans, Journ. Bomb. N. H. S. xxx, p. 639 (1925).

Amblypodia fulla, Seitz, Macrolep. ix, p. 967 (1927).

Me Song forest, Prae, N. Siam; Sen Tor (Nam Pat), N. E. Siam.

Distribution: Sikkim to Burma; Siam.

Other subspecies are found in the Andamans and Amboina.

SURENDRA-Group.

498. *Surendra quercetorum quercetorum* Moore.

Surendra quercetorum Moore, Cat. Lep. Mus. E. I. C. p. 42 (1857).

Surendra quercetorum quercetorum, Seitz, Macrolep. ix, p. 943, fig. 149 a (1926).

Me Song forest, Prae, and Chiengrai, N. Siam; Pak Jong, E. Siam.

Distribution: N. India to Burma; Siam.

Subspecies are found in Tonkin, S. India, and Ceylon.

499. *Surendra todara distorta* (de Nicév.).

Surendra todara Moore, Proc. Zool. Soc. Lond. p. 530 (1883); Seitz, Macrolep. ix, p. 107 (1926).

Rapala distorta de Nicéville, Proc. Zool. Soc. Lond. p. 461 (1877).

Surendra todara distorta Evans, Journ. Bomb. N. H. S. xxx, p. 756 (1925).

1 ♂: Nong Khai Ploi, Sriracha forest, S. E. Siam, April 1914.

Distribution: Sikkim to Shan States; Siam.

Evans (*l. c.*) recognizes a subspecies from the Karen and Dawna Hills.

LOXURA-Group.

500. *Loxura atymnus continentalis* Fruhst.

Papilio atymnus Cramer, Pap. Exot. iv, p. 82 (1780); Evans (*Loxura*), Journ. Bomb. N. H. S. xxx, p. 757 (1925).

Loxura atymnus continentalis Fruhstorfer, Berl. Ent. Zeit. lvi, p. 221 (1912); Seitz, Macrolep. ix, p. 996 (1926).

Widely spread over Siam.

Distribution: Seitz (*l. c.*) refers *continentalis* to "the Himalayan countries as far as Siam and Burma." Evans, however, does not recognize it as an Indian form.

The species ranges from India and China to Malaya.

501. *Yasoda tripunctata* (Hew.).

Loxura tripunctata Hewitson, Ill. Di. Le p. p. 26 (1863); Seitz (*Yasoda*), Macrolep. ix, p. 995, fig. 157 C. d. (1926).

This record is based on a single specimen obtained by Dr. Erick Haase at Chantabun, S. E. Siam, in 1891.

Distribution: Sikkim to Burma; Siam; Perak.

502. *Yasoda pita* (Horsf.).

Loxura pita Horsfield, Cat. Lep. E. I. C. p. 122 (1829); Seitz (*Yasoda*) Macrolep. ix, p. 995 (1926).

Khao Luang (2,000-3,000 ft.), Nakom Sritamarat (*Pendlebury*); Ban Sai Kao and Khao Kalakiri, Pattani (*Kerr*)—all in P. Siam.

Distribution: P. Siam; Malay Peninsula; Sumatra; Java.

503. *Drina donina* (Hew.).

Myrina donina Hewitson, Ill. Di. Lep., p. 39 (1865); Seitz (*Drina*), Macrolep. ix, p. 987, fig. 160 a (1926).

Me Song forest, Prae, N. Siam and Sriracha forest, S. E. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*); Klong Suan Mak, W. Siam (*Stockley*).

Distribution: Burma; Siam; Malacca.

504. *Drina maneia*. (Hew.).

Myrina maneia Hewitson, Ill. Di. Lep., p. 29, pl. xii, figs. 14, 15 (1863); Seitz (*Drina*), Macrolep. ix, p. 987 (1926).

1 ♂: Kao Keo Range (c. 600 m.), Satul, P. Siam, 12. 3. 28 (*Kerr*).

Distribution: Mergui; P. Siam; Singapore; Malacca; Borneo.

SPINDASIS-Group.

505. *Spindasis vulcanus tavoyana* Evans.

Papilio vulcanus Fabricius, Syst. Ent. p. 519 (1775); Seitz (*Aphnaeus*), Macrolep. ix, p. 938 (1926).

Spindasis vulcanus tavoyana Evans, Journ. Bomb. N. H. S. xxx, p. 758 (1925).

Phar Mieng, near Chiengrai, N. Siam ; Sriracha, S. E. Siam.

Distribution : Tavoy ; Siam.

The typical form occurs in India and subspecies are found in Ceylon and Java.

506. *Spindasis maximus* (Elw.).

Aphnaeus maximus Elwes, Proc. Zool. Soc. Lond., p. 637 (1892); Evans (*Spindasis*), Journ. Bomb. N. H. S. xxx, p. 759 (1925).

Aphnaeus ictis maximus Seitz, Macrolep. ix, p. 939 (1926).

1 ♂ : Chiengmai district, N. Siam, Jan. 1921 (*Tennent*).

Distribution : Burma ; Siam.

Seitz (*l. c.*) treats *maximus* as a subspecies of *ictis* Hew., but both Riley* and Evans regard it as a good species.

507. *Spindasis syama peguanus* (Moore).

Amblypodia syama Horsfield, Cat. Lep. Mus. E. I. C. p. 107 (1829).

Spindasis syama peguanus Evans, Journ. B. N. H. S. xxx, p. 759 (1925).

Aphnaeus lohita peguanus Seitz, Macrolep. ix, p. 937 (1926).

Me Song forest, Prae, and Chiengrai, N. Siam ; Sriracha, S. E. Siam ; Bangkok, C. Siam.

Distribution : Sikkim to Burma ; Siam.

The species ranges from India to the Philippines.

508. *Spindasis lohita batina* Fruhst.

Amblypodia lohita Horsfield, Cat. Lep. Mus. E. I. C. p. 186 (1829).

Aphnaeus lohita batina Fruhstorfer, Berl. Ent. Zeit., p. 218 (1912) ; Seitz, Macrolep. ix, p. 937, fig. 157 a (1926).

Chiengmai, N. Siam (*Tennent*) ; Pak Jong, E. Siam and Bangkok, C. Siam (*Godfrey*).

Distribution : Siam ; S. Annam.

The species ranges from India and China to the Philippines.

509. *Dacalana penicilligera* (de Nicév.).

Arrhenothrix penicilligera de Nicéville, Butt. Ind. iii, p. 337 (1890) ;

Evans (*Dacalana*), Journ. Bomb. N. H. S. xxx, p. 760 (1925).

Dacalana vidura penicilligera Seitz, Macrolep. ix, p. 970, fig. 155 b (1926).

Me Song forest, Prae, N. Siam.

* Riley, *in litt.* 9. 8. 27.

Distribution: Bhutan and Assam to Borneo and Java.

This species was placed in *Arrhenothrix* by de Nicéville on account of having only 4 branches to the s. c. vein (there are 5 in *Dacalana*), but this character is highly variable and it is most probable that *penicilligera* is merely a subspecies of *D. vidura* Horsf. Seitz (*l. c.*) presumes that it is "only a large form" of that species.

510. *Tajuria jangala ravata* (Moore).

Amblypodia jangala Horsfield, Cat. Lep. Mus. E. I. C. p. 113 (1829).

Myrina ravata Moore, Proc. Zool. Soc. Lond. p. 776 (1865).

Remelana jangala ravata Seitz, Macrolep. ix, p. 977, fig. 146 c (1926).

Me Song forest, Prae, N. Siam; Sriracha forest, S. E. Siam.

Distribution: Sikkim to Burma; Siam.

The species extends to the Philippines.

511. *Tajuria cippus maxentius* Fruhst.

Hesperia cippus Fabricius, Ent. Syst. Suppl. v, p. 429 (1798).

Tajuria cippus maxentius Fruhstorfer, Berl. Ent. Zeit. xlvi, p. 210 (1912); Seitz, Macrolep. ix, p. 972 (1926).

Muak Lek, E. Siam; Bangkok, C. Siam.

Distribution: Siam; Malay Peninsula; Sumatra.

The typical form occurs in India and Burma, and subspecies are found in Hainan, Ceylon and Java.

512. *Tajuria jalindra* (Horsf.).

Amblypodia jalindra Horsfield, Cat. Lep. Mus. E. I. C., p. 109 (1829); Seitz (*Tajuria*), Macrolep. ix, p. 973 (1926).

1 ♂: Klawng Kampuam, Ranawng (Renong), P. Siam, 4. 2. 29 (*Kerr*).

The species ranges from India to Java, Sumatra, and Nias.

JACOONA-Group.

513. *Jacoona anasuja* Feld.

Jacoona anasuja Felder, Novara Reise Lep. ii, p. 237 (1865).

Jacoona anasuja anasuja Seitz, Macrolep. ix, p. 992, fig. 146 f. (1926).

1 ♂: Chiengmai district, N. Siam, Jan. 1921 (*Tennent*); 1 ♂: Ta Samet, Nakon Sritamarat, P. Siam, 27. 1. 28 (*Kerr*).

Distribution: Burma; Siam; Malay Peninsula.

514. **Neucheritra amrita** (Feld.).

Myrina amrita Felder, Wien. Ent. Monats. iv, p. 395 (1860); Seitz (*Neucheritra*), Macrolep. ix, p. 991, fig. 159 a (1926).

1 ♂ : Ta Samet, Nakon Sritamarat, P. Siam, 28. 1. 28 (*Kerr*).

Distribution : Burma ; Siam ; Malay Peninsula ; Nias ; Billiton ; Sumatra ; Borneo.

Moulton* treats *theodora* Druce (which Seitz refers to Borneo) as a synonym of *amrita*.

515. **Purlisa gigantea gigantea** (Dist.).

Joalus (Purlisa) gigantea Distant, Ent. Mo. Mag. xvii, p. 245 (1881).

Tajuria gigantea gigantea Seitz, Macrolep. ix, p. 973, fig. 155 f. (1926).

1 ♂ : Koh Pa Nga, Chumporn, P. Siam, August 1919.

Distribution : S. Burma ; Siam ; Penang (*Type locality*).

A very rare species of which only a few examples are known.

Seitz (*l. c.*) separates the Bornean form as *P. g. borneana* Fruhst.

516. **Suasa lisides** (Hew.).

Myrina lisides Hewitson, Ill. Di. Lep. p. 33 (1863); Seitz (*Suasa*) Macrolep. ix, p. 980, fig. 157 h (1926).

1 ♀ : Renong, P. Siam (*Adams Coll., British Museum*).

Distribution : Assam to Burma ; Siam ; Perak.

MARMESSUS-Group.

517. **Cheritrella truncipennis** de Nicév.

Cheritrella truncipennis de Nicéville, Proc. Zool. Soc. p. 456 (1887) ; Seitz, Macrolep. ix, p. 991, fig. 159 a (1926).

1 ♂ : Chiengrai district, N. Siam, Jan. 1921 (*Tennent*).

Distribution : Sikkim ; Assam ; N. Burma ; N. Siam.

518. **Neomyrina hiemalis** (Godm. & Salv.).

Myrina hiemalis Godman and Salvin, Proc. Zool. Soc. p. 640 (1878) ; Seitz (*Neomyrina*), Macrolep. ix, p. 991, fig. 159 b (1926).

Me Song forest, Prae, N. Siam ; Sriracha forest, S. E. Siam.

Distribution : Burma ; Siam ; Malacca ; Sumatra.

* Moulton, Journ. Straits Branch Roy. As. Soc. No. 60, p. 158 (1912).

519. ***Cheritra freja freja* (Fabr.).**

Hesperia freja Fabricius, Ent. Syst. iii, p. 263 (1793).

Cheritra freja freja Seitz, Macrolep. ix, p. 993, figs. 146 g, 159 b (1926).

Me Song forest and Doi Chang (4,500 ft.), N. Siam; Sriracha forest, S. E. Siam (*Godfrey*); Me Ping valley, W. Siam (*Stockley*).

Distribution: N. India to Burma; Tonkin; Siam.

The species ranges from India to the Malay Peninsula, Sumatra, and Java.

520. ***Ticherra acte* (Moore).**

Myrina acte Moore, Cat. Lep. Mus. E. I. C., p. 47 (1857); Seitz (*Ticherra*), Macrolep. ix, p. 994, figs. 146 g, 158 h (1926).

2 ♂♂, 3 ♀♀: Me Song forest, Prae, N. Siam, April 1916.

Distribution: Sikkim to Burma; Siam; Sumatra.

521. ***Biduanda cyara* (Hew.).**

Myrina cyara Hewitson, Ill. Di. Lep. Suppl. p. 26 (1878); Riley and Godfrey (*Biduanda*), Journ. N. H. S. Siam, iv, p. 187 (1921); Seitz (*Biduanda*), Macrolep. ix, p. 988, fig. 146 B. d (1926).

Biduanda melisa cyara Evans, Journ. Bomb. N. H. S. xxx, p. 766 (1925).

Me Song forest, Prae, N. Siam.

Distribution: Sikkim to N. Burma; N. Siam.

Evans (*l. c.*) treats *cyara* as a subspecies of *melisa* Hew., but Riley and Godfrey (*l. c.*) regard it as a good species.

522. ***Biduanda melisa* (Hew.).**

Myrina melisa Hewitson, Ill. Di. Lep. Suppl. p. 6 (1869); Seitz (*Biduanda*), Macrolep. ix, p. 988, fig. 146 B. d (1926).

Sriracha forest, S. E. Siam.

Distribution: S. Burma; S. E. Siam.

523. ***Biduanda thesmia* (Hew.).**

Myrina thesmia Hewitson, Ill. Di. Lep. Lyc. p. 32 (1863).

Distribution: Burma to the Malay Peninsula, Sumatra, Nias, Borneo and Palawan.

(a) ***Biduanda thesmia fabricii* (Moore).**

Drupadia fabricii Moore, Journ. Linn. Soc. Zool. p. 32 (1884).

Biduanda thesmia fabricii Seitz, Macrolep. ix, p. 987 (1926).

2 ♂♂, 1 ♀ : Me Song forest, Prae, N. Siam, April 1918; 1 ♀ April 1916.

Distribution : Burma ; N. Siam.

(b) **Biduanda thesmia thesmia** (Hew.).

Biduanda thesmia thesmia Seitz, op. cit., p. 987.

Khao Ram and Khao Luang, P. Siam (*Pendlebury*).

Distribution : Mergui ; Siam ; Malay Peninsula ; Sumatra.

524. Marmessus lisias boisduvali Moore.

Papilio lisias Fabricius, Mant. Ins. ii, p. 65 (1787).

Marmessus lisias boisduvali Seitz, Macrolep. ix, p. 989, fig. 159 c. (1926).

Me Song forest, Prae, N. Siam and Sriracha forest, S. E. Siam (*Godfrey*) ; Ban Samui, W. Siam (*Stockley*).

Distribution : Burma ; Siam ; Annam.

The species ranges from Burma to Sumatra and Borneo.

525. Marmessus moorei (Dist.).

Sithon moorei Distant, Ann. Mag. N. H. (5) x, p. 246 (1882) ; Seitz (*Marmessus*), Macrolep. ix, p. 989 (1926).

Chiengrai, N. Siam (*Godfrey*) ; Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : Siam to the Malay Peninsula, Borneo and Sumatra.

526. Eooxylides tharis (Hübner).

Oxylides tharis Hübner, Zutr. exot. Schmett. figs. 883, 884 (1857) ; Seitz (*Eooxylides*), Macrolep. ix, p. 992 (1926).

1 ♂ : Ban Natawn, Nakon Sritamarat, P. Siam, 10. 5. 28 (*Kerr*).

Distribution : India to the Malay Peninsula, Sumatra, Nias, Billiton, Borneo, Sumatra, and Java.

527. Thamala miniata Moore.

Thamala miniata Moore, Proc. Zool. Soc. Lond. p. 834 (1878) ; Seitz, Macrolep. ix, p. 986, figs. 158 d, 156 g (1926).

Khao Ram and Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : S. Burma ; Mergui ; P. Siam ; Malay Peninsula.

528. **Thamala marciana** (Hew.).

Myrina marciana Hewitson, Ill. Di. Lep. Lyc. p. 34 (1863); Seitz (*Thamala*), Macrolep. ix, p. 986, fig. 158 e (1926).

Khao Ram, Nakon Sritamarat, P. Siam (*Pendlebury*); Kao Pawta Luang Keo, c. 1,300 m., Ranawng (Renong), P. Siam (*Kerr*).

Distribution: P. Siam; Malacca; Sumatra; Borneo.

HORAGA-Group.

529. **Horaga achaja** Fruhst.

Horaga achaja Fruhstorfer, Berl. Ent. Zeit. lvi, p. 233 (1912); Riley and Godfrey, Entomol. lviii, p. 141, pl. 3, fig. 1, ♀; Seitz, Macrolep. ix, p. 982, fig. 158 a (1926).

1 ♂ : Hinlap, E. Siam, Jan. 1900 (*Fruhstorfer*); 6 ♂♂, 4 ♀♀ : Me Song forest, Prae, N. Siam, April 1916 (*Godfrey*).

Distribution: Confined to Siam.

A very distinct species described by Fruhstorfer from a single ♂ which he obtained at Hinlap on the Bangkok-Korat line.

530. **Horaga halba** Dist.

Horaga halba Distant, Rhop. Malay. p. 460 (1882-1886).

Horaga onyx halba Seitz, Macrolep. ix, p. 982 (1926).

1 ♂ : Kao Pawta Luang Keo, c. 1,300 m., Ranawng (Renong), P. Siam, 2, 2, 29 (*Kerr*).

Distribution: Mergui; P. Siam; Malacca; Sumatra.

CATAPOECILMA-Group.

531. **Catapoecilma elegans** (Druce).

Hypochnysops elegans Druce, Proc. Zool. Soc., p. 350 (1873).

Distribution: India; Ceylon; Siam; Malay Peninsula; Nias; Sumatra; Borneo; Java; Philippines.

(a) **Catapoecilma elegans major** (Druce).

Catapoecilma elegans major Seitz, Macrolep. ix, p. 983 (1926).

Me Song forest, Prae, N. Siam.

Distribution: Mussoorie to Burma; N. Siam.

(b) **Catapoecilma elegans emas** Fruhst.

Catapoecilma elegans emas Seitz., op. cit., p. 983.

1 ♂ : Kao Samui, P. Siam, May 1928 (*Kerr*).

Distribution : P. Siam ; Malacca ; Sumatra.

532. *Catapoecilma subochracea* Elw.

Catapoecilma subochracea Elwes, Proc. Zool. Soc. Lond. p. 640 (1892) ; Seitz, Macrolep. ix, p. 983 (1926).

1 ♂ , 1 ♀ : Chiengmai, N. Siam, Feb. 1921 (*Tennent*).

Distribution : Naga Hills to S. Burma ; N. Siam.

HYPOLYCAENA-Group.

533. *Chliaria othona othona* (Hew.).

Hypolycaena othona Hewitson, Ill. Di. Lep. p. 50 (1865).

Chliaria othona othona Seitz, Macrolep. ix, p. 980 (1926).

Me Song forest, Prae, N. Siam (*Godfrey*) ; Me Wong, W. Siam (*Stockley*).

Distribution : Sikkim to Burma ; Andamans ; Siam.

Seitz (*l. c.*) recognizes a subspecies, *Ch. o. matiana* Fruhst., from Tonkin.

534. *Chliaria merguia watsoni* Swinh.

Chliaria merguia Doherty, Journ. As. Soc. Beng. 58 (2), p. 427 ; Seitz, Macrolep. ix, p. 981 (1926).

Chliaria merguia watsoni Evans, Journ. Bomb. N. H. S. xxx, p. 769 (1925).

1 ♂ , 1 ♀ : Me Wong (2,500 ft.), 12. 4. 20 ; 1 ♂ Klong Klung, 29. 2. 24—W. Siam (*Stockley*).

Distribution : Burma ; Siam.

The species ranges from Burma to Sumatra, Borneo, and Java.

535. *Hypolycaena thecloides thecloides* (Feld.).

Myrina thecloides Felder, Wien. Ent. Monats. iv, p. 395 (1860).

Hypolycaena thecloides thecloides Seitz, Macrolep. ix, p. 984, fig. 158 B. c (1926).

1 ♂ : Khanduli, P. Siam, Sep. 1918.

Distribution : S. Burma ; Siam ; Malacca ; Sumatra ; Java ; Borneo.

Seitz recognizes as good subspecies :—*vardura* Fruhst. from “Isle of Siargao near Borneo” ; *extensa* Fruhst., from Engano ; and

philippina Staud. from Palawan. Evans has described a subspecies, *nicobarica*, from the Nicobars.

536. *Hypolycaena phemis* H. H. Druce.

Hypolycaena phemis H. H. Druce, Proc. Zool. Soc. p. 605 (1895).

Chliaria merguia phemis Seitz, Macrolep. ix, p. 981, fig. 147 b (1926).

Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: P. Siam; Borneo.

Seitz (*l. c.*) treats *phemis* as a subspecies of *Ch. merguia* Doh. and remarks that Fruhstorfer placed it under *Ch. amabilis* de Nicév.

Riley* informs me that an examination of the type of *phemis* proves that it is undoubtedly correctly referred to *Hypolycaena*, and is not a *Chliaria*.

537. *Hypolycaena erylus himavantus* Fruhst.

Polyommatus erylus Godart, Ency. Méth. ix, p. 633 (1823).

Hypolycaena erylus himavantus Seitz, Macrolep. ix, p. 984, figs. 148 a, 158 c. (1926).

Widely spread over Siam.

Distribution: Sikkim to Burma; Tonkin; Siam; Annam.

The species ranges from India to Sumatra, Borneo, Java, and the Philippines.

538. *Zeltus etolus etolus* (Fabr.).

Papilio etolus Fabricius, Mant. Ins. ii, p. 66 (1787).

Zeltus etolus etolus Seitz, Macrolep. ix, p. 990, fig. 146 g (1926).

Prae, N. Siam; Muak Lek, E. Siam; Sriracha, S. E. Siam (*Godfrey*); Klong Pong Kapo, W. Siam (*Stockley*).

Distribution: Sikkim to Burma; Tonkin; Siam.

The species ranges from Sikkim to Malacca, Sumatra, Borneo, and Java.

ARTIPE-Group.

539. *Deudorix epijarbas amatius* Fruhst.

Dipsas epijarbas Moore, Cat. Lep. Mus. E. I. C. i, p. 32 (1857).

*Riley, *in litt.*, 22. 9. 27.

Deudorix epijarbas amatius Fruhstorfer, Berl. Ent. Zeit. lvi, p. 265 (1912); Seitz, Macrolep. ix, p. 999 (1926).

Me Song forest, Prae; Khun Than (4,000 ft.)—N. Siam.

Distribution: Sikkim to Burma; Andamans and Nicobars; Tonkin; Siam.

The species ranges from N. India to Formosa and the Philippines, south to Malaya.

540. *Virachola perse perse* (Hew.).

Deudorix perse Hewitson, Ill. Di. Lep. p. 18 (1863).

Deudorix perse perse Seitz, Macrolep. ix, p. 999, pl. 161 b (1926).

Hue Tak So, W. Siam (*Stockley*).

Distribution: Sikkim to N. Burma; Siam.

Subspecies are found in S. Burma and the Andamans, and in the Philippines.

541. *Rapala abnormis* Elwes.

Rapala abnormis Elwes, Proc. Zool. Soc. Lond. p. 642, pl. xliv, fig. 2, ♂ (1892); Seitz, Macrolep. ix, p. 1007 (1926).

1 ♂: Kao Pawta Luang Keo, c. 900 m., Ranawng (Renong) P. Siam, 21. I. 29 (*Kerr*).

Distribution: Karen Hills to S. Burma; P. Siam; Borneo; Sumatra; Java.

This species is apparently rare.

542. *Rapala suffusa suffusa* (Moore).

Deudorix suffusa Moore, Proc. Zool. Soc. Lond., p. 834 (1878).

Rapala suffusa suffusa Evans, Journ. Bomb. N. H. S. xxx, p. 771 (1925).

Rapala xenophon suffusa Seitz, Macrolep. ix, p. 1002 (1926).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Assam to Burma; Siam; (?) Malacca.

Evans (*l. c.*) recognizes a subspecies from the Andaman Islands.

Seitz (*l. c.*) incorrectly places *suffusa* under *xenophon* Fabr.

543. *Rapala sphinx* (Fabr.).

Papilio sphinx Fabricius, Syst. Ent. p. 520 (1775); Seitz (*Rapala*), Macrolep. ix, p. 1005, fig. 160 e (1926).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Assam to Borneo, Sumatra Java and the Philippines.

544. *Rapala rhoecus* de Nicév.

Rapala rhoecus de Nicéville, Journ. Bomb. N. H. S. p. 319, pl. P, fig. 47, ♂ (1895); op. cit., p. 182, pl. 7, fig. 40, ♀ (1896); Riley and Godfrey, Journ. N. H. S. Siam, iv, p. 186 (1921).

Rapala sphinx rhaecus Evans, Journ. Bomb. N. H. S. xxx, p. 772 (1925); Seitz, Macrolep. ix, p. 1005 (1926).

1 ♂: Me Song forest, Prae, N. Siam.

Distribution: Evans (*l. c.*) gives the range of *rhoecus* as "S. Shan States—S. Burma." There are in the British Museum four males: 2 from Sumatra, 1 from Moulmein, Tenasserim, and 1 from Bhutan (*G. C. Dudgeon*).

Both Seitz and Evans treat *rhoecus* as a subspecies of *sphinx* Fabr., but Riley and Godfrey (*l. c.*) regard it as a very distinct species. The male cannot be mistaken as it has a small very clearly defined sex-mark, triangular in shape, occupying the bases of areas 2 and 3; this is entirely absent in the male of *sphinx*.

545. *Rapala varuna orseis* Hew.

Thecla varuna Horsfield, Cat. Lep. Mus. E. I. C. p. 91 (1829).

Rapala varuna orseis Seitz, Macrolep. ix, p. 1004, fig. 146 e (1926).

Me Song forest, Prae, N. Siam; Pak Jong, E. Siam.

Distribution: Sikkim to Burma; Andamans; Siam.

The species ranges from N. India to Formosa, south to the Malay Archipelago.

546. *Rapala schistacea* (Moore).

Deudorix schistacea Moore, Proc. Zool. Soc. Lond. p. 140 (1879); Seitz (*Rapala*), Macrolep. ix, p. 1005, fig. 146 a (1926).

Me Song forest, Prae, N. Siam; Sriracha forest, S. E. Siam.

Distribution: India; Ceylon; Andaman Islands; Siam; Borneo; Sumatra; Java.

547. *Rapala scintilla* de Nicév.

Rapala scintilla de Nicéville, Butt. Ind. iii, p. 461 (1890); Moulton, Journ. Straits Branch Roy. As. Soc. No. 60, p. 171 (1912); Seitz, Macrolep. ix, p. 1005, fig. 160 d (1926).

Pak Jong, E. Siam.

Distribution: Sikkim to Burma; Siam; Sumatra; Borneo.

548. *Rapala pheretimus petosiris* (Hew.).

Deudorix pheretima Hewitson, Ill. Di. Lep. p. 21 (1863).

Deudorix petosiris Hewitson, op. cit., p. 22; Seitz (*Rapala*), Macrolep. ix, p. 1003, fig. 146 b (1926).

Rapala pheretimus petosiris Evans, Journ. Bomb. N. H. S. xxx, p. 773 (1925).

Me Song forest, Prae, N. Siam; Muak Lek and Pak Jong, E. Siam; Bangkok, C. Siam.

Distribution: Sikkim to Burma; Siam.

The typical form occurs in S. Mergui.

549. *Rapala dieneces dieneces* Hew.

Rapala dieneces Hewitson, Hew. Ill. Di. Lep. Suppl. p. 31.

Rapala dieneces dieneces Evans, Journ. Bomb. N. H. S. Vol. xxx (3), p. 773 (1925).

Rapala xenophon dieneces Seitz, Macrolep. ix, p. 1002 (1926).

Pak Jong, E. Siam.

Distribution: Bengal; Assam to Burma; Siam; (?) Malacca.

Evans (*l. c.*) recognizes *R. d. intermedia* Staud., from the Andamans, as a good subspecies.

Seitz (*l. c.*) treats *dieneces* as a subspecies of *xenophon*.

550. *Rapala jarbas* (Fabr.).

Papilio jarbas Fabricius, Mant. Ins. ii, p. 68 (1787); Evans (*Rapala*), Journ. Bomb. N. H. S. xxx, p. 773 (1925).

Rapala melampus jarbas Seitz, Macrolep. ix, p. 1001 (1926).

Prae, N. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam.

Distribution: Sikkim to Burma; Siam; Annam.

Seitz (*l. c.*) treats *jarbas* as a subspecies of *melampus* Cram., but remarks that the two forms:—"are both so extraordinarily similar that it appears almost impossible to separate them."

551. *Sinthusa nasaka amba* Kirby.

Thecla nasaka Horsfield, Cat. Lep. Mus. E. I. C. p. 91 (1829); Seitz (*Sinthusa*), Macrolep. ix, p. 979, fig. 157 f (1926).

Sinthusa nasaka amba Evans, Journ. B. N. H. S. xxx, p. 774 (1925).

Me Song forest, Prae, N. Siam ; Sriracha forest, S. E. Siam.

Distribution : Sikkim to Burma ; Siam.

The species ranges from N. India to Borneo, Sumatra and Java.

552. Bindahara phocides ines Swinh.

Hesperia phocides Fabricius, Ent. Syst. iii, p. 282 (1793) ; Seitz (*Bindahara*) Macrolep. ix, p. 994 (1926).

Bindahara ines Swinhoe, Lep. Ind. ix, p. 27 (1911).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution : Burma ; Siam ; Borneo ; Sumatra.

The species ranges from the Himalayas to Australia.

553. Araotes lapithis lapithis (Moore).

Myrina lapithis Moore, Cat. Lep. Mus. E. I. C. i., p. 48 (1857).

Araotes lapithis lapithis Seitz, Macrolep. ix, p. 978, fig. 157 e (1926).

Me Song forest, Prae, and Doi Chang (2,200 ft.)—N. Siam.

Distribution : Sikkim to Burma ; Siam.

The species ranges from India to the Malay Peninsula, Borneo, Sumatra, and Java.

LIPHYRINAE.

554. Liphyra brassolis Westw.

Liphyra brassolis Westwood, Proc. Ent. Soc. Lond. (3) ii, p. 31 (1864) ; Grünberg, in Seitz, Macrolep. ix, p. 802 (1915).

Bangkok (*Godfrey* ; *Ladell*).

Distribution : N. India to Australia.

HESPERIIDAE.

555. Hasora lizetta anura de Nicév.

Ismene lizetta Plötz, Stett. Ent. Zeit. xlv, p. 59 (1884).

Hasora anura de Nicéville, Journ. Bomb. N. H. S. iv, p. 170 (1889) ; Seitz, Macrolep. ix, p. 1048 (1927).

Hasora lizetta anura Evans, Journ. Bomb. N. H. S. xxxi, p. 59 (1926).

Me Taw, W. Siam, Jan. 1920 (*Stockley*) ; Chiengmai, N. Siam (*Tennent*).

Distribution : Mussoorie to N. Burma ; Siam ; C. and W. China.

The typical form occurs in the Malay Peninsula, Sumatra, Java, Nias, and Sumbawa.

556. Hasora proxissima Elw. & Edw.

Hasora proxissima Elwes and Edwards, Trans. Zool. Soc. Lond. p. 302 (1897); Seitz, Macrolep. ix, p. 1051, fig. 166 d (1927).

Nong Yai Boo, Sriracha forest, S. E. Siam.

Distribution: Siam; Borneo; Philippines.

Apparently a very rare species.

557. Hasora badra badra (Moore).

Goniloba badra Moore, Cat. Lep. Mus. E. I. C. p. 295 (1857).

Hasora badra badra Evans, Journ. Bomb. N. H. S. xxxi, p. 59 (1926); Seitz, Macrolep. ix, p. 1048 (1927).

Chiengrai, N. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam.

Distribution: S. India; Sikkim to Burma and Siam; Andamans; China; Formosa; Malay Peninsula; Borneo; Sumatra; Java.

Evans (*l. c.*) recognizes *lanka* Ev. from Ceylon and *quadripunctata* Mab. from the Philippines, Celebes and Moluccas as good races, but sinks Fruhstorfer's races *godama*, *sankarya* and *madatta*.

558. Hasora vitta (Butl.) [= *chabrona* Plötz].

Ismene vitta Butler, Trans. Ent. Soc. Lond. p. 498 (1870); Seitz (*Hasora*), Macrolep. ix, p. 1050 (1927).

Ismene chabrona Plötz, Stett. Ent. Zeit. xlv, p. 56 (1884).

Me Song forest, Prae, N. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*); Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: Sikkim to Burma; S. W. China; Andamans; Siam; Malay Peninsula; Sumatra; Java; Borneo; Philippines.

559. Hasora alexis alexis (Fabr.) [= *chromus Cram.].**

Papilio alexis Fabricius, Ent. Syst. p. 533 (1775).

Hasora alexis alexis Seitz, Macrolep. ix, p. 1049 (1927).

Hup Bon and Nong Yai Boo, Sriracha forest, S. E. Siam.

Distribution: Ceylon; India; Burma; China; Siam; Celebes.

* The synonymy was set right by Aurivillius from an examination of the Fabrician type, which he discovered to be the species figured by Cramer as *chromus*.

The species ranges from India to China, Hainan, and Formosa; south to Australia.

560. Hasora taminatus almea Swinh.

Coeliades taminatus Hübner, Zutr. Ex. Schmett. figs. 193, 194 (1818).

Hasora almea Swinhoe, Ann. Mag. N. H. p. 90 (1909); Seitz, Macrolep. ix., p. 1049 (1927).

Hasora taminatus almea Evans, Journ. Bomb. N. H. S. xxxi, p. 60 (1926).

Recorded by Evans from Siam.

Distribution: S. Burma to Siam, the Malay Peninsula, and Borneo.

Evans recognizes as good races of *taminatus*:—*H. t. taminatus* Hüb. (Ceylon and S. India), and *H. t. malayana* Feld. (Andamans and Nicobars).

Seitz treats *almea* as a separate species confined to Borneo.

561. Hasora schönherri (Godt.) [= *chuza* Hew.].

Hesperia schönherri Godart, Enc. Meth. ix, p. 242 (1823).

Ismene chuza Hewitson, Ex. Butt. iv, Ismene, pl. 1. (1867).

Hasora schönherri chuza Seitz, Macrolep. ix, p. 1051, fig. 166 e (1927).

Hup Bon, Sriracha forest, S. E. Siam (*Godfrey*); Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: Assam to Burma; Siam; the Malay Peninsula; Sumatra; Borneo; Java.

The species ranges from Assam to Australia.

562. Ismene mahinthia Moore.

Ismene mahinthia Moore, Proc. Zool. Soc. Lond. p. 575 (1874); Seitz, Macrolep. ix, p. 1055 (1927).

1 ♂: Chet Ton (1,950 ft.), N. E. Siam, Dec. 1914 (*Godfrey*); 1 ♂, 1 ♀: Chiengmai district, N. Siam, March 1921 (*Tennent*).

Distribution: Assam; Burma; Siam.

563. Ismene oedipodea Swains.

Ismene oedipodea Swainson, Zool. Ill. i, p. 116 (1820).

Distribution: India to Tonkin, Hainan and the Philippines; south to the Malay Peninsula, Borneo and Java.

(a) **Ismene oedipodea athena** Fruhst.

Ismene oedipodea athena Fruhstorfer, Iris, p. 61 (1911); Seitz, Macrolep. ix, p. 1054 (1927).

Muak Lek and Pak Jong, E. Siam.

Distribution: E. Siam.

(b) **Ismene oedipodea oedipodea** Swains.

Ismene oedipodea oedipodea, Seitz, Macrolep. ix, p. 1053, fig. 167 b (1927).

Nakon Sritamarat, P. Siam (*Pendlebury; Kerr*).

Distribution: P. Siam; Malay Peninsula and Archipelago; Philippines.

564. Ismene jaina Moore.

Ismene jaina Moore, Proc. Zool. Soc. Lond. p. 782 (1865).

Distribution: India; Burma; Formosa; Siam; Celebes.

(a) **Ismene jaina jaina** Moore.

Ismene jaina jaina Seitz, Macrolep. ix, p. 1054, fig. 167 d (1927).

3 ♂♂: Me Song forest, Prae, N. Siam, April 1918.

Distribution: Mussoorie to Sikkim; N. Siam.

(b) **Ismene jaina margana** Fruhst.

Ismene jaina margana Fruhstorfer, Iris, p. 59 (1911); Seitz, Macrolep. ix, p. 1054 (1927).

Pak Jong and Hinlap, E. Siam.

Distribution: This subspecies, which was originally described from Siam, has since been found in Burma.

565. Ismene anadi (de Nicév.).

Choaspes anadi de Nicéville, Journ. As. Soc. Beng. p. 83 (1883); Seitz (*Ismene*), Macrolep. ix, p. 1054, fig. 171 i (1927).

1 ♀: Pak Jong, E. Siam, 3. 12. 14; 1 ♂, 1 ♀: 3. 1. 15.

Distribution: Mussoorie to the Karen Hills; Siam.

Apparently a very rare species.

566. Ismene harisa purpurea Ril. & Godf.

Ismene harisa Moore, Proc. Zool. Soc. Lond., p. 782 (1865); Seitz, Macrolep. ix, p. 1053, fig. 167 a (1927).

Ismene harisa purpurea Riley and Godfrey, Entomol. lviii, p. 143, pl. 3, fig. 7 (1925).

Me Song forest, Prae, N. Siam; Pak Jong and Muak Lek, E. Siam.

Distribution: Confined to Siam.

The species ranges from India to Tonkin, south to the Malay Peninsula and Archipelago.

567. *Ismene amara* Moore.

Ismene amara Moore, Proc. Zool. Soc. Lond. p. 783 (1865); Seitz, Macrolep. ix, p. 1053 (1927).

1 ♂: Me Wong (2,700 ft.), W. Siam, I. I. 24 (*Stockley*).

Distribution: Sikkim to the Shan States; N. Siam.

568. *Bibasis sena sena* (Moore).

Goniloba sena Moore, Proc. Zool. Soc. Lond. p. 778 (1865).

Bibasis sena sena Seitz, Macrolep. ix, p. 1052 (1927).

Me Song forest, Prae, N. Siam; Muak Lek and Pak Jong, E. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam.

Distribution: N. India to Burma, Siam, and the Malay Peninsula.

The species ranges from India to Borneo, Sumatra, Java, and the Philippines.

569. *Rhopalocampta benjamini* crawfurdii (Dist.).

Thymele benjamini Guérin, Delessert's Souv. Voy. Ind. ii, p. 79 (1843).

Ismene crawfurdii Distant, Ann. Mag. Nat. Hist. (5), x, p. 247 (1882).

Rhopalocampta benjamini *crawfurdii* Evans, Journ. Bomb. N. H. S. xxxi, p. 63 (1926).

Rhopalocampta subcaudata *crawfurdii* Seitz, Macrolep. ix, p. 1055 (1927).

Khao Ram, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: S. Burma; P. Siam; Malay Peninsula; Sumatra; Nias; Borneo.

The typical form is found in Ceylon, S. India, Simla to the Karen Hills, and China.

Seitz (*l. c.*) treats *crawfurdii* as a race of *subcaudata* Feld. and refers it to Malacca and Borneo.

570. *Badamia exclamacionis* (Fabr.).

Hesperia exclamacionis Fabricius, Syst. Ent. p. 530 (1775); Seitz (*Badamia*), Macrolep. ix, p. 1052 (1927).

Widely spread over Siam.

Distribution: Throughout the Indo-Australian region.

571. *Orthophaetus phaenus lalita* (Doh.).

Eudamus phaenus Hewitson, Descr. Hesp. p. 14 (1877).

Erionota lalita Doherty, Journ. As. Soc. Beng. p. 263 (1886).

Orthophaetus phaenus lalita Seitz, Macrolep. ix, p. 1032 (1927).

Me Song forest, Prae and Nam Phi (Sen Tor), N. E. Siam (Godfrey); Klong Klung and Me Lamoung, W. Siam (Stockley).

Distribution: Burma; Siam.

The species ranges from Assam to Perak, Borneo, and Sumatra.

Evans (Journ. Bomb. N. H. S. xxxi, p. 64, 1926) regards *phaenus* and *lalita* as separate species.

572. *Capila mackwoodi* (Evans) [= *barroni* Ril. & Godf.]

Orthophaetus mackwoodi Evans, Journ. Bomb. N. H. S. xxiii, p. 303 (1914); Evans (*Capila*), op. cit., xxxi, p. 65 (1926).

Orthophaetus barroni Riley and Godfrey, Journ. N. H. Soc. Siam, iv, p. 187, pl. 6, figs. 10, 11 (1921).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: N. Shan States to the Karen Hills; S. E. Siam.

Riley and Godfrey in describing *O. barroni* unfortunately overlooked Evans' description of *O. mackwoodi*.

The male types of both *mackwoodi* and *barroni* are in the British Museum and there is nothing by which they can be separated, except individual characteristics.

Mackwoodi was obtained at Aniskan, near Maymo, Burma, 24. 3. 14, and *barroni* at Hup Bon, 24. 4. 14.

This species is not recorded by Seitz.

573. *Charmion ficalnea ficalnea* (Hew.).

Hesperia ficalnea Hewitson, Deser. Hesp. p. 37 (1868).

Charmion ficalnea ficalnea Evans, Journ. Bomb. N. H. S. xxxi, p. 65 (1926); Seitz, Macrolep. ix, p. 1038, fig. 164 e (1927).

Recorded from Siam by Evans (*l. c.*) and by Swinhoe.*

Distribution: S. Burma; Siam; Malay Peninsula; Sumatra; Borneo.

Subspecies are known from Nias and the Celebes.

Seitz (*l. c.*) gives *queda* Plötz as the Siamese race, but Evans sinks this under *ficulnea*.

574. *Celaenorrhinus leucocera leucocera* (Kollar).

Hesperia leucocera Kollar, in Hugel's Kaschmir, iv, p. 454 (1844).

Celaenorrhinus leucocera leucocera Seitz, Macrolep. ix, p. 1036, fig. 163 g (1927).

Me Song forest, Prae and Chiengrai, N. Siam (*Godfrey*); Me Lamoung, W. Siam (*Stockley*).

Distribution: Sikkim to Burma; Siam; Annam; Malay Peninsula.

A subspecies is known from W. China and another from Sumatra, Java, Bali, and Borneo.

575. *Celaenorrhinus maculicornis* Elw. & Edw.

Celaenorrhinus maculicornis Elwes and Edwards, Trans. Zool. Soc. Lond. p. 116 (1897); Seitz, Macrolep. ix, p. 1036, fig. 164 (1927).

Muak Lek, E. Siam and Sriracha forest, S. E. Siam; Chiengrai, N. Siam.

Distribution: Assam to Manipur; Siam.

576. *Celaenorrhinus asmara aditta* Fruhst.

Hesperia asmara Moore, Cat. Lep. Mus. E. I. C. p. 253 (1857).

Celaenorrhinus asmara aditta Fruhstorfer, Iris, p. 63 (1910); Seitz, Macrolep. ix, p. 1037 (1927).

Me Song forest, Prae, N. Siam; Pak Jong and Muak Lek, E. Siam.

Distribution: Fruhstorfer (*l. c.*) restricts this subspecies to Siam.

The species ranges from Burma to Sumatra, Java and the Philippines.

* Swinhoe, Lep. Ind. x, p. 28.

577. *Celaenorrhinus nigricans nigricans* (de Nicév.).

Plesioneura nigricans de Nicéville, Journ. As. Soc. Beng. p. 123 (1885).

Celaenorrhinus nigricans nigricans Evans, Journ. Bomb. N. H. S. xxxi, p. 68 (1926); Seitz, Macrolep. ix, p. 1037 (1927).

1 ♂: Me Lamoung (2,000 ft.), W. Siam, 14. 4. 20 (Stockley).

Distribution: Sikkim to S. Burma; Siam; Malay Peninsula.

A subspecies is known from Borneo.

578. *Celaenorrhinus aurivittata* (Moore).

Plesioneura aurivittata Moore, Proc. Zool. Soc. Lond. p. 843 (1878).

Distribution: Assam to the Malay Peninsula, Sumatra, and Borneo.

(a) *Celaenorrhinus aurivittata aurivittata* (Moore).

Celaenorrhinus aurivittata aurivittata Seitz, Macrolep. ix, p. 1038, fig. 164 c (1927).

Me Song forest, Prae, N. Siam and Sriracha forest, S. E. Siam (Godfrey); Me Lamoung, W. Siam (Stockley).

Distribution: Assam to Mergui; Siam.

(b) *Celaenorrhinus aurivittata cameroni* (Dist.).

Plesioneura cameroni Distant, Ann. Mag. N. H. p. 248 (1882).

Celaenorrhinus aurivittata cameroni Seitz, op. cit., p. 1038.

Tai Sai River (Pendlebury); Wang Sai (Kerr) - Nakon Sritamarat, P. Siam.

Distribution: S. Mergui; P. Siam; Malay Peninsula; Sumatra.

579. *Achalarus bifasciatus liliana* (Atks.).

Eudamus bifasciatus Bremer and Grey, Schmett. N. China, p. 10 (1853).

Plesioneura liliana Atkinson, Proc. Zool. Soc. Lond. p. 216 (1871); Seitz (*Lobocla*), Macrolep. ix, p. 1032 (1927).

Achalarus bifasciatus liliana Evans, Journ. Bomb. N. H. S. xxxi, p. 69 (1926).

2 ♂♂: Doi Chang (4,500 ft.), Chiengrai, N. Siam, 13. 4. 21.

Distribution: Assam to the Karen Hills; Yunnan; N. Siam.

The typical form is found in China and a subspecies, *A. b. easyapa* Moore, occurs from Kashmir to Kumaon.

580. *Tagiades atticus* (Fabr.).

Hesperia atticus Fabricius, Ent. Syst. iii (1), p. 339 (1793).

Tagiades atticus Evans, Journ. Bomb. N. H. S. xxxi, p. 71 (1926).

Evans (*l. c.*) treats this species as follows:—

T. a. atticus Fabr. (= *ravi* Moore; *lugens* Mab.; *utanus* Plotz; *yotissa* Fruhst.).

Distribution: Tavoy to the Malay Peninsula; Pulo Condore; Sumatra.

T. a. khasiana Moore. (= *epicharmus* Fruhst.).

Distribution: Central Provinces to Bengal; Dun to Tavoy; Siam.

T. a. helferi Feld. (= *ravina* Fruhst.).

Distribution: Andamans and Nicobars.

T. a. rajaghra Fruhst. (= *balana* Fruhst.).

Distribution: Borneo.

Seitz in the "Macrolepidoptera of the World" makes no mention of *atticus*. He regards *ravi* as a "doubtful species", and recognizes as subspecies of it *yotissa* Fruhst, *rajaghra* Fruhst, *khasiana* Moore, *epicharmus* Fruhst, *balana* Fruhst. and *ravina* Fruhst.

Two forms of *atticus* occur in Siam.

(a) **Tagiades atticus atticus f. atticus** (Fabr.).

Me Song forest, Prae, N. Siam; Pak Jong and Muak Lek, E. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam.

This is the wet season form.

(b) **Tagiades atticus atticus f. khasiana** Moore.

Tagiades khasiana Moore, Journ. As. Soc. Beng. liii, p. 51 (1884).

Found in the same localities as the above.

This is the dry season form.

581. **Tagiades gana gana** Moore.

Tagiades gana Moore, Proc. Zool. Soc. Lond. p. 180 (1865); Seitz, Macrolep. ix, p. 1040, fig. 164 h (1927).

Pak Jong, E. Siam.

Distribution: Sikkim to the Karen Hills; Siam; Cambodia.

The species ranges from Sikkim to the Malay Archipelago and the Philippines.

582. **Tagiades menanto** Plötz.

Tagiades menanto Plötz, Berl. Ent. Zeit. xxix, p. 231 (1885); Seitz

Macrolep. ix, p. 1042, fig. 165 d (1927).

Tagiades gana menanto Evans, Journ. Bomb. N. H. S. xxxi, p. 71 (1926).

Khao Ram and Khao Luang, Nakon Sritamarat, P. Siam (Pendlebury).

Distribution : Dawnas to S. Burma ; Siam ; Malay Peninsula ; Nias ; Sumatra ; Borneo ; Java.

Evans (*l. c.*) treats *menanto* (= *elongata* and *niasana* Mab. ; *avala*, *jainas* and *parra* Fruhst.) as a subspecies of *gana* Moore.

Riley* regards it as quite a distinct species, identical with *nana* Elw. & Edw.

583. *Tagiades japerus meetana* Moore (= *patimoka* Fruhst.).

Papilio japerus Cramer, Pap. Exot. iv, p. 365 (1782).

Tagiades meetana Moore, Proc. Zool. Soc. Lond. p. 593 (1877).

Tagiades obscurus meetana Evans, Journ. Bomb. N. H. S. xxxi, p. 72 (1926).

Tagiades obscurus alica Seitz, Macrolep. ix, p. 1039 (1927).

Tagiades japerus patimoka Seitz, Macrolep. ix, p. 1039 (1927).

Muak Lek, E. Siam ; Sriracha forest, S. E. Siam.

Distribution : Karen Hills to Mergui ; Tonkin ; Siam.

Evans (*l. c.*) treats *meetana* as a subspecies of *obscurus* Mab. Riley*, however, does not consider *obscurus* a good species ; he thinks it is probably the S. Indian race of *japerus*.

Seitz (*l. c.*) regards *meetana* as a synonym of *alica* Moore, which he gives as a subspecies of *obscurus*.

Fruhstorfer's *patimoka*, which Seitz (*l. c.*) regards as "perhaps a distinct species from Siam", must be considered an absolute synonym of *meetana* Moore.

584. *Tagiades toba* de Nicév.

Tagiades toba de Nicéville, Journ. Bomb. N. H. S. x, p. 19 (1896) ; Evans, Journ. Bomb. N. H. S. xxxi, p. 72 (1926) ; Seitz, Macrolep. ix, p. 1042, fig. 165 d (1927).

Evans (*l. c.*) records this species from P. Siam.

Distribution : According to Evans, the species occurs in S. Burma, P. Siam, the Malay Peninsula, Sumatra, Nias, and Borneo.

* Riley, *in litt.*, 11. 8. 27.

585. *Tagiades waterstradti* Elw.

Tagiades waterstradti Elwes, Trans. Zool. Soc. p. 143 (1897); Evans, Journ. Bomb. N. H. S. xxxi, p. 72 (1926).

Tagiades menaka waterstradti, Seitz, Macrolep. ix, p. 1041 (1927).

Evans (*l. c.*) records this species from P. Siam and assigns it to P. Siam, the Malay Peninsula, N. Borneo and Java.

Both Seitz (*l. c.*) and Riley* regard *waterstradti* as a purely Bornean form.

586. *Tagiades menaka menaka* (Moore).

Pterygospidea menaka Moore, Proc. Zool. Soc. p. 778 (1865); Seitz (*Tagiades*), Macrolep. ix, p. 1041 (1927).

2 ♂♂: Huey Tak So, W. Siam, 23. 2. 24 (Stockley).

Distribution: Kashmir to the Karen Hills; W. China; Tonkin; Siam.

According to Seitz, the species ranges from N. India to New Guinea.

587. *Tagiades litigiosa litigiosa* Mösch. (= *atticus* Moore nec Fabr.).

Tagiades litigiosa Möschler, Verh. Zeol-bot. Ges. Wien. xlvi, p. 230 (1878).

Tagiades litigiosa litigiosa Evans, Journ. Bomb. N. H. S. xxxi, p. 73 (1926).

Tagiades menaka litigiosa Seitz, Macrolep. ix, p. 1041 (1927).

Me Song forest, Prae, N. Siam; Muak Lek, E. Siam; Sriracha forest, S. E. Siam; Bangnara, P. Siam.

Distribution: Sikkim to Burma; Andamans; W. China; Hainan; Siam.

A subspecies, *T. l. rajuna* Fruhst., is known from Ceylon and S. India.

Seitz (*l. c.*) treats *litigiosa* as a subspecies of *menaka* Moore, but both Riley and Evans regard it as a separate species.

588. *Odina decoratus* (Hew.).

Eudamus decoratus Hewitson, Deser. Hesp. p. 17 (1867); Seitz (*Odina*), Macrolep. ix, p. 1044, fig. 165 f (1927).

*Riley, *in litt.*, 11. 8. 27.

Muak Lek, E. Siam (*Fruhstorfer**); Klung, Chantabun, and Kao Sabap - S. E. Siam (*Kerr*).

Distribution: Sikkim to Burma; Tonkin; Siam.

589. *Mooreana trichoneura pralaya* Moore.

Pterygospidea trichoneura Felder, Wien. Ent. Monats. iv, p. 402 (1860).

Mooreana trichoneura pralaya Evans, Journ. Bomb. N. H. S. xxxi, p. 73 (1926).

Tagiades trichoneura pralaya Seitz, Macrolep. ix, p. 1043 (1927).

Me Song forest, Prae, N. Siam; Sriracha forest, S. E. Siam.

Distribution: Sikkim to N. Burma; Siam.

According to Evans, the typical form is found from the Karen Hills to S. Burma, in the Malay Peninsula, Sumatra, Nias, and Java.

590. *Daimio phisara* (Moore).

Satarupa phisara Moore, Journ. As. Soc. Beng. p. 50 (1884).

Satarupa bhagava phisara Seitz, Macrolep. ix, p. 1034, fig. 163 f (1927).

Me Song forest, Prae, N. Siam (*Godfrey*); Me Wong, W. Siam (*Pendlebury*).

Distribution: Sikkim to Burma; Siam; Perak.

591. *Daimio bhagava bhagava* (Moore) [= *milliana* Swinh.]

Satarupa bhagava Moore, Proc. Zool. Soc. Lond. p. 781 (1865); Seitz, Macrolep. ix, p. 1034 (1927).

Me Song forest, Prae, N. Siam; Muak Lek, E. Siam; Sriracha forest, S. E. Siam.

Distribution: Sikkim to Burma; Siam.

A subspecies *D. b. andamanica* Wood-Mason, is known from the Andaman Islands.

592. *Daimio limax dirae* de Nicéville.

Tagiades limax Plötz, Jahrb. Nassau. Ver. xxxvii, p. 54 (1884).

Daimio dirae de Nicéville, Journ. Bomb. N. H. S. ix, p. 369 (1895); Seitz (*Satarupa*), Macrolep. ix, p. 1033, fig. 163 e (1927).

Daimio limax dirae Evans, Journ. Bomb. N. H. S. xxxi, p. 75 (1926).

Khao Luang, Nakon Sritamarat, P. Siam (*Pendlebury*; *Kerr*).

* Fruhstorfer, Iris, xxiv, p. 70 (1910).

Distribution: S. Burma to Tavoy; P. Siam; Malay Peninsula; Borneo; Sumatra; Nias.

The typical form occurs in Java, and a subspecies is known from the Philippines.

593. *Daimio tabrica pinwilli* (Butl.).

Pterygospidia tabrica Hewitson, Exot. Butt. v, Pterygospidea, pl. 1 fig. 8 (1873).

Plesioneura pinwilli Butler, Trans. Linn. Soc. Zool. Lond. p. 556 (1877); Seitz (*Tagiades*), Macrolep. ix, p. 1043, pl. 165 e (1927).

Daimio tabrica pinwilli Evans, Journ. Bomb. N. H. S. xxxi, p. 75 (1926).

2 ♂♂: Me Song forest, Prae, N. Siam, April 1918.

Distribution: Assam to Burma; Siam; Malay Peninsula; Sumatra; Borneo.

Evans (*l. c.*) refers the typical form to Darjiling, and accepts *D. t. bowringi*, Joicey and Talbot (Hainan), as a "very distinct race".

594. *Coladenia indrani uposatha* Fruhst.

Hesperia indrani Moore, Cat. Lep. Mus. E. I. C. i, p. 253 (1857).

Coladenia indrani uposatha Evans, Journ. Bomb. N. H. S. xxxi, p. 75 (1926); Seitz, Macrolep. ix, p. 1034 (1927).

Me Song forest, Prae, N. Siam; Sriracha forest, S. E. Siam.

Distribution: N. Burma to the Karen Hills; Siam.

According to Evans (*l. c.*), the typical form occurs from Mussoorie to Sikkim, and good subspecies are known from S. India and Ceylon.

595. *Coladenia dan dhyana* Fruhst.

Papilio dan Fabricius, Mant. Ins. ii, p. 88 (1787).

Coladenia dan dhyana Fruhstorfer, Iris, p. 66 (1910); Seitz, Macrolep. ix, p. 1035 (1927).

Widely spread over Siam.

Distribution: S. Burma to the Karen Hills; Siam; Malay Peninsula.

The species ranges from India to China and the Philippines, south to Borneo, Sumatra, and Java.

596. *Coladenia laxmi* (de Nicév.).

Plesioneura laxmi de Nicéville, Journ. As. Soc. Beng. p. 290 (1888); Seitz (*Coladenia*), Macrolep. ix, p. 1035 (1927).

1 ♂, ♀ : Me Song forest, Prae, N. Siam, 14. 3. 16.

Distribution : Sikkim to Burma; Siam; Perak; Sumatra; Borneo; and (?) the Philippines.

597. Sarangesa dasahara dasahara (Moore).

Nisoniades dasuhara Moore, Proc. Zool. Soc. Lond. p. 787 (1865).

Sarangesa dasahara dasahara Evans, Journ. Bomb. N. H. S. xxxi, p. 76 (1926); Seitz, Macrolep. ix, p. 1044 (1927).

Chiengrai, N. Siam; Muak Lek, E. Siam; Sriracha forest, S. E. Siam (*Godfrey*); Me Lamoung, W. Siam (*Stockley*).

Distribution : Central Provinces, India; Burma; Siam, Tonkin.

Evans (*l. c.*) recognizes as good subspecies :—*S. d. albicilia* Moore (Ceylon), and *S. d. davidsoni* Swinh. (S. India to Bombay).

598. Darpa pteria (Hew.) [= *dealbata* Dist.].

Pterygospidea pteria Hewitson, Descr. Hesp. p. 51 (1868); Evans (*Darpa*), Journ. Bomb. N. H. S. xxxi, p. 77 (1926); Seitz (*Tagiades*), Macrolep. ix, p. 1042, pl. 165 d (1927).

1 ♂ : Doi Chang (4,500 ft.), Chiengrai, N. Siam, 13. 4. 21, Also obtained by Stockley on the Me Wong, W. Siam.

Distribution : S. Burma; Siam; Malay Peninsula; Borneo; Philippines.

599. Tapena thwaitesi minuscula Elw. & Edw.

Tapena thwaitesi Moore, Lep. Ceylon i, p. 181 (1881).

Tapena minuscula Elwes and Edwards, Proc. Zool. Soc. Lond., p. 659 (1892).

Tapena thwaitesi minuscula Evans, Journ. Bomb. N. H. S. xxxi, p. 77 (1926); Seitz, Macrolep. ix, p. 1044, fig. 165 g (1927).

Chiengrai, N. Siam and Sriracha forest, S. E. Siam (*Godfrey*) Me Wong, W. Siam (*Stockley*).

Distribution : Assam to S. Burma; Siam; Malay Peninsula; Borneo and Sumatra.

According to Evans the typical form occurs in Ceylon.

600. Ctenoptilum vasava vasava (Moore).

Achyloides vasava Moore, Proc. Zool. Soc. p. 786 (1865); Seitz (*Ctenoptilum*), Macrolep. ix, p. 1044, fig. 165 g (1927),

4 ♂♂ : Huey Tak So, W. Siam, 23. 2. 24 (*Stockley*).

Distribution : Mussoorie to Burma; Siam.

Seitz (*l. c.*) recognizes as a good subspecies *Ct. v. chinensis* Leech, from China.

601. Ctenoptilum multiguttata de Nicév.

Ctenoptilum multiguttata de Nicéville, Journ. Bomb. N. H. S. p. 221 (1890); Seitz, Macrolep. ix, p. 1044 (1927).

Me Song forest, Prae, N. Siam and Pak Jong, E. Siam (*Godfrey*), Me Lamoung, W. Siam (*Stockley*).

Distribution: Manipur to Burma; Siam.

602. Odontoptilum angulata (Feld.).

Pterygospidea angulata Felder, Verh. Zool-bot Ges. Wien. xi, p. 482 (1862); Evans (*Odontoptilum*), Journ. Bomb. N. H. S. xxxi, p. 77 (1926); Seitz (*Odontoptilum*), Macrolep. ix, p. 1045, fig. 165 g (1927).

Me Song forest, Prae, N. Siam and Sriracha forest, S. E. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*); Me Wong, W. Siam (*Stockley*).

Distribution: S. India; Kulu to Burma; S. China; Siam; Malay Peninsula; Sumatra; Java; Borneo; Lombok; Sumbawa; Philippines.

Under *angulata*, Evans (*l. c.*) sinks the following forms:—
sura Moore; *sumatrana mahabina*, *subangulata* and *hyperides* Fruhst.; *helisa* Semp.; and *hypocedes* Doh.

603. Caprona ransonnettii siamica Swinh.

Pterygospidia ransonnettii Felder, Vehr. Zool-bot Ges. Wien., p. 284 (1868).

Abaratha siamica Swinhoe, Ann. Mag. Nat. Hist. p. 432 (1907).

Abaratha alida siamica Seitz, Macrolep. ix, p. 1046, fig. 165 h (1927).

Caprona ransonnettii alida Evans, Journ. Bomb. N. H. S. xxxi, p. 78 (1926).

Me Song forest, Prae, N. Siam (*Godfrey*); N. Siam (*Swinhoe*).

Distribution: Siam.

The species ranges from N. India to Ceylon.

I retain the name *siamica* for Siamese specimens, but probably it is merely a synonym of *alida* de Nicév.

604. Caprona agama (Moore).

Pyrgus agama Moore, Cat. Lep. Mus. E. I. C. i, p. 249 (1857); Evans (*Caprona*), Journ. Bomb. N. H. S. xxxi, p. 78 (1926).

Abaratha syriethus agama Seitz, Macrolep. ix, p. 1046 (1927).

Caprona elwesi Watson, Journ. Bomb. N. H. S. x, p. 674 (1896).

Chiengrai, N. Siam; Sriracha forest, S. E. Siam.

Distribution:—S. India to Mussoorie and Burma; Siam; Tonkin; Java; Bali.

C. elwesi, recorded from Siam by Mabille (Bull. Soc. Ent. France, p. 101, 1917), is treated by Evans as a variety of *agama* which occurs not only in Siam, but from Assam to the Shan States.

Under *agama*, Evans sinks *syriethus* Feld., *pelias* Fruhst., and *parvopunctata* Mab.

605. *Hesperia galba* (Fabr.).

Papilio galba Fabricius, Ent. Syst. iii (1), p. 352 (1793); Evans (*Hesperia*), Journ. Bomb. N. H. S. xxxi, p. 79 (1926); Seitz (*Hesperia*), Macrolep. ix, p. 1046, fig. 165 i (1927).

Muak Lek and Pak Jong, E. Siam (*Godfrey*); Kao Tok, Kanburi, W. Siam (*Kerr*).

Distribution: Evans (*l. c.*) gives the distribution of this species as:—"Ceylon, S. India to Karachi, India generally to Shan States".

606. *Astictopterus jama olivascens* Moore (= *kada* Swinh.).

Astictopterus jama Felder, Wien. Ent. Monats. iv, p. 401 (1860).

Astictopterus jama olivascens Evans, Journ. Bomb. N. H. S. xxxi, p. 429 (1926).

Astictopterus fuligo olivascens Seitz, Macrolep. ix, p. 1097, fig. 175 b (1927).

Chiengmai, N. Siam (*Tennent*); Sen Tor (Nam Pat), N. E. Siam, and Bangkok, C. Siam (*Godfrey*).

Distribution: S. India; Kumaon to Tavoy; Andamans; Tonkin; Siam.

The typical form occurs in Mergui and the Malay Peninsula, and a subspecies is found in W. China and Hainan.

Seitz (*l. c.*) treats *olivascens* as a subspecies of *fuligo* Mab., but Evans sinks the latter name under *jama* Feld.

607. *Ochus subvittatus* (Moore).

Cyclopides subvittatus Moore, Proc. Zool. Soc. Lond. p. 692 (1878); Seitz (*Ochus*), Macrolep. ix, p. 1073, fig. 170 f (1927); Evans (*Ochus*), Journ. Bomb. N. H. S. xxxi, p. 430 (1926).

Me Song forest, Prae, and Chinegrai, N. Siam (*Godfrey*); Tai

Sai river and Khao Ram, P. Siam (*Pendlebury*) ; Ban Ton, Satul, P. Siam (*Kerr*).

Distribution : Kumaon to the Dawnas ; Tonkin ; Siam.

608. *Ampittia dioscorides* (Fabr.) [= *maro* Fabr.]

Hesperia dioscorides Fabricius, Syst. Ent. iii, p. 329 (1793) ; Evans (*Ampittia*), Journ. Bomb. N. H. S. xxxi, p. 430 (1926).

Ampittia maro Seitz, Macrolep. ix, p. 1074 (1927).

Widely spread over Siam.

Distribution : According to Evans (*l. c.*) the range of this species is :—“Ceylon, India, Burma, Siam, Shanghai, Tonkin, Malay Peninsula, Sumatra, Bali.”

609. *Aeromachus pygmaeus pygmaeus* (Fab.).

Aeromachus pygmaeus pygmaeus Evans, Journ. Bomb. N. H. S. xxxi, p. 432 (1926).

1 ♂ : Tung Wa, Satul, P. Siam, 4. 1. 28 (*Kerr*).

Distribution : Evans (*l. c.*) refers typical *pygmaeus* to “S. India, Assam to Burma, Malay Peninsula, Sumatra,” and recognizes as a good race *A. p. musca* Mab. from Luzon.

Dr. Kerr’s specimen was identified by Riley, and is now in the British Museum.

610. *Arnetta atkinsoni* (Moore).

Isoteinon atkinsoni Moore Proc. Zool. Soc. Lond. p. 844 (1878) ; Evans (*Arnetta*), Journ. Bomb. N. H. S. xxxi, p. 432 (1926) ; Seitz (*Arnetta*) ; Macrolep. ix, p. 1064 (1927).

1 ♂ : Me Song forest, Prae, N. Siam, April 1918.

Distribution : Sikkim to Tavoy ; Siam.

Evans (*l. c.*) regards *subtestaceus* and *khasianus* Moore as synonyms of *atkinsoni*.

611. *Iambrix tytleri woolletti* (Ril.).

Iambrix tytleri Evans, Journ. Bomb. N. H. S. xxiii, p. 302 (1914).

Mimambrix woolletti Riley, Entomol. Ivi, p. 37, pl. 1, fig. 5 n (1923) ; Riley and Godfrey, op. cit., lviii, p. 142, pl. 3, fig. 6 ♀ (1925).

1 ♀ : Phar Mieng, near Chiengrai, N. Siam, 16. 4. 21 (*in British Museum*).

Distribution : Borneo ; Siam.

According to Evans, the typical form occurs in the Naga Hills.

612. *Iambrix salsala salsala* (Moore).

Nisionades salsala Moore, Proc. Zool. Soc. Lond. p. 786 (1865).

Iambrix salsala salsala Evans, Journ. Bomb. N. H. S. xxxi, p. 433 (1926); Seitz, Macrolep. ix, p. 1063, fig. 168 (1927).

Widely spread over Siam.

Distribution : Sikkim to Burma ; China ; Tonkin ; Siam ; Malay Peninsula and Archipelago.

Evans (*l. c.*) recognizes as a good race *I. s. luteipalpus* Plötz from Ceylon and S. India, but sinks Fruhstorfer's *vasuba* from Tonkin.

613. *Iambrix stellifer* (Butl.).

Astictopterus stellifer Butler, Trans. Linn. Soc. ser. 2, Zool. 1, p. 555 (1877); Seitz (*Iambrix*), Macrolep. ix, p. 1063, fig. 168 d (1927).

Me Wong, W. Siam (*Stockley*) ; Bangnara, P. Siam (*Godfrey*) ; Nakon Sritamarat, P. Siam (*Pendlebury* ; *Kerr*).

Distribution : Burma to the Malay Peninsula, Sumatra, Nias, and Java.

614. *Suastus gremius gremius* (Fabr.).

Hesperia gremius Fabricius, Ent. Syst. Suppl. p. 433 (1798).

Suastus gremius gremius Evans, Journ. Bomb. N. H. S. xxxi, p. 434 (1926); Seitz, Macrolep. ix, p. 1062 (1927).

Muak Lek and Pak Jong, E. Siam ; Bangkok, C. Siam.

Distribution : India ; Burma ; S. China ; Formosa ; Siam.

Evans (*l. c.*) recognizes *subgrisea* Moore from Ceylon as a good race ; Seitz (*l. c.*) sinks it.

615. *Suastus rama rama* (Mab.).

Pamphila rama Mabille, Bull. Soc. Ent. France, p. 198 (1876).

Suastus rama rama Evans, Journ. Bomb. N. H. S. xxxi, p. 435 (1926).

Suastus sala Seitz, Macrolep. ix, p. 1062 (1927).

Pak Jong and Muak Lek, E. Siam ; Sriracha, S. E. Siam.

Distribution : Sikkim to Burma ; Siam.

Evans (*l. c.*) recognizes as good races :—*S. r. aditus* Moore from the Andaman and Cocos Islands, and *S. r. bipunctus* from the Nilgiris.

Seitz makes no mention of *rama*. He regards *aditus* as a synonym of *sala* Hew., which he treats as a separate species ranging from Sikkim to Java.

Evans sinks Seitz' *sala* under *rama*, remarking that true *sala* is a *Plastingia*.

616. *Scobura cephalata* (Hew.).

Hesperia cephalata Hewitson, Ent. Mo. Mag. p. 152 (1876); Seitz (*Scobura*), Macrolep. ix, p. 1069 (1927).

Me Song forest, Prae and Chiengrai, N. Siam.

Distribution: Sikkim to Burma; Siam; Malay Peninsula; Sumatra; Borneo.

617. *Suada swerga swerga* (de Nicév.).

Hesperia swerga de Nicéville, Journ. As. Soc. Beng. p. 89 (1883); Seitz (*Suada*), Macrolep. ix, p. 1062, fig. 168 c (1927).

Suada swerga swerga Evans, Journ. Bomb. N. H. S. xxxi, p. 435 (1926).

Me Song forest, Prae, and Chiengrai, N. Siam; Sriracha, S. E. Siam.

Distribution: Sikkim to Burma; Siam; Malay Peninsula; Sumatra.

Evans (*l. c.*) recognizes a subspecies from Java; and another from Borneo, Pulo Laut and the Philippines.

618. *Koruthaialos rubecula* (Plötz).

Lychnuchus rubecula Plötz, Berl. Ent. Zeit. xxvi, p. 264 (1882); Evans (*Koruthaialos*), Journ. Bomb. N. H. S. xxxi, p. 436 (1926); Seitz (*Koruthaialos*), Macrolep. ix, p. 1072, fig. 170 d (1927).

1 ♂, 2 ♀ ♀: Me Song forest, Prae, N. Siam April 1916; 5 ♂ ♂, 4 ♀ ♀, April 1918.

Distribution: Evans (*l. c.*) gives the distribution of this species as "Assam to Burma, Malay Peninsula, and Islands, Philippines, Luzon, Natunas, Celebes", but remarks:—"I have verified true *rubecula* from Assam to the Malay Peninsula and Sumatra, but the other localities require check".

619. *Koruthaialos butleri butleri* (W-M. & de Nicév.).

Astictopterus butleri Wood-Mason and de Nicéville, Journ. As. Soc. Beng. p. 98 (1883); Seitz (*Koruthaialos*) Macrolep. ix, p. 1072 (1927).

Koruthaialos butleri butleri Evans, Journ. Bomb. N. H. S. xxxi, p. 436 (1926).

1 ♂: Pak Jong, E. Siam, Jan. 1915.

Distribution: Sikkim to Burma; Siam.

Evans (*l. c.*) recognizes subspecies from S. Burma and S. Annam; from Sumatra and Nias; and from Java.

620. Koruthaialos xanites xanites (Butl.).

Astictopterus xanites Butler, Trans. Ent. Soc. Lond. p. 510 (1870); Evans (*Koruthaialos*), Journ. Bomb. N. H. S. xxxi, p. 436 (1926); Seitz (*Koruthaialos*), Macrolep. ix, p. 1072, fig. 170 d (1927).

Me Song forest, Prae, N. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*); Kao Kalakiri, Pattani (*Kerr*).

Distribution: S. Burma; Siam; Malay Peninsula; Borneo; Philippines.

Evans (*l. c.*) recognizes as good subspecies:—*K. x. gopaka* Fruhst. (Assam to Burma, Tonkin); *K. x. kerala* de Nicév. (Sumatra); and *K. x. javanites* Staud. (Java, Bali, Batu).

621. Sancus pulligo subfasciatus Moore.

Tagiades pulligo Mabille, Ann. Ent. Soc. France, p. 272 (1876).

Sancus pulligo subfasciatus Evans, Journ. Bomb. N. H. S. xxxi, p. 437 (1926); Seitz, Macrolep. ix, p. 1037 (1927).

Me Song forest, Prae, N. Siam and Sriracha forest, S. E. Siam (*Godfrey*); Kambengpet, W. Siam (*Stockley*).

Distribution: S. India; Burma; Siam; Annam; Malay Peninsula.

The species ranges from India to Sumatra, Borneo, Java, Bali, the Celebes, and the Philippines.

622. Kerana nigrita (Godt.).

Hesperia nigrita Godart, Enc. Meth. ix, p. 788 (1823); Evans (*Kerana*), Journ. Bomb. N. H. S. xxxi, p. 437 (1926).

Kerana diocles Seitz, Macrolep. ix, p. 1097 (1927).

Pak Jong, E. Siam and Sriracha, S. E. Siam (*Godfrey*); Me Wong, W. Siam (*Stockley*).

Distribution: Bengal; Kumaon to Burma; Siam; Malay Peninsula and Archipelago; Philippines.

Under *nigrita* Evans (*l. c.*) sinks *diocles* Moore; *maura* Snell.; *fumatus* Mab.; *evaira*, *sumata* and *prabha* Fruhst.

Evaira was described from Siam.

623. Kerana armata armata (Druce).

Astictoperus armatus Druce, Proc. Zool. Soc. p. 359 (1873).

Kerana armata armata Evans, Journ. Bomb. N. H. S. xxxi, p. 437

(1926); Seitz, Macrolep. ix, p. 1097, fig. 175 c (1927).

2 ♂♂: Chawng, Trang, P. Siam, 14. 4. 28 (Kerr).

Distribution: S. Mergui; P. Siam; Malay Peninsula; Sumatra; Borneo.

Evans has described a race, *K. a. niasana*, from Nias.

624. Kerana fulgur de Nicév.

Kerana fulgur de Nicéville, Journ. As. Soc. Beng. p. 86 (1883); Seitz, Macrolep. ix, p. 1098, fig. 175 c (1927).

Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: P. Siam; Malay Peninsula; Sumatra; Borneo.

625. Udaspes folus (Cram.).

Papilio folus Cramer, Pap. Exot. i, pl. 74, fig. 7 (1779); Seitz (*Udaspes*), Macrolep. ix, p. 1096, fig. 175 a (1927).

Widely spread over Siam, but not common.

Distribution: Throughout the Indo-Australian region.

626. Notocrypta volux volux (Mabille).

Plesioneura volux Mabille, Ann. Soc. Ent. Belg. p. 56 (1883).

Notocrypta volux volux Evans, Journ. Bomb. N. H. S. xxxi, p. 438 (1926).

Notocrypta alydos volux Seitz, Macrolep. ix, p. 1095 (1927).

Me Song forest, Prae, N. Siam (*Godfrey*); Me Wong and Upper Me Klong, W. Siam (*Stockley*); Nakon Sritamarat, P. Siam (*Pendlebury*; *Kerr*).

Distribution: Karen Hills to Burma; Siam; Cambodia; Malay Peninsula; Borneo; Philippines.

Evans has described a subspecies *N. v. niasana* from Nias.

Seitz (*l. c.*) treats *volux* as a subspecies of *alydos* Moore.

Both Riley and Evans regard *alydos* as a subspecies of *feisthamelii* Boisd.

627. Notocrypta paralysos devadatta Fruhst.

Plesioneura paralysos Wood-Mason and de Nicéville, Journ. As. Soc. Beng. p. 257 (1881).

Notocrypta devadatta Fruhstorfer, Iris, p. 24 (1911).

Notocrypta paralysos devadatta Evans, Journ. Bomb. N. H. S. xxxi, p. 438 (1926).

Notocrypta alydos devadatta Seitz, Macrolep. ix, p. 1094 (1927).

Me Song forest Prae, N. Siam; Muak Lek, E. Siam; Hup

Bon, S. E. Siam.

Distribution : Dawnas to S. Burma; Siam; French Laos; Malay Peninsula; Sumatra; Nias.

Evans (*l. c.*) recognizes as good races :—*N. p. paralyos* W-M. & de Nicév. (Andamans); *N. p. alysia* Ev. (Ceylon, S. India, Mussoorie to the Karen Hills); and *N. p. sidha* Fruhst. (Java).

628. Notocrypta feisthamelii alydos (Moore).

Thymele feisthamelii Boisduval, Voy. Astrolabe, Lep. p. 159 (1832).

Plesioneura alydos Moore, Proc. Zool. Soc. Lond. p. 789 (1865).

Notocrypta feisthamelii alydos Evans, Journ. Bomb. N. H. S. xxxi, p. 439 (1926).

Notocrypta alydos alydos Seitz, Macrolep. ix, p. 1094, fig. 174 g (1927).

Me Song forest, Prae, N. Siam and Nam Phi, N. E. Siam (*Godfrey*); Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution : Murree to Burma; W. China; Siam; French Laos; Malay Peninsula; Sumatra; Borneo.

Evans (*l. c.*) recognizes as good races :—*N. f. avattana* Fruhst. (Java, Lombok, and Sumbawa); *N. f. alinkara* Fruhst. (Philippines); *N. f. celebensis* Staud. (Celebes); and *N. f. feisthamelii* Boisd. (Moluccas to Queensland).

629. Notocrypta curvifascia (Feld.).

Plesioneura curvifascia Felder, Wien. Ent. Mon. vi, p. 29 (1862); Seitz (*Notocrypta*), Macrolep. ix, p. 1094 (1927).

Me Song forest, Prae, N. Siam and Sriracha forest, S. E. Siam (*Godfrey*); Me Lamoung, W. Siam (*Stockley*).

Distribution : Ceylon; S. India; Sikkim to Burma; China; Siam; Malay Peninsula; Sumatra; Borneo; Java; Philippines.

630. Gangara thyrsis thyrsis (Fabr.).

Papilio thyrsis Fabricius, Syst. Ent. p. 532 (1775).

Gangara thyrsis thyrsis Seitz, Macrolep. ix, p. 1071, figs. 170 a, b (1927).

Sriracha, S. E. Siam and Bangkok, C. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*).

Distribution : Sikkim to Burma; Tonkin; Siam; the Malay Peninsula and Archipelago.

Subspecies are known from Ceylon, the Andamans, and the Philippines.

631. **Erionota thrax** (Linn.).

Erionota thrax Linnaeus, Syst. Nat. ii, p. 764 (1767).

Distribution: India to China, south to the Sunda and Molucca Islands.

(a) **Erionota thrax thrax** (Linn.).

Erionota thrax thrax Seitz, Macrolep. ix, p. 1071, fig. 107 a (1927).

Widely spread over Siam.

Distribution: India; Burma; Siam; Malay Peninsula and Archipelago; Philippines.

(b) **Erionota thrax thrax f. acroleuca** (W-M. & de Nicév.).

Telegonus acroleucus Wood-Mason and de Nicéville, Proc. As. Soc. Bengal, p. 143 (1881).

Bangkok, C. Siam (*Godfrey*; *Aagaard*).

Distribution: Occurs as a rare variety of *thrax* from Assam to Burma, in Siam, the Malay Peninsula, and in Borneo.

632. **Matapa aria** (Moore).

Ismene aria Moore, Proc. Zool. Soc. Lond. p. 784 (1865); Seitz (*Matapa*), Macrolep. ix, p. 1067, figs. 169 d, e (1927).

Sriracha, S. E. Siam and Bangkok, C. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*); Me Wong W. Siam (*Stockley*).

Distribution: Ceylon; S. India; Sikkim to Burma; China; Tonkin; Hainan; Siam; Malay Peninsula and Archipelago; Philippines.

633. **Matapa sasivarna** (Moore).

Ismene sasivarna Moore, Proc. Zool. Soc. Lond. p. 784 (1865); Seitz (*Matapa*), Macrolep. ix, p. 1068, fig. 169 f (1927).

1 ♂ : Me Song forest, Prae, N. Siam, April 1918.

Distribution: Sikkim to Burma; Siam; Malay Peninsula; Sumatra.

634. **Matapa shalgrama** de Nicév.

Matapa shalgrama de Nicéville, Journ. As. Soc. Beng. p. 85 (1883); Seitz (*Matapa*), Macrolep. ix, p. 1068, fig. 169 f (1927).

1 ♂ : Me Song forest, Prae, N. Siam, April 1918.

Distribution: Sikkim to Burma; Andamans; Tonkin; Siam; Java; Bali; Pulo Laut.

635. **Hyarotis adrastus praba** (Moore).

Papilio adrastus Cramer, Pap. Exot. iv, pl. 319, figs. F, G (1780).

Hyarotis adrastus praba Seitz, Macrolep. ix, p. 1066, figs. 169 a, b (1927).

Me Song forest, Prae and Chiengrai, N. Siam; Muak Lek, E. Siam; Bangkok, C. Siam.

Distribution: Kangra (Punjab) to Burma; Andamans; Hong Kong; Siam.

The species ranges from India and Ceylon to the Malay Peninsula, Pulo Condore, Sumatra, Borneo, and Java.

636. *Zographetus satwa* (de Nicév.).

Isoteinon satwa de Nicéville, Journ. As. Soc. Beng. p. 86 (1883); Seitz (*Zographetus*), Macrolep. ix, p. 1067, fig. 169 c (1927).

Chiengrai, N. Siam; Pak Jong, E. Siam; Sriracha, S. E. Siam.

Distribution: Sikkim to Burma; Siam; Malay Peninsula; Sumatra; Java.

637. *Zographetus ogygia ogygia* (Hew.).

Hesperia ogygia Hewitson, Trans. Ent. Soc. Lond. p. 500 (1866); Seitz (*Zographetus*), Macrolep. ix, p. 1067, fig. 169 d (1927).

Zographetus ogygia ogygia Evans, Journ. Bomb. N. H. S. xxxi, p. 442 (1926).

Me Song forest, Prae, N. Siam and Muak Lek, E. Siam (*Godfrey*); Me Wong, W. Siam (*Stockley*).

Distribution: Sikkim to Burma; S. India; Siam; Malay Peninsula and Archipelago.

Evans has described a subspecies, *Z. o. andamana* from the Andamans.

638. *Isma protoclea* (H.-Schäf.).

Goniloba protoclea Herrich-Schäffer, Prodr. Syst. Lep. p. 195 (1869); Seitz, Macrolep. ix, p. 1066, fig. 169 b (1927).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Karen Hills to S. Burma; Siam; Malay Peninsula; Java; Borneo.

639. *Isma vulso bicolor* Evans.

Pamphila vulso Mabille, Ann. Soc. Ent. Belg. xxxvii, p. 55 (1893).

Isma vulso bicolor Evans, Journ. Bomb. N. H. S. xxxi, p. 443 (1926).

Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: Evans (*l. c.*) refers this subspecies to the Karen and Dawna Hills and to the Malay Peninsula. He gives Sumatra,

Java and Bali as the habitat of the typical form and recognizes *purpurascens* Elw. & Edw. as the Bornean race.

Seitz (p. 1066) treats *vulso* as a subspecies of *protoclea* H.-Schäff.

640. Sepa feralia (Hew.).

Hesperia feralia Hewitson, Descr. Hesp. p. 31 (1868); Seitz (*Scobura*), Macrolep. ix, p. 1069, figs. 169 h, i (1927).

Khao Ram, Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: P. Siam; Malay Peninsula; Sumatra; Borneo; Java; Pulo Laut.

641. Sepa fenestra (Elw. & Edw.).

Scobura fenestra Elwes and Edwards, Trans. Zool. Soc. Lond. p. 206 (1897); Seitz (*Scobura*), Macrolep. ix, p. 1069, fig. 169 i (1927).

Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: S. Mergui; P. Siam; Malay Peninsula; Sumatra; Borneo; Java; Pulo Laut.

642. Sepa inarime (de Nicév.).

Isma inarime de Nicéville, Journ. Bomb. N. H. S. p. 391 (1891); Seitz (*Scobura*), Macrolep. ix, p. 1069, fig. 169 i (1927).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Mergui; Siam; Malay Peninsula; Sumatra; Java; Borneo; Pulo Laut.

643. Plastingia callineura burmana Evans

Hesperia callineura Felder, Nov. Reise Lep. iii, p. 573 (1866); Seitz (*Plastingia*), Macrolep. ix, p. 1092 (1927).

Plastingia callineura burmana Evans, Journ. Bomb. N. H. S. xxxi, p. 444 (1926).

Kao Sabap, Chantabun, S. E. Siam.

Distribution: Assam to Burma; Siam.

The species ranges from Assam to the Malay Peninsula, Sumatra, Nias, Java, Borneo and Palawan.

644. Plastingia latoia latoia (Hew.).

Hesperia latoia Hewitson, Descr. Hesp. p. 34 (1868).

Plastingia latoia latoia Evans, Journ. Bomb. N. H. S. xxxi, p. 444 (1926).

Plastingia callineura latoia Seitz, Macrolep. ix, p. 1092, fig. 174 e (1927).

Hup Bon, Sriracha forest, S. E. Siam (*Godfrey*); Nakon Sritamarat, P. Siam (*Kerr*).

Distribution: Dawna Hills to S. Burma; Siam; Malay Peninsula; Sumatra; Java; Borneo; Labuan; Philippines.

Evans (*l. c.*) recognizes as a good subspecies *P. l. margherita* Doh. (Upper Assam; Tenasserim).

Seitz (*l. c.*) treats *latoia* as a subspecies of *callineura* Feld., and refers it to Malacca.

645. *Plastingia sala* (Hew.).

Hesperia sala Hewitson, Trans. Ent. Soc. Lond. p. 500 (1866); Evans (*Plastingia*), Journ. Bomb. N. H. S. xxxi, p. 445 (1926); Seitz (*Suastus*), Macrolep. ix, p. 1062 (1927).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Assam to Burma; Siam; Malay Peninsula.

According to Evans (op. cit., p. 434), *sala* does not belong to the genus *Suastus* (in which Seitz places it), but is a *Plastingia*.

646. *Plastingia naga* (de Nicév.).

Hesperia naga de Nicéville, Journ. As. Soc. Beng. p. 89 (1883); Seitz, (*Plastingia*), Macrolep. ix, p. 1091, fig. 174 a (1927).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Assam to Burma; Siam; Malay Peninsula; Sumatra; Borneo; Java; (?) Philippines.

647. *Lotongus calathus* (Hew.).

Eudamus calathus Hewitson, Ann. Mag. Nat. Hist. (4) xvii, p. 353 (1876); Seitz (*Lotongus*), Macrolep. ix, p. 1098, fig. 175 e (1927).

Me Song forest, Prae, N. Siam; Sriracha, S. E. Siam; Khanduli, P. Siam (*Godfrey*); Kao Pawta Luang Keo, c. 900 m., Ranawng (Renong), P. Siam (*Kerr*).

Distribution: Burma to the Malay Peninsula; Sumatra; Borneo; Java; Palawan.

648. *Lotongus schaedia* (Hew.).

Hesperia schaedia Hewitson, Ann. Mag. N. H. (4) xviii, p. 452 (1876); Evans (*Lotongus*), Journ. Bomb. N. H. S. xxxi, p. 446 (1926).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Siam; Malay Peninsula; Sumatra; Nias; Borneo.

Seitz treats *schaedia* as a synonym of *calathus*, but both Riley and Evans accept it as a good species.

649. *Zela zeus optimus* Fruhst.

Zela zeus de Nicéville, Journ. Bomb. N. H. S. ix, p. 338 (1895).

Zela zeus optimus Evans, Journ. Bomb. N. H. S. xxxi, p. 615 (1926).

Lotongus zeus optimus Seitz, Macrolep. ix, p. 1099 (1927).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Assam; Tavoy; Siam.

The species ranges from Assam to Sumatra, Borneo, Pulo Laut, and Mindoro.

650. *Acerbas anthea* (Hew.).

Hesperia anthea Hewitson, Descr. Hesp. p. 29 (1868); Seitz (*Acerbas*), Macrolep. ix, p. 1070 (1927).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Burma to the Malay Peninsula and Archipelago.

651. *Unkana attina* (Hew.) [= *batara* Moore].

Hesperia attina Hewitson, Trans. Ent. Soc. Lond. p. 489 (1866); Seitz (*Unkana*), Macrolep. ix, p. 1099, fig. 175 g (1927).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: India (Bassein) to S. Burma; Siam; Malacca; Sumatra; Borneo; Java; and Palawan.

The name *batara* was published by Moore in 1857 in Cat. Lep. Mus. E. I. C., but without a description of the species. It was stated that a description would be published later but, as this never appeared, Hewitson's name *attina*, published in 1865, has priority.

652. *Hidari irava* (Moore).

Hesperia irava Moore, Cat. Lep. Mus. E. I. C. p. 254 (1857); Seitz (*Hidari*), Macrolep. ix, p. 100, fig. 175 h (1927).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution: Burma; Siam; Malacca; Sumatra; Borneo; Java.

653. *Pirdana hyela* (Hew.).

Hesperia hyela Hewitson, Descr. Hesp. p. 23 (1867); Evans (*Pirdana*), Journ. Bomb. N. H. S. xxxi, p. 617 (1926); Seitz (*Pirdana*), Macrolep. ix, p. 1093, fig. 174 e (1927).

Me Song forest, Prae, N. Siam and Sriracha forest, S. E. Siam (*Godfrey*); Chiengmai, N. Siam (*Tennent*).

Distribution: Assam to Burma, China, Hainan and Palawan; south to the Malay Peninsula and Archipelago.

Under *hyela*, Evans sinks *rudolphii* Elw. and de Nicév., and *scanda* Fruhst.

654. *Pithauria murdava* (Moore).

Ismene murdava Moore, Proc. Zool. Soc. Lond. p. 784 (1865); Seitz (*Pithauria*), Macrolep. ix, p. 1091, fig. 171 h (1927).

6 ♂ ♂ : Phar Mieng, near Chiengrai, N. Siam, April 1921.

Distribution : Sikkim to Burma; Siam; Malay Peninsula; Borneo.

655. *Halpe albipectus* de Nicév.

Halpe albipectus de Nicéville, Journ. Bomb. N. H. S. p. 389 (1891); Seitz, Macrolep. ix, p. 1090, fig. 171 g (1927).

Me Song forest, Prae, N. Siam (*Godfrey*); Me Wong, W. Siam (*Stockley*).

Distribution : Shan States to S. Burma; Siam.

656. *Halpe cerata* (Hew.).

Hesperia cerata Hewitson, Ent. Mo. Mag. p. 152 (1876); Seitz (*Halpe*), Macrolep. ix, p. 1088 (1927).

Me Song forest, Prae, and Chiengrai, N. Siam.

Distribution : Sikkim to Burma; Siam; (?) Philippines.

657. *Halpe masoni* (Moore).

Pamphila masoni Moore, Proc. Zool. Soc. Lond. p. 842 (1878); Seitz (*Halpe*), Macrolep. ix, p. 1090, fig. 171 f (1927).

1 ♂ : Me Song forest, Prae, N. Siam, April 1916 (*Godfrey*);
1 ♂ : Klong Pong Kapo, W. Siam (*Stockley*).

Distribution : Chin Hills to S. Burma; Tonkin; Siam.

658. *Halpe burmana* Swinh.

Halpe burmana Swinhoe, Lep. Ind. x, p. 269 (1913); Evans, Journ. Bomb. N. H. S. xxxi, p. 620 (1926).

7 ♂ ♂ , 4 ♀ ♀ : Me Song forest, Prae, N. Siam, April 1916;
2 ♂ ♂ , 1 ♀ : April 1918 (*Godfrey*); 1 ♂ : Klong Pong Kapo, W. Siam, April 1920 (*Stockley*).

Distribution : Burma; Siam.

Seitz makes no mention of this species.

659. *Halpe moorei* Wats. (= *beturina* Fruhst.).

Halpe moorei Watson, Proc. Zool. Soc. Lond. p. 109 (1893).

Halpe moorei beturina Seitz, Macrolep. ix, p. 1088 (1927).

Me Song forest, Prae, and Chiengrai, N. Siam; Muak Lek, E. Siam; Sriracha forest, S. E. Siam; Bangkok, C. Siam.

Distribution: Mussoorie to Burma; Hongkong; Siam; Annam; Sumatra; Borneo; Celebes.

660. *Halpe zema zema* (Hew.).

Hesperia zema Hewitson, Ann. Mag. N. H. p. 77 (1877).

Hesperia zema zema Seitz, Macrolep. ix, p. 1088, fig. 173 b (1927).

Me Song forest, Prae, N. Siam (*Godfrey*); Klong Pong Kapo, W. Siam (*Stockley*).

Distribution: Sikkim to S. Burma; Tonkin; Siam; Malay Peninsula.

The species ranges from Sikkim to Sumatra, Nias, Java, and Palawan.

661. *Halpe sikkima sikkima* Moore.

Halpe sikkima Moore, Proc. Zool. Soc. Lond. p. 407 (1882).

Halpe sikkima sikkima Seitz, Macrolep. ix, p. 1089, fig. 171 b (1927).

Sriracha forest, S. E. Siam and Bangkok, C. Siam (*Godfrey*); Klong Pong Kapo, W. Siam (*Stockley*).

Distribution: Sikkim to Burma; Siam; Malay Peninsula; Sumatra; Borneo.

Subspecies are found in Nias and Palawan.

662. *Halpe homolea homolea* (Hew.).

Hesperia homolea Hewitson, Descr. Hesp. p. 29 (1868).

Halpe homolea homolea Evans, Journ. Bomb. N. H. S. xxxi, p. 622 (1926); Seitz, Macrolep. ix, p. 1089, fig. 171 d (1927).

Sriracha forest, S. E. Siam; Chiengrai, N. Siam.

Distribution: Sikkim to Burma; Siam; Singapore.

Evans (*l. c.*) recognizes as a good race *H. h. ceylonica* Moore, from Ceylon.

663. *Onryza siamica* Ril. & Godf.

Onryza siamica Riley and Godfrey, Entomol. lviii, p. 142, pl. 3, fig. 5 (1925).

Halpe meiktila de Nicéville, Journ. Bomb. N. H. S. p. 57 (1891); Evans (*Onryza*), op. cit., xxxi, p. 622 (1926).

Suan Dawk and Phar Mieng, near Chiengrai, N. Siam.

Distribution: N. Siam.

Evans treats *siamica* as a subspecies of *meiktila* de Nicév. (Meiktila to the Dawna Hills).

664. Cupitha purreea (Moore).

Pamphila purreea Moore, Proc. Zool. Soc. Lond. p. 594 (1877); Seitz (*Cupitha*), Macrolep. ix, p. 1093 (1927).

Me Song forest, Prae, N. Siam; Pak Jong and Muak Lek, E. Siam.

Distribution : S. India; Sikkim to Burma; Andamans; Siam; Malay Peninsula and Archipelago; Philippines.

665. Taractrocera maevius flaccus (Fabr.) [= *lineata* Druce].

Hesperia maevius Fabricius, Ent. Syst. iii, p. 352 (1793); Seitz (*Taractrocera*), Macrolep. ix, p. 1074 (1927).

Taractrocera maevius flaccus Evans, Journ. Bomb. N. H. S. xxxi, p. 624 (1926).

Nakon Chaisee, W. Siam (Druce); Muak Lek, E. Siam and Bangkok, C. Siam (Godfrey).

Distribution : India; Burma; Siam.

According to Evans, the typical form occurs in Ceylon.

T. lineata was originally described from Siam.

Seitz sinks *flaccus* under *maevius*.

666. Taractrocera ziclea samadha Fruhst.

Thymelicus ziclea Plötz, Stett. Ent. Zeit. xlv, p. 289 (1884).

Taractrocera ziclea samadha Seitz, Macrolep. ix, p. 1075 (1927).

Chiengrai, N. Siam; Bangkok, C. Siam.

Distribution : Burma; Siam.

The species ranges from Burma to the Philippines, south to the Malay Archipelago.

667. Taractrocera archias quinta Swinh.

Pamphila archias Felder, Sitz.-Ber. Ak. Wiss. Wien. xl, p. 462 (1860); Seitz (*Taractrocera*), Macrolep. ix, p. 1075 (1927).

Taractrocera archias quinta Evans, Journ. Bomb. N. H. S. xxxi, p. 625 (1926).

Doi Chang (4,500 ft.), N. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam.

Distribution : Burma; Siam; Annam; Malay Peninsula.

The species ranges from Burma to the Malay Archipelago.

668. *Padraona bola* (Moore).

Padraona bola Moore, Proc. Zool. Soc. Lond. p. 596 (1877); Evans (*Nicervillea*), Journ. Bomb. N. H. S. xxxi, p. 623 (1926); Seitz (*Padraona*), Macrolep. ix, p. 1077, fig. 170 i (1927).

Muak Lek, E. Siam; Sriracha, S. E. Siam; Bangkok, C. Siam.

Distribution: India to China and the Philippines, south to the Malay Peninsula and Archipelago.

669. *Padraona rectifasciata rectifasciata* (Elw. & Edw.).

Telicota rectifasciata Elwes and Edwards, Trans. Zool. Soc. Lond. xiv, p. 254 (1897).

Padraona rectifasciata rectifasciata Seitz, Macrolep. ix, p. 1078, fig. 168, h (1927).

Padraona kobros rectifasciata Evans, Journ. Bomb. N. H. S. xxxi, p. 626 (1926).

2 ♂♂ : Me Song forest, Prae, April 1918.

Distribution: Karen Hills to S. Burma; Siam.

Seitz recognizes subspecies from Java and Nias.

Evans (*l. c.*) treats *rectifasciata* as a subspecies of *kobros* Plötz (Burma to New Guinea and Australia).

670. *Padraona moesoides moesoides* (Butl.).

Pamphila moesoides Butler, Trans. Linn. Soc. Zool. p. 554 (1879).

Padraona moesoides moesoides Evans, Journ. Bomb. N. H. S. xxxi, p. 626 (1926).

Padraona dara moesoides Seitz, Macrolep. ix, p. 1077, fig. 171 i (1927).

Chiengrai, N. Siam; Nam Phi, N. E. Siam; Muak Lek, S. E. Siam; Khanduli, P. Siam.

Distribution: Dawna Hills to S. Burma; Siam; Malay Peninsula; Sumatra; Banka.

Evans has described a subspecies, *P. m. ottalina*, from the Andamans.

Seitz (*l. c.*) treats *moesoides* as a subspecies of *dara* Koll. and refers it to Malacca.

671. *Padraona dara serina* Plötz.

Hesperia dara Kollar, in Hugel's Kaschmir iv, p. 455 (1848); Seitz (*Padraona*), Macrolep. ix, p. 1076 (1926).

Padraona dara serina Evans, Journ. Bomb. N. H. S., p. 627 (1926).

Widely spread over Siam.

Distribution : S. Burma ; Siam ; Malay Peninsula ; Sumatra ; Java ; Lombok ; Nias.

In addition to *P. d. serina*, Evans (*l. c.*) recognizes as distinct races : *P. d. dara* Koll. (Chitral to Kumaon) ; *P. d. pseudomoesa* Moore (Ceylon ; S. India) ; *P. d. confucius* Feld. (Mussoorie to Shan States and China) ; and *P. d. dilutior* (Borneo ; Pulo Laut).

Seitz makes no mention of *serina*.

672. Padraona sunias tropica (Plötz).

Hesperia tropica Plötz, Stett. Ent. Zeit. p. 230 (1883).

Padraona sunias tropica Evans, Journ. Bomb. N. H. S. xxxi, p. 627 (1926).

Telicota tropica tropica Seitz, Macrolep. ix, p. 1079, fig. 172 a (1927).

2 ♂ ♂, 1 ♀ : Chiengrai, N. Siam, April 1921.

Distribution : S. India ; Sikkim to Burma, Siam, the Malay Peninsula and Archipelago.

Evans refers typical *sunias* to the Moluccas.

Seitz regards both *tropica* and *sunias* as separate species.

673. Padraona nitida (Mab.).

Pamphila nitida Mabille, Petit. Nouv. ii, p. 114 (1877) ; Evans (*Padraona*), Journ. Bomb. N. H. S. xxxi, p. 627 (1926).

Padraona dara nitida Seitz, Macrolep. ix, p. 1077, fig. 170 i (1927).

Bangkok, C. Siam (*Godfrey*).

Distribution : Ataran and Tavoy; Siam ; Philippines.

674. Padraona trachala (Mab.).

Padraona trachala Mabille, Petit. Nouv. i, p. 237; Evans (*Padraona*), Journ. Bomb. N. H. S. xxxi, p. 628 (1926).

Padraona dara trachala Seitz, Macrolep. ix, p. 1077, figs. 170 h, i (1926).

Nakon Sritamarat, P. Siam (*in F. M. S. Museums' Coll.*).

Distribution : Assam to Burma ; P. Siam ; Malay Peninsula ; Sumatra ; Borneo ; Java.

Seitz (*l. c.*) treats *trachala* as a subspecies of *dara* and refers it to Java.

675. Telicota augias augias (Linn.).

Papilio augias Linnaeus, Syst. Nat. i, p. 794 (1767).

Telicota augias augias Seitz, Macrolep. ix, p. 1079, fig. 171 b (1927).
 Sriracha, S. E. Siam and Bangkok, C. Siam (*Godfrey*); Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: India to China, Formosa, Hongkong, and the Philippines; south to the Malay Archipelago.

A subspecies is known from N. Australia.

676. *Telicota pythias bambusae* (Moore).

Pamphila pythias Mabille, Pet. Nouv. ii, p. 234 (1878).

Pamphila bambusae Moore, Proc. Zool. Soc. Lond. p. 691 (1878); Seitz (*Telicota*), Macrolep. ix, p. 1078 (1927).

Telicota pythias bambusae Evans, Journ. Bomb. N. H. S. xxxi, p. 628 (1926).

Sriracha, S. E. Siam; Bangkok, C. Siam.

Distribution: Ceylon; India; Burma; Andamans; China; Hainan; Tonkin; Siam; Annam.

According to Evans, the species ranges from India to New Guinea and Australia.

677. *Telicota palmarum palmarum* (Moore).

Pamphila palmarum Moore, Proc. Zool. Soc. p. 690 (1878); Evans (*Telicota*), Journ. Bomb. N. H. S. xxxi, p. 269 (1926).

Telicota palmarum palmarum Seitz, Macrolep. ix, p. 1080, figs. 172 a, b (1927).

Doi Chang (4,500 ft.), Chiengrai, N. Siam; Bangkok, C. Siam.

Distribution: Calcutta to Assam; Burma; Siam; Malay Peninsula and Archipelago.

The species ranges from India to Australia.

678. *Augiades siva* (Moore).

Pamphila siva Moore, Proc. Zool. Soc. Lond. p. 692 (1878); Seitz (*Augiades*), p. 1082, fig. 172 f (1927).

♂♂: Doi Chang (4,500 ft.), Chiengrai, N. Siam, April 1921.

Distribution: Assam to Burma; Siam.

679. *Iton semamora* (Moore).

Hesperia semamora Moore, Proc. Zool. Soc. Lond. p. 791 (1865); Seitz (*Iton*), Macrolep. ix, p. 1090 (1927).

Me Song forest, Prae, N. Siam (*Godfrey*); Nakon Sritamarat, P. Siam (*Pendlebury*).

Distribution: Sikkim to Burma; Siam; Malay Peninsula;

Sumatra ; Borneo ; Philippines.

680. Iton watsoni (de Nicév.).

Parnara watsoni de Nicéville, Journ. Bomb. N. H. S. p. 223 (1890) ; Seitz (*Iton*), Macrolep. ix, p. 1070, fig. 171 f (1927).

Me Song forest, Prae, N. Siam (*Godfrey*) ; Kanburi, W. Siam (*Fruhstorfer*).

Distribution : Burma ; Siam ; Perak.

681. Baoris oceia farri Moore.

Baoris oceia Hewitson, Descr. Hesp. p. 31 (1868) ; Seitz (*Parnara*), Macrolep. ix, p. 1086 (1927).

Baoris farri Moore. Proc. Zool. Soc. Lond. p. 688 (1878).

Baoris oceia farri Evans, Journ. Bomb. N. H. S. xxxi, p. 632 (1926).

Distribution : S. India ; Sikkim to Burma ; Andamans ; Tonkin ; Hongkong ; Siam ; Malay Peninsula ; Sumatra ; Nias ; Java.

Evans refers typical *oceia* to the Philippines, and recognizes subspecies from China, Ceylon and Borneo.

682. Baoris brunnea (Snell.).

Pamphila brunnea Snellen, Titj. v Ent. xix, p. 164 (1876) ; Evans (*Baoris*), Journ. Bomb. N. H. S. xxxi, p. 632 (1926) ; Seitz (*Parnara*), Macrolep. ix, p. 1087, fig. 173 a (1927).

Hup Bon, Sriracha forest, S. E. Siam.

Distribution : Sikkim to Burma ; Siam ; Annam ; Java ; Bali ; Borneo.

683. Baoris mathias (Fabr.).

Hesperia mathias Fabricius, Ent. Syst. Suppl. p. 433 (1798) ; Seitz (*Parnara*), Macrolep. ix, p. 1087 (1927).

Widely spread over Siam.

Distribution : Throughout the Indo-Australian region.

684. Baoris pagana (de Nicév.).

Parnara pagana de Nicéville, Proc. Zool. Soc. Lond. p. 465 (1887) ; Seitz, Macrolep. ix, p. 1085, fig. 173 c (1927).

Baoris pagana Evans, Journ. Bomb. N. H. S. xxxi, p. 633 (1926).

Me Song forest, Prae, N. Siam ; Bangkok, C. Siam.

Distribution : Sikkim to Burma ; Siam ; Sumatra ; Borneo.

685. Baoris cahira (Moore).

Hesperia cahira Moore, Proc. Zool. Soc. Lond. p. 593 (1877) ; Seitz (*Parnara*), Macrolep. ix, p. 1085, fig. 173 c (1927).

Distribution: Sikkim to W. China, Tonkin, and the Philippines; south to the Malay Peninsula and Archipelago.

(a) **Baoris cahira nirwana** Plötz

Baoris cahira nirwana Evans, Journ. Bomb. N. H. S. xxxi, p. 633 (1926).

Chiengmai, N. Siam (*Tennent*); Me Lamoung, W. Siam (*Stockley*).

Distribution: Tonkin; Siam; Malay Peninsula and Archipelago.

(b) **Baoris cahira sirius** Evans

Baoris cahira sirius Evans, op. cit. p. 633.

Pak Jong, E. Siam; Bangkok, C. Siam.

Distribution: Burma; Siam.

686. Baoris cormasa (Hew.).

Hesperia cormasa Hewitson, Ann. Mag. Nat. Hist. (4) xviii, p. 457 (1876); Seitz (*Parnara*), Macrolep. ix, p. 1085 (1927).

1 ♂: Bangkok, C. Siam, 23. 9. 13; 1 ♂: 21. 7. 27.

Distribution: Sikkim to Burma; Siam; Malay Peninsula and Archipelago.

687. Baoris tulsi (de Nicév.).

Parnara tulsi de Nicéville, Journ. As. Soc. Beng. p. 86 (1883); Seitz, Macrolep. ix, p. 1086, fig. 173 e (1927).

Nakon Sritamarat, P. Siam (*in F. M. S. Museums' Coll.*).

Distribution: Sikkim to Burma; P. Siam; Java.

688. Baoris eltola (Hew.).

Hesperia eltola Hewitson, Exot. Butt iv, Hesp. pl. 4, fig. 40 (1869); Seitz (*Parnara*), Macrolep. ix, p. 1084, fig. 173 b (1927).

Me Song forest and Doi Chang (4,500 ft.), N. Siam; Sriracha, S. E. Siam.

Distribution: Mussoorie to the Karen Hills; Siam.

689. Baoris contigua (Mab.).

Pamphila contigua Mabille, Bull. Soc. Zool. France, p. 232 (1877); Seitz (*Parnara*), Macrolep. ix, p. 1084, fig. 173 e (1927).

Me Song forest and Chiengsen, N. Siam; Muak Lek, E. Siam; Khanduli, P. Siam (*Godfrey*); Umpang, W. Siam (*Stockley*).

Distribution: India to China, Tonkin and Formosa; south to the Malay Peninsula and Archipelago.

690. *Baoris guttatus bada* (Moore).

Eudamus guttatus Bremer and Grey, Schmett. N. China, p. 10 (1853).

Hesperia bada Moore, Proc. Zool. Soc. Lond. p. 688 (1878).

Baoris guttatus bada Evans, Journ. Bomb. N. H. Soc. xxxi, p. 636 (1926).

Parnara guttatus bada Seitz, Macrolep. ix, p. 1083 (1927).

Widely spread over Siam.

Distribution: India to Tonkin, Hainan, Formosa and the Philippines, south to Malaya.

Evans gives the range of typical *guttatus* as:—"Chitral to Assam Hills, 2,900 ft, China and Japan."

691. *Baoris zelleri colaca* (Moore).

Hesperia zelleri Lederer, Verh. Zool.-bot. Ges. Wien. p. 194 (1855).

Hesperia colaca Moore, Proc. Zool. Soc. Lond. p. 594 (1877); Seitz, Macrolep. ix, p. 1084, fig. 173 a (1927).

Baoris zelleri colaca Evans, Journ. Bomb. N. H. S. xxxi, p. 636 (1926).

1 ♂, 1 ♀ : Chiengrai, N. Siam, 1. 5. 21.

Distribution: Evans (*l. c.*) gives the range of *colaca* as:—"Ceylon, India, Burma, Andamans, Nicobars, Formosa, Hainan, Malay Peninsula and Islands, Philippines, Moluccas, N. Australia."

Evans refers typical *zelleri* to Syria and N. Africa.

692. *Baoris bevani* (Moore).

Hesperia bevani Moore, Proc. Zool. Soc. Lond. p. 688 (1878); Seitz (*Parnara*), Macrolep. ix, p. 1084, figs. 173 a, b (1927).

1 ♂ : Chiengrai, N. Siam, 30. 4. 21.

Distribution: Sikkim to Borneo, Palawan, and Formosa.

INDEX *

to

A REVISED LIST OF SIAMESE BUTTERFLIES.

A	alemon 334 alesia 282 alexis 353 alica 361 alida 366 alinkara 373 aliris 262 alitaeus 331 allica 203, 305 Allotinus 312 almama 272 almea 354 alompra 305 alyisia 373 alysos 372, 373 amabilis 348 amantes 335 amarra 356 Amathusia 259 Amathuxidia 260 amatius 348 amatrix 335 amba 351 ambica 297 Amblypodia 329 ambrysus 290 ameria 328 ameithystus 260 amharina 286 ammoq 337 ammonoides 337 Ampittia 368 amrita 343 amythaon 260 Anadebis 247 anadi 355 ananga 317 anapita 244 anarte 329 anaxias 251 anaxioides 251 aneon 312 andamana 375 andam nica (Arhopala) 339 andamanica (Daimio) 363 andersoni 291 angelica 300 angularis 244 angulata (Abisara) 306 angulata (Odontoptilum) 366 angusta 320 anita 327 annamita, 295 	anthea (Acerbas) 378 anthelus 329 antimuta 336 antiphates 216 anunda 329 anura 352 aoris 268 apama 292 Apatura 297, 298 apelles 256 Aphnaeus 341 aphya 322 Appias 225 arama 332 Araotes 352 arca 264 arcesilaus 203, 257 archias 381 arcturus 215 argentina 244 argiades 316 argiola 318 Argynnis 271 argyridina 226 Arhopala 329 aria 374 aricia 330 ariadne 263 ariana 332 aristeus 216 aristogiton 303 aristolochiae 208 arja 301 Arnetta 368 area 333
---	--	---

* Names of all genera printed in black type in the text are included in this index, but not their synonyms (printed in italics). All specific and subspecific names are fully indexed.

atraz 331
atticus 359, 360, 362
attina 378
atymnus 339
Augiades 384
augias 383
aurelia (Rahinda) 278
aurelia (Amblypodia) 334
aurelius 261
aurivittata 359
avalokita 299
avanta 243
avatar 231
avattana 373
axion 217
Azanus 325

B

bada 387
Badamia 357
badra 353
balana 360
baldus 243
bambooae 384
Baoris 385
barata 207
baronesa 253
barroni 357
batara 378
bataviana 234
bathycles 218
bathycloides 218
batina 341
batmara 247
bazalus 335
beata 294
belisama 300
belladonna 222
bellata 295
benjamini 356
berenice 322
berthae 309
beturina 379
bevani 387
bhagava 363
bhavama 298
bhima 247
Bibasis 356
biblis 270
bicolor 375
Bidiunda 344
bifasciatus 359
biggsii 312
Bindahara 352
binghami (Penthema) 266
binghami (Dodona) 305
bipunctus 369
birmana 337
bisaltide 275
blanda 230
boehus 323
boeticus 320
boisduvali (Miletus) 312

boisduvali (Marmessus) 345
bolina 275
borneana 343
Bothrinia 316
bouletti 308
bowringi 364
bradamantha 311
brahma 242
brassolis 352
brunnea 385
buddha 298, 299
bulis 326, 327
burmania (Gandaca) 231
burmania (Plastingia) 376
burmania (Halpe) 379
busiris 258
butleri 370
butlerianus 214

C

cahra 385
calathus 377
caleta 315
Calinaga 298
callineura 376, 377
calydonia 300
cama 286
camadeva 258
camadevoidea 258
camaralzaman 238
Cambodia 259
camdeo 329
cameroni 359
canace 273
canidia 224
Capila 357
Caprona 366
cardia 318
cartica 282
Castalius 315
castelnaui 265
castor 211
easyapa 359
Catapoecilma 346
catilla 229
Catochrysops 316, 320
Catopsis 228
Celaenorhinus 358
celebensis 373
celeno 324
centaurus 234
cephala 370
cerata 379
cerberus 206
Cethosia 270
ceylonica 380
chabrona 353
chandica 246
chandra 298
chaon 212
charaka 250
Charaxes 302
Charmion 357
chelensis 257

chenelli 316
Cheritra 344
Cheritrella 343
cheronus 218
chersias 260
Chersonesia 277
chersonesia (Hestia) 233
chersonesia (Zeuxidia) 261
chersonesia (Symbrenthia) 274
chersonesia (Euthalia) 293
Chilades 318, 319
chinensis (Heliophorus) 326
chinensis (Ctenoptilum) 366
chione 269
chiron 218
Chliaria 347, 348
chromus 353
chrysippus 233
chula 292
chuza 354
ciphus 342
circuta 238
ciris 222
Cirrochroa 268
clarissa 270
clemanthe 223
cleodus 323
clericia 285
cloanthus 217
elytia 210
cneius 316
coecles 203, 276
cocytina 291
coeytus 289
Coelites 248
coerulea 324
cognata 289
colaca 387
Coladenia 364
columella 282
confusa (Papilio) 227
confusa (Lethe) 245
confucius 383
consimilis 300
contigua 386
continentalis (Discophora) 263
continentalis (Nacaduba) 321
continentalis (Loxura) 339
coon 208
core 238
cormasa 386
cornuta 317
corus 241
crameri 237
crassa 242
crawfuri 356
crisalda 253
crisia 253
critias 253
critolaus 253
crocale 228

croton 311
Ctenoptilum 365
Cupha 266
Cupitha 381
Curetis 326
curius 220
curvifascia 373
cyanæ 270
Cyaniris 316, 317
cyanipardus 297
cyara 344
cymbia 325
cynis 232
Cynthia 267
Cyrestis 276

D

Dacalana 341
daedalion 255
Daimio 363
damajanti 307
dan 364
Danaida 234
Danaida 232
danisepa 210
daos 233
dapha 224
dara (*Elymnias*) 255
dara (*Padraona*) 382, 383
darada 227
daraxa 287
darlisa 265
Darpe 365
daruea 273
dasahara 365
dasarada 207
davidsoni (*Terias*) 230
davidsoni (*Sarangesa*) 365
dealbata 365
decidia 315
decoratus 362
deione 240
dejeani 242
delesserti 220
Delias 221
delphis 362
demoleus 211
demolion 211
deo 263
deodata 306
Deramas 311
Dercas 228
descombesi 222
Deudorix 348, 349
devadatta 372
dhyana 364
diademoïdes 247
diardi 338
dices 351
dicta 318
dilucida 260
dilutior 383
dindinga 278
diocles 371

dioctetianus 242
diores 261
dioscorides 368
diphilus 209
dirae 363
dirtea 296
Discophora 262
discrepans 256
dispar 330
distanti (*Appias*) 227
distanti (*Euploea*) 238
distantius 214
distorta 339
djata 294
Dodona 305
dohertyi (*Ypthima*) 243
dohertyi (*Simiskhima*) 310
Doleschallia 275
dolon 302
donina (*Nacaduba*) 323
donina (*Drina*) 340
dorelia 281
doson 217
doubledayi (*Papilio*) 208
doubledayi (*Dercas*) 228
doubledayi (*Euploea*) 239
doubledayi (*Zeuxidia*) 261
doubledayi (*Neurosigma*)
 288
Drina 340
druc ei 241
drumila 312
dubiosa 323
Ducapa 267
duessa 336
dufresne 240
dunya 296
durnfordi 304
duryodana 279
dyrta 245

E

ebusa 282
echerius 306
editha 259
egeon 362
elegans 346
elephenor 214
elna 315
elongata 361
elpis 324
eltola 386
elwesi 367
Elymnias 255
emalea 269
emas 346
emolus 325
Eniope 263
Eooxyliides 345
ephorus 308
epicharmus 360
epicles 326
epijarbas 348
epimete 336

epius 313
epixantha 255
epycides 239
Ergolis 263
Eriboea 301
Erionota 374
eriphyte 292
Erites 244
erot a 267
erotella 267
Erycinidae 304
erycinoides 310
erylus 348
erymanthis 266
eryx 236
ethion 315
ethologa 237
etolus 348
euanthes 270
Euchrysops 316
eudamippus 301
eudoxia 274
Eulaceura 298
eumeus 257
eumolphus 335
euplea 322
Euploea 237
euproteinor 214
eurinus 300
Euripus 299
europa 245
euryplus 218
eurysaces 324
Euthalia 288
enthymius 263
evaira 371
evelina 295
Everes 316
exclamationis 357
extensa 347

F

fabius 302
fabricii 344
falcata 270
farri 385
fasciata (*Ducapa*) 267
fasciata (*Taxila*) 309
Faunis 257
faunula 258
feisthamelii 372, 373
fenestra 376
feralia 376
ficulnea 357
flaccus 381
flanona 246
flavobrunnea 268
flegyas 305
flora 290
florella 228
folus 372
fortunatus 327
francisca 251
frankii 300

- | | | |
|---|--|--|
| <p>freja 344
fuentia 244
fugator 302
fulgida 338
fuliginosa 283
fuligo 367
fulgur 372
fulla 339
fumatus 371
fusea 325
fuseum 250
ylla 306
fyllaria 306</p> <p style="text-align: center;">G</p> <p>gaika 319
galaesus 286
galba (Appias) 225
galba (Hesperia) 367
gambara 245
gambrius 288
gana 360, 361
ganapati 254
Gandaca 231
ganessa 214
Gangara 373
garlinda 298
garuda 292
gautama 235
geoffroyi 305
Gerydus 311, 312
gigantea 343
gisca 317
glaucescens 265
glaucippe 231
gloriosa 326
godama 353
godarti (Euploea) 238
godarti (Euthalia) 290
godfreyi 259
gola 382
gonatina 280
goniopeltis 208
goodrichi 295
gopaka 371
gotama 250
grahami 292
grammica 236
grandis 302
gremius 369
grisea 312
gupta 295
guttatus 387
gyas 216
gythion 322</p> <p style="text-align: center;">H</p> <p>hadeni 232
halba 346
Halpe 379
halitherses 299
hamada 313
haquinus 309
haraldus 317</p> | <p>harina 231
harisa 355
harrisi 240
harita 281
Hasora 352
hauxwelli 305
Hebomoia 231
hecate 230
hegessipus 234
helena 206
helenus 212
helferi (Prioneris) 223
helferi (Tagiades) 360
heliodore 281
Heliophorus 326
helisa 366
henrici (Anadebis) 248
henrici (Dodona) 305
henricus 211
hermocrates 216
Herona 299
Hesperia 367
Hesperiidae 352
Hestia 232
Hestina 298
hewitsoni (Poritia) 310
hewitsoni (Amblypodia) 334
Hidari 378
hiemalis 343
hieras 302
hierta 272
himachala 247
himavantus 348
hippia 231
hippoclus 273
hippooides 225
hipponous 213
homolea 380
Horaga 546
hordonia 278
horsfieldi (Laringa) 265
horsfieldi (Precis) 271
Horsfieldia 327
huebneri 242
Huphina 224
Hyarotis 374
hyela 378
hylas 279
hylax 314
hyparete 221
hypatia 274
hyperides 366
hyperbius 271
hypermnestra (Hestia) 232
hypermnestra (Elymnias)
 255
hypochira 209
Hypolimnas 274
Hypolycaena 347
hypomuta 333
hypselis 273</p> <p style="text-align: center;">I</p> <p>Iambrix 368</p> | <p>ictis 341
Ideopsis 233
ignara 338
Ilerda 326
imperatrix 317
inachus 275
inarime 376
incerta 257
indica (Vanessa) 273
indica (Euthalia) 293
indicus 338
indochinensis 219
indosinica (Pantoporia) 283
indosinica (Curetis) 326
indra 226
indrani 364
ines 352
infumata 221
inornata 333
insularis 326, 327
intermedia (Thauria) 262
intermedia (Terinos) 269
intermedia (Rapala) 351
iphita 271
Iraota 327
irava 378
irroratus 312
isaeus 264
iswara 212
Isma 375
Ismene 354
ismene (Melanitis) 254
ismene (Lebadea) 287
Issoria 267
ithiela 222
Iton 384
ivena 256
Ixias 227</p> <p style="text-align: center;">J</p> <p>Jacoona 342
jadeitina 296
jahrides 291
jahmu 291
jaina 355
jainas 361
jalindra 342
jalysus 301
jama 367
Jamides 323
jangala 342
janus 210
japetus 361
jarbas 351
jasoda 311
javaniates 371
jugurtha 223
julii 289
jyntearna 318</p> <p style="text-align: center;">K</p> <p>Kahruba 303
kada 367
Kallima 275</p> |
|---|--|--|

kanara 283
 Kankena 324
 kansa 246
 kanwa 284
 kausambi 307
 kephisos 219
 kerala 371
 Kerana 371
 kerriana 320
 Kesava 292
 khasiana 360
 khasianus 368
 klugi 242
 kobros 382
 Koruthaiilos 370

L

lacordairei 240
 lacteola 230
 Lade 226
 laeta 229
 laius 318
 lalage 226
 lalagela 226
 lalassis 226
 Lampides 320
 langsonensis 217
 lanka 353
 laos 208
 lapithis 352
 Laringa 265
 larynna 284
 lathyi 262
 latoia 376
 Laxita 307
 laxmi 364
 layardi 238
 lea 224

Lebadea 287
 leda 254
 ledereri 241
 lemonias 272
 lena 258
 lepcha 249
 lepidea 289
 lepita 304
 Leptocircus 220
 Leptosia 220
 Lethe 245
 leucacantha 222
 leucoceera 385
 leucogony 241
 leuconoe 233
 leucothoe 219
 leucostictos 241
 libera 304
 Libthyea 304
 libthyea (Appias) 225
 libthyea (Terias) 229
 lilacinus 288
 liliana 359
 limax 363
 limbata 317
 limbatus 317

limborgi (*Euploea*) 240
 limborgi (*Kallima*) 275
 Liminitis 287
 limniace 235
 lindos 219
 lineata 381
 linteata 232
 lioneli 256
 Liphyra 352
 lisias 345
 lisides 343
 lithargyria 320
 litigiosa 362
 livens 311
 lizetta 352
 Lobocla 359
 Logania 313
 lohita 341
 lotis 266
 Lotongus 377, 378
 louisa 258
 lowi 247
 Loxura 339
 lubentina 293
 lucipor 261
 ludomia 294
 lugens 360
 luteipalpus 369
 Lycaenesthes 325
 Lycaenidae 309
 lycaenina 325
 Lycaenopsis 317
 Lycambe 325
 lyclene 308
 lyncens 232
 lynecestis 308
 lyneida 225

M

macareus 219
 mackwoodi 357
 macayi 290
 macromalayana 249
 madatta 353
 maevius 381
 maghada 279
 malha 319
 mahabina 366
 mahadeva 210
 Mahathala 328
 mahintha 354
 maineas 252
 major 346
 malaya (*Huphina*) 224
 malaya (*Megisba*) 314
 malayana (*Danaida*) 234
 malayana (*Lebadea*) 287
 malayana (*Hasora*) 354
 malayanus 211
 malayica (*Euploea*) 237
 malayica (*Curetis*) 326
 malelas 256
 mallus 311
 malsara 250

mamaja 279
 mamerta 250
 Mandarinia 253
 maneia 340
 marathus 299
 maro 368
 marcia 325
 marciana 346
 margana 355
 margherita 377
 marginata 209
 marmax 303
 Marmessus 345
 marsdeni 237
 martabana 282
 martha 287
 martini 269
 masoni (*Zeuxidia*) 260
 masoni (*Halpe*) 379
 massalia 313
 Matapa 374
 mathias 385
 matiana 347
 maura 371
 maxentius 342
 maximus 341
 maxwelli 335
 mazares 241
 medus 253
 meetana 361
 megarus 220
 meges 220
 Megisba 314
 mehala 211
 meiktila 380
 mekara 246
 melaneus 236
 melania 227
 melanooides 236
 melanippus 236
 Melanitis 254
 melampus 351
 melema 265
 melisa 344
 melissa 235
 memnon 213
 menaka 362
 menanto 360
 mendesia 313
 menetriesi 240
 meraka 282
 mergua 347, 348
 merguiana 321
 meridionalis 255
 merione 264
 merta 293
 metamuta 331
 metana 314
 metarete 222
 methora 243
 methypsea 270
 miah 283
 midamus 241
 Miletus 311

- | | |
|--|--|
| <p>milliana 363
 <i>Mimanbrix</i> 368
 <i>mimetica</i> 266
 <i>minerva</i> 245
 <i>mineus</i> 248
 <i>miniata</i> 345
 <i>minuscula</i> 365
 <i>mirabella</i> 331
 <i>miranda</i> 297
 <i>misippus</i> 274
 <i>mithila</i> 268
 <i>mnasicles</i> 248
 <i>modesta</i> 238
 <i>moesoides</i> 382
 <i>monilis</i> 295
 <i>monina</i> 292
 <i>monticola</i> 239
 <i>moolaina</i> 331
 <i>Mooreana</i> 363
 <i>moorei</i> (<i>Pantoporia</i>) 284
 <i>moorei</i> (<i>Marmessus</i>) 345
 <i>moorei</i> (<i>Halpe</i>) 379
 <i>muirheadi</i> 247
 <i>mulciber</i> 241
 <i>multicaudata</i> 328
 <i>multiguttata</i> 366
 <i>murdava</i> 379
 <i>musca</i> 368
 <i>Mycalesis</i> 248
 <i>myrrha</i> 305
 <i>mystes</i> 252</p> <p style="text-align: center;">N</p> <p><i>Nacaduba</i> 320
 <i>nadina</i> 224
 <i>naga</i> 377
 <i>nama</i> 298
 <i>namouna</i> 298
 <i>nana</i> 361
 <i>nandina</i> 279
 <i>narada</i> 328
 <i>narina</i> 304
 <i>nasaka</i> 351
 <i>nautilus</i> 250
 <i>nefte</i> 286
 <i>Neocheritra</i> 343
 <i>Neomyrina</i> 343
 <i>Neope</i> 247
 <i>neophron</i> 307
 <i>neophyta</i> 247
 <i>Neopihecopis</i> 314
 <i>Neorina</i> 247
 <i>neovisala</i> 249
 <i>nepenthes</i> 301
 <i>nephelus</i> 212
 <i>Neptis</i> 279
 <i>neptunus</i> 208
 <i>nerissa</i> 224
 <i>nero</i> 225
 <i>nesaea</i> 256
 <i>nesia</i> 279
 <i>Neurosigma</i> 288
 <i>niasana</i> 361
 <i>Nicevillea</i> 382</p> | <p><i>nicobarica</i> (<i>Curetis</i>) 326
 <i>nicobarica</i> (<i>Hypolyceaena</i>)
 348
 <i>nigrescens</i> 256
 <i>nigricans</i> 359
 <i>nigrita</i> 371
 <i>nigrobasalis</i> 301
 <i>niladana</i> 245
 <i>nilgiriensis</i> 245
 <i>ninus</i> 223
 <i>Niphanda</i> 325
 <i>nirwana</i> 386
 <i>nitida</i> 383
 <i>nivalis</i> 276
 <i>nivea</i> 276
 <i>noblei</i> 211
 <i>nolana</i> 283
 <i>noliteia</i> 315
 <i>nomius</i> 216
 <i>nonius</i> 288
 <i>nora</i> 323
 <i>nothis</i> 248
 <i>Notoecrypta</i> 372
 <i>Nymphalidae</i> 263
 <i>nymphidia</i> 309</p> <p style="text-align: center;">O</p> <p><i>oberthüri</i> 257
 <i>obseurus</i> 361
 <i>obsolecsens</i> 285
 <i>oezia</i> 385
 <i>Ochus</i> 367
 <i>ocuale</i> 272
 <i>odana</i> 262
 <i>odilina</i> 289
 <i>Odina</i> 362
 <i>Odontoptilum</i> 366
 <i>oedipodea</i> 354
 <i>oenome</i> 272
 <i>ogygia</i> 375
 <i>olivacea</i> 268
 <i>olivascens</i> 367
 <i>Onryza</i> 380
 <i>onyx</i> 346
 <i>opalina</i> (<i>Pantoporia</i>) 285
 <i>opalina</i> (<i>Amblypodia</i>) 330
 <i>optimus</i> 378
 <i>orientalis</i> 285
 <i>orissa</i> 269
 <i>orithya</i> 272
 <i>oroatis</i> 252
 <i>orontobates</i> 239
 <i>orphana</i> 308
 <i>orseis</i> (<i>Mycalesis</i>) 250
 <i>orseis</i> (<i>Rapala</i>) 350
 <i>Orthophaetus</i> 357
 <i>Orsotriaena</i> 253
 <i>osteria</i> 298
 <i>othona</i> 347
 <i>otis</i> 319
 <i>ottalina</i> 382</p> <p style="text-align: center;">P</p> <p><i>pactolus</i> 321</p> <p><i>Padraona</i> 382
 <i>paganus</i> 385
 <i>palinurus</i> 215
 <i>pallidior</i> 264
 <i>palmarum</i> 384
 <i>pandara</i> 327
 <i>pandava</i> 316
 <i>Pandita</i> 287
 <i>panope</i> 210
 <i>Pantoporia</i> 283
 <i>Papilio</i> 206
 <i>Papilionidae</i> 206
 <i>papone</i> 210
 <i>paradoxa</i> 210
 <i>paraka</i> 278
 <i>paralysos</i> 372
 <i>paramuta</i> 336
 <i>parangesa</i> 337
 <i>paravatar</i> 231
 <i>Pareba</i> 304
 <i>Pareronia</i> 231
 <i>parhasius</i> 316
 <i>paris</i> 215
 <i>parisatis</i> 297
 <i>Parnara</i> 386, 387
 <i>parra</i> 361
 <i>parthenope</i> 223
 <i>Parthenos</i> 288
 <i>pasira</i> 310
 <i>patimoka</i> 361
 <i>patna</i> 255
 <i>pavana</i> 321
 <i>pavonica</i> 311
 <i>payeni</i> 217
 <i>peducaeae</i> 225
 <i>peguanus</i> 341
 <i>pelea</i> 238
 <i>pelias</i> 367
 <i>penanga</i> 257
 <i>penicilligeria</i> 341
 <i>Penthema</i> 265
 <i>peraka</i> 278
 <i>perakana</i> (<i>Ideopsis</i>) 233
 <i>perakana</i> (<i>Cethosia</i>) 270
 <i>periander</i> 276
 <i>peridoneus</i> 303
 <i>perimuta</i> 336
 <i>perius</i> 284
 <i>perna</i> 248
 <i>perse</i> 349
 <i>perseoides</i> 249
 <i>perseus</i> 248
 <i>persimilis</i> 237
 <i>petosiris</i> 351
 <i>pfeifferae</i> 300
 <i>phaenops</i> 333
 <i>phalantha</i> 266
 <i>phalia</i> 311
 <i>pharis</i> 264
 <i>pharyge</i> 310
 <i>phedima</i> 254
 <i>phemis</i> 348
 <i>phemius</i> 293
 <i>pheretimus</i> 351</p> |
|--|--|

- | | | |
|--------------------------|--------------------------|----------------------------|
| phidippus 259 | pyrene 227 | sangaica 252 |
| philippina 348 | pythias 384 | sanguinalis 305 |
| philomela 281 | | sankara 282 |
| philoxenus 207 | Q | sankarya 353 |
| phisara 363 | quadripuncta 353 | Sarangesa 365 |
| phocides 352 | quercetorum 339 | sari 230 |
| phraatica 310 | queda 358 | saronis 326 |
| Pieris 224 | quulta 282 | sarpedon 217 |
| pila 316 | quinta 381 | sasivarna 374 |
| pinwilli 364 | | Satarupa 363 |
| Pirdana 378 | R | satropaces 290 |
| pishuna 262 | rafflesii 332 | saturnus 212 |
| pita 340 | Ragadia 253 | satwa 375 |
| Pithauria 379 | Rahinda 278 | Satyridae 242 |
| Pithecopus 314 | rahria 277 | savara 244 |
| pitmani 213 | rahrioides 277 | savitri 307 |
| placida 317 | rajuna 362 | scanda 379 |
| Plastingia 376 | rama (Amblypodia) 335 | schaedia 377 |
| plataniston 236 | rama (Suastus) 369 | schistacea 350 |
| plexippus 234 | ramosa 335 | schönherri 354 |
| plinius 319 | ranga 285 | scintilla 350 |
| polyctor 214 | ransonnettii 366 | Scobura 370, 376 |
| polyuectes (Papilio) 207 | Rapala 349 | scylla 229 |
| polyuectes (Miletus) 311 | rajaghra 360 | selanophora 285 |
| polynice 274 | ravata 342 | semanora 384 |
| Polyommatus 320 | ravi 360 | semiflava 227 |
| polytes 213 | ravina 360 | sena 356 |
| polyxena 302 | recta 295 | Sepa 376 |
| pomona 229 | rectifasciata 832 | Sephisa 298 |
| pompilius 216 | regalis 253 | septentrionis 235 |
| Poritia 309, 310, 311 | regia 332 | serina 382 |
| position 313 | reinwardti 232 | shalgrama 374 |
| praba 374 | rekka 313 | shania 280 |
| prabha 371 | Remelana 342 | sidha 373 |
| praecox 206 | repetitus 303 | siamensis (Papilio) 213 |
| pralaya 363 | reta 284 | siamensis (Huphina) 224 |
| pravara 283 | rhetenor 214 | siamensis (Hestia) 233 |
| Precis 271 | Rhinopalpa 274 | siamensis (Stichophthalma) |
| principalis 227 | rhoeucus 350 | 258 |
| Prioneris 223 | Rhopalocampa 356 | siamensis (Thauria) 262 |
| procrys 287 | rileyi 316 | siamensis (Cirrochroa) 268 |
| protenor 214 | risa 277 | siamensis (Doleschallia) |
| Prothoë 300 | robertsia 269 | 275 |
| protoecia 375, 376 | rohini 304 | siamensis (Kallima) 275 |
| proxissima 353 | rohrria 245 | siamensis (Cyrestis) 276 |
| pseudaliris 262 | rosimon 315 | siamensis (Pantoporia) 284 |
| Pseudergolis 265 | rotundata 268 | siamensis (Apatura) 297 |
| pseudoinsularis 327 | roxana 315 | siamensis (Abisara) 306 |
| pseudolaeta 229 | roxus 315 | siamensis (Miletus) 312 |
| pseudoleis 227 | rubecula 370 | siamica (Mycalesis) 251 |
| pseudomoesa 383 | rudis 250 | siamica (Eenthalia) 294 |
| pseudomonta 332, 333 | rudolphii 379 | siamica (Caprona) 366 |
| pteria 365 | | siamica (Onryza) 380 |
| pulligo 371 | S | sidra 229 |
| pura 323 | sabina 274 | sikkima (Megisba) 314 |
| Purlisa 343 | sakala 281 | sikkima (Halpe) 380 |
| purpuraseens 376 | sala 369, 370 | silhetensis 332 |
| purpurea 355 | salangana 291 | similis (Papilio) 220 |
| purreea 381 | salsala 368 | similis (Danaida) 237 |
| puseda 291 | samadha 381 | Simiskina 310 |
| puspa 317 | samatha 301 | singhapura 221 |
| pygmaeus 368 | sanatana (Mycalesis) 251 | sinha 267 |
| pyramus 223 | sanatana (Curetis) 326 | sinis 223 |
| pyranthe 228 | Saneus 371 | sinope 287 |

- | | | |
|--|---|---|
| <p>sinoria 287
 <i>Sinthusa</i> 351
 <i>sintica</i> 246
 <i>siponta</i> 253
 <i>sirius</i> 386
 <i>sita</i> 236
 <i>siva</i> 384
 <i>sivoka</i> 323
 <i>slateri</i> 209
 <i>sobrina</i> 243
 <i>sodalis</i> 230
 <i>soma</i> 280
 <i>sordice</i> 304
 <i>Spalgis</i> 313
 <i>specularia</i> 264
 <i>speithis</i> 326
 <i>sphendale</i> 329
 <i>sphinx</i> 349, 350
 <i>Spindasis</i> 340
 <i>stellifer</i> 369
 <i>Stiboges</i> 309
 <i>Stichophthalma</i> 258
 <i>strabo</i> 320
 <i>Suada</i> 370
 <i>Suasa</i> 343
 <i>Suastus</i> 369
 <i>subangulata</i> 366
 <i>subcaudata</i> 356
 <i>subfasciata</i> 329
 <i>subfuscatus</i> 371
 <i>subgrisea</i> 369
 <i>subochracea</i> 347
 <i>subtestaceus</i> 368
 <i>subvittatus</i> 367
 <i>sudassana</i> 298
 <i>suffusa</i> 349
 <i>sulphureus</i> 302
 <i>sumatrae</i> 309
 <i>sumatrana</i> 366
 <i>sunias</i> 383
 <i>sura</i> 366
 <i>Surendra</i> 339
 <i>surkha</i> 252
 <i>surya</i> 268
 <i>susrutina</i> 280
 <i>swerga</i> 370
 <i>swinhoei</i> 316
 <i>syama</i> 341
 <i>sycorax</i> 206
 <i>sylvia</i> 288
 <i>Symbrenthia</i> 273
 <i>Syntarucus</i> 319
 <i>syriethus</i> 367</p> <p style="text-align: center;">T</p> <p><i>tabrica</i> 364
 <i>Tagiades</i> 359, 364
 <i>Tajuria</i> 342
 <i>Talicada</i> 314
 <i>taminatus</i> 354
 <i>Tanaëcia</i> 288
 <i>Tapena</i> 365
 <i>Taractrocera</i> 381
 <i>Taraka</i> 313</p> | <p><i>Tarucus</i> 319
 <i>tavoyana</i> (<i>Poritia</i>) 310
 <i>tavoyana</i> (<i>Spindasis</i>) 340
 <i>tavoyanus</i> 209
 <i>Taxila</i> 308
 <i>telearchus</i> 210
 <i>telesia</i> 308
 <i>Telicota</i> 383
 <i>Terias</i> 229
 <i>Terinos</i> 269
 <i>terpander</i> 269
 <i>tessellata</i> (<i>Enispe</i>) 263
 <i>tessellata</i> (<i>Niphanda</i>) 325
 <i>teuta</i> 294
 <i>Thaduka</i> 328
 <i>Thamala</i> 345
 <i>tharis</i> 345
 <i>Thaumantis</i> 261
 <i>Thauria</i> 262
 <i>theclodes</i> 347
 <i>theodora</i> 343
 <i>Theope</i> 247
 <i>thesmia</i> 344
 <i>thestylis</i> 223
 <i>thomsoni</i> 206
 <i>thrax</i> 374
 <i>thronion</i> 226
 <i>thuisto</i> 308
 <i>thwaitesi</i> 365
 <i>Thymphia</i> 243
 <i>thydamas</i> 277
 <i>thyra</i> 222
 <i>thyrsis</i> 373
 <i>thysbe</i> 223
 <i>Ticherra</i> 344
 <i>tilaha</i> 230
 <i>timandra</i> 256
 <i>timoleon</i> 327
 <i>timora</i> 263
 <i>tira</i> 236
 <i>toba</i> 361
 <i>todara</i> 339
 <i>toungva</i> 336
 <i>trachala</i> 383
 <i>transpecta</i> 317
 <i>trichoneura</i> 363
 <i>tripunctata</i> 340
 <i>tritogenia</i> 245
 <i>triumphator</i> 214
 <i>trochilus</i> 319
 <i>tropica</i> 383
 <i>truncipennis</i> 343
 <i>tullia</i> 262
 <i>tulsi</i> 386
 <i>tunicula</i> 252
 <i>tushita</i> 280
 <i>tyche</i> 268
 <i>tytleri</i> 368</p> <p style="text-align: center;">U</p> <p><i>Udaina</i> 232
 <i>Udaspes</i> 372
 <i>Unkana</i> 378
 <i>uposatha</i> 364</p> | <p><i>urios</i> 325
 <i>ustulata</i> 252
 <i>utanus</i> 360</p> <p style="text-align: center;">V</p> <p><i>vaga</i> 246
 <i>vajura</i> 321
 <i>valeria</i> 231
 <i>vallona</i> 296
 <i>Vanessa</i> 273
 <i>vardara</i> 347
 <i>varro</i> 330
 <i>varuna</i> (<i>Papilio</i>) 206
 <i>varuna</i> (<i>Rapala</i>) 350
 <i>vasava</i> (<i>Appias</i>) 225
 <i>vasava</i> (<i>Ctenoptilum</i>) 365
 <i>vasuba</i> 369
 <i>vasudeva</i> 257
 <i>vasuki</i> 274
 <i>venata</i> 229
 <i>verhueli</i> 228
 <i>verma</i> 246
 <i>verna</i> 227
 <i>vesta</i> 304
 <i>veturia</i> 311
 <i>vidura</i> 341
 <i>vikasi</i> 281
 <i>vilma</i> 300
 <i>vindyha</i> 246
 <i>viola</i> 321
 <i>violetta</i> 255
 <i>Virachola</i> 349
 <i>viraja</i> 283
 <i>virescens</i> 220
 <i>visala</i> 249
 <i>vitta</i> 353
 <i>volux</i> 372
 <i>vulcanus</i> 340
 <i>vulgaris</i> 237
 <i>vulso</i> 375</p> <p style="text-align: center;">W</p> <p><i>wallacei</i> 272
 <i>waterstradii</i> 362
 <i>watsoni</i> (<i>Ypthima</i>) 243
 <i>watsoni</i> (<i>Chliaria</i>) 347
 <i>watsoni</i> (<i>Iton</i>) 385
 <i>wedah</i> 265
 <i>woolleyi</i> 368</p> <p style="text-align: center;">X</p> <p><i>xanites</i> 371
 <i>Xanthotaenia</i> 258
 <i>xenocles</i> 219
 <i>xenophon</i> 349, 351
 <i>xiphia</i> 220</p> <p style="text-align: center;">Y</p> <p><i>Yasoda</i> 340
 <i>yerburii</i> 280
 <i>Yoma</i> 274
 <i>yotissa</i> 360
 <i>Ypthima</i> 242</p> |
|--|---|---|

Z

zaida 280
zal 262
zaleucus 207
zalmora 314
zamoia 298
zanella 311
Zarona 311

zephyretta 337
Zela 378
zelleri 387
zelmira 225
Zeltus 348
zema 380
Zemeros 305
zeroa 286
zeugitana 246

zeus 378
Zeuxidia 260
ziclea 381
zistra 328
zitenuis 254
Zizera 319
Zographetus 375
zuehara 246

