

## A SUPPLEMENTARY LIST OF THE BUTTERFLIES OF SIAM.

By E. J. GODFREY, B. Sc., F. E. S.

The following additions to my revised list\* of the butterflies of Siam bring up the total number of species known to occur in the country from 692 to 727.

Several species obtained recently by Dr. A. F. G. Kerr in N. E. Siam have not yet been identified.

For convenience of reference, I have numbered the new records in sequence, numbers in brackets referring to the corresponding ones in my previous list.

## PAPILIONIDAE.

693. *Papilio aidoneus* Doubl.

*Papilio aidoneus* Doubleday, A. M. N. H. xvi, p. 178 (1845); Jordan, in Seitz Macrolep. ix, p. 29, figs. 17b, c (1909).

2 ♂♂ : Pang Tawn, Chiengmai Province, N. Siam, 26. 4. 31.

*Distribution* : Sikkim to the Shan States; Siam; Tonkin; Hainan.

## PIERIDAE.

694. *Delias singhapura singhapura* (Wall.).

*Thyca singhapura* Wallace, Trans. Ent. Soc. ser. 3. vol. iv, p. 353, n. 29, t. 7, fig. 2 (1867).

*Delias singhapura singhapura* Fruhstorfer, in Seitz, Macrolep. ix, p. 124 (1919).

2 ♂♂, 2 ♀♀ : Kaw Chang, Ranawng [Renong], P. Siam, 11.1. 29 (Dr. A. F. G. Kerr).

*Distribution* : Burma; P. Siam; Malay Peninsula.

*D. s. singhapura* is extremely rare; the race *singhapura indistincta* Fruhst., from Borneo, is commoner.

[63c] *Delias aglaia tonkiniana* Fruhst.

*Delias aglaia tonkiniana* Fruhstorfer, in Seitz, Macrolep. ix, p. 133 (1910).

1 ♂ : Sriracha, S. E. Siam, October 1930.

*Distribution* : Tonkin; Siam.

---

\* Journ. Siam. Soc., Nat. Hist. Suppl., Vol. vii, No. 4, pp. 203-397 (1930).

*D. a. thyra* Fruhst. has been found in N. and S. E. Siam and *D. a. parthenope* Wall. in P. Siam.

**695. Aporia agathon agathon (Gray).**

*Pieris agathon* Gray, Zool. Misc., p. 33 (1832).

*Aporia agathon agathon* Fruhstorfer, in Seitz, Macrolep. ix, p. 139 (1910); Evans, Ident. Ind. Butt. p. 68 (1932).

1 ♂ : Chiet, Kemarat, N. E. Siam. 21. 5. 32. (*Dr. A. F. G. Kerr*).

*Distribution* : Nepal to the Shan States; N. Siam.

According to Evans, other good subspecies are *a. phryxe* Boisd. (Kashmir-Mussoorie) and *a. caphusa* Moore (Mussoorie-Kumaon).

DANAIDAE.

**696. Euploea aegyptus Butl.**

*Euploea aegyptus* Butler, Proc. Zool. Soc. Lond. p. 277 (1866); Fruhstorfer, in Seitz, Macrolep ix, p. 248 (1910).

1 ♂ : Kantuli, P. Siam. 12. 10. 31.

Riley\* informs me that there is nothing exactly like this specimen in the British Museum and that it suggests a new subspecies. He says, however, that he would hesitate to describe it without other examples from the same district for comparison.

*Distribution* : The species occurs in Singapore, Borneo, Java, Sumatra, and Nias.

SATYRIDAE.

SATYRINAE.

**697. Erebia annada orixa (Moore).**

*Callerebia orixa* Moore, Proc. Zool. Soc. Lond., p. 555 (1872).

*Callerebia annada orixa* Fruhstorfer, in Seitz, Macrolep. ix, p. 301 (1911).

1 ♂ : Pu Wieng, Kawnken, N. E. Siam, 17. 3. 32. (*Dr. A. F. G. Kerr*).

*Distribution* : Assam to N. Burma; N. Siam.

Other subspecies are found in Sikkim, Nepal, Manipur, and the Chin Hills.

**698. Neorina hilda Westw.**

*Neorina hilda* Westwood, in Dblday., Westw. & Hew., Gen. Di. Lep.

---

\* Riley, *in litt.*, 29. 11. 31.

p. 370, pl. 65, fig. 2 (1851); Fruhstorfer in Seitz, Macrolep. ix, p. 326, fig. 94 d (1911).

1 ♂: Doi Chang (2500ft.), Chiengmai Province, N. Siam, 12. 4. 21.

This specimen should have been recorded in my previous list.  
*Distribution*: Sikkim to Assam; N. Siam.

**699. *Coelites epiminthia epiminthia* Westw.**

*Coelites epiminthia* Westwood (Doubl. & Hew.), Gen. Di. Lep. p. 368, n. 2, note (1851).

*Coelites epiminthia epiminthia* Fruhstorfer, in Seitz, Macrolep. ix, p. 329 (1911).

1 ♂: Doi Sutep (4500 ft), Chiengmai Province, N. Siam, Feb. 1931; 1 ♂: Puket, P. Siam, 8. 3. 29.

*Distribution*: Siam; Perak; Borneo; Sumatra.

Other subspecies are found in Burma and the Celebes.

**[182a] *Ragadia crisilda critolina* Evans.**

*Ragadia crisilda* Fruhstorfer, in Seitz, Macrolep. ix, p. 360 (1911).

*Ragadia crisilda critolina* Evans, Ident. Ind. Butt. p. 124 (1932).

1 ♂: Kapong, Takuapa, P. Siam, 17. 2. 29. (*Dr. A. F. G. Kerr*).

*Distribution*: Dawnas to S. Burma; P. Siam.

Replaced in N. Siam by *R. c. critolaus* de Nicév.

**ELYMNIINAE.**

**[190e] *Elymnias hypermnestra violetta* ♀-f *obfuscata* Ril.**

*Elymnias hypermnestra violetta* ♀-f *obfuscata* Riley, Journ. Siam Soc., Nat. Hist. Suppl., Vol. viii, No. 4, p. 249 (1932).

1 ♀: Sriracha, S. E. Siam, May 1931 (*J. J. McBeth*).

**NYMPHALIDAE.**

**[226a] *Penthema binghami godfreyi* Ril.**

*Penthema binghami* Wood-Mason, Journ. As. Soc. Beng. p. 87 (1881).

*Penthema binghami godfreyi* Riley, Journ. Siam Soc., Nat. Hist. Suppl., Vol. viii, No. 4, p. 250, pl. 22, figs. 9, 10 (1932).

4 ♂ ♂: Upper Me Lem, Me Song forest, Prae, N. Siam, April

1918.

*Distribution*: N. Siam.

Replaced in P. Siam by *P. b. mimetica* Lathy.

700. **Neptis nashona aagaardi** Ril.

*Neptis nashona* Swinhoe, A. M. N. H. (6), xvii, p. 357 (1896).

*Neptis nashona aagaardi* Riley, Journ. Siam Soc., Nat. Hist. Suppl. Vol. viii, No. 4, p. 249, pl. 22, fig. 6 (1932).

*Neptis cartica nashona* Fruhstorfer, in Seitz, Macrolep. ix, p. 613 (1912).

1 ♂: Doi Sutep (4500 ft.), Chiengmai Province, N. Siam, Feb. 1931 (*C. J. Aagaard*).

Distribution: N. Siam.

Fruhstorfer (*l. c.*) treats *nashona* as a subspecies of *cartica* Moore, but Riley regards it as a good species with two subspecies —*nashona* in Assam and *aagaardi* in Siam.

701. **Neptis manasa** Moore

*Neptis manasa* Moore, Cat. Lep. Mus. E. I. C. i, p. 165, pl. 4 a, fig. 2 (1857); Fruhstorfer, in Seitz, Macrolep. ix, p. 610 (1912).

1 ♂: Doi Sutep (4500 ft.), Chiengmai Province, N. Siam, Feb. 1931.

Distribution: Sikkim; S. Shan States; N. Siam.

[281a] **Neptis vikasi omeroda** Moore.

*Neptis vikasi omeroda* Fruhstorfer, in Seitz, Macrolep. ix, p. 612 (1912).

1 ♂: Kao Pawta Luang Keo, Ranawng [Renong], P. Siam, 3. 2. 29 (*Dr. A. F. G. Kerr*).

Distribution: P. Siam; Penang; Sumatra.

Replaced in N., C., and E. Siam by *N. v. sakala* Fruhst.

702. **Neptis anjana hyria** Fruhst.

*Neptis anjana* Moore, Trans. Ent. Soc. Lond. p. 309 (1881).

*Neptis anjana hyria* Fruhstorfer, in Seitz, Macrolep. ix, p. 620 (1912).

1 ♂: Kao Pawta Cheng Dong, Ranawng [Renong], P. Siam, 4. 2. 29 (*Dr. A. F. G. Kerr*).

Distribution: P. Siam; Perak; (?) Sumatra.

Other subspecies are found in Borneo, Java, and Nias.

703. **Pantoporia asura** (Moore).

*Athyma asura* Moore, Cat. Lep. Mus. E. I. C. i, p. 171 (1857);

Fruhstorfer (*Pantoporia*), in Seitz, Macrolep. ix, p. 626 (1912).

1 ♂: Doi Sutep (4500 ft.), Chiengmai Province, N. Siam, Feb. 1931.

*Distribution:* Kulu to Tavoy; N. Siam.

**704. Pantoporia idita (Moore)**

*Athyma idita* Moore, Proc. Zool. Soc. Lond. p. 16, pl. 51, fig. 3, ♂ (1858).

*Pantoporia asura idita* Fruhstorfer, in Seitz, Macrolep. ix, p. 627 (1912); Evans, Ident. Ind. Butt. p. 164 (1932).

1 ♀: Doi Sutep (4500 ft.), Chiengmai Province, N. Siam Feb. 1931.

*Distribution:* S. Burma to Mergui; Siam.

Both Fruhstorfer and Evans treat *idita* as a subspecies of *asura*, but Riley\* regards it as a good species. He writes:—"The obvious difference between *asura* and *idita* is in the discal row of white spots on the upperside hindwing. They are black-centred in *asura* and clear in *idita*..... Series of the two placed side by side are very different."

**705. Liminitis dudu Westw.**

*Liminitis dudu* Westwood, in Dbdlay., Westw. & Hew., Gen. Di. Lep. ii, p. 276 (1850); Fruhstorfer, in Seitz, Macrolep. ix, p. 638 (1912).

1 ♂: Doi Sutep (4500 ft.), Chiengmai Province, N. Siam, Feb. 1931.

*Distribution:* Sikkim to Burma; Siam; Sumatra; Formosa.

**706. Eriboea hebe chersonesus Fruhst.**

*Charaxes hebe* Butler, Proc. Zool. Soc. Lond. p. 634, n. 46, t. 37, fig. 3 (1865).

*Eriboea hebe chersonesus* Fruhstorfer, in Seitz Macrolep. ix, p. 721 (1914).

1 ♂: Kao Kalakiri, Pattani, P. Siam, 29. 3. 28 (*Dr. A. F. G. Kerr*).

*Distribution:* P. Siam; Victoria Point; Malacca; Penang.

---

\* Riley, in *litt.*, 17. 12. 32.

**707. *Eriboea schreiberi assamensis* Roths. & Jord.**

*Nymphalis schreiberi* Godart, Encycl. Méth. ix, Suppl. p. 825 (1823).

*Eriboea schreiberi assamensis* Fruhstorfer, in Seitz, Macrolep. ix, p. 725 (1914).

1 ♂ : Tung Luang, Petchaburi, W. Siam, 8. 11. 31. (*Dr. A. F. G. Kerr*).

*Distribution* : The species ranges from India to the Philippines, south the Malay Peninsula, Sumatra, Borneo, and Java.

*E. s. assamensis* is represented in the British Museum only by the male recorded above, and one male from the Malay Peninsula.

**ERYCINIDAE.****708. *Dodona dipoea dipoea* Hew.**

*Dodona dipoea* Hewitson, Ex. Butt. iii, pl. 1, fig. 3 (1866).

*Dodona dipoea dipoea* Fruhstorfer, in Seitz, Macrolep. ix, p. 775, pl. 139 d (1914).

1 ♂, 1 ♀ : Doi Angka (4500 ft.), Chiengmai Province, N. Siam, Feb. 1932.

*Distribution* : Sikkim to Assam ; N. Siam.

Other subspecies are found in Mussoorie and N. Burma.

**LYCAENIDAE.****709. *Logania obscura* (Röb.).**

*Allotinus obscura* Röber, Iris, p. 532 (1885) ; Fruhstorfer (*Logania*), in Seitz, Macrolep. ix, p. 807 (1915).

1 ♂ : Tambon Kao Panom (c. 400 ft.), Krabi, P. Siam, 20. 3. 30 (*Dr. A. F. G. Kerr*).

*Distribution* : Tombugu, E. Celebes (*Fruhstorfer*).

This is an extremely rare species.

**710. *Hypochrysops kerri* Ril.**

*Hypochrysops kerri* Riley, Journ. Siam Soc., Nat. Hist. Suppl., Vol. viii, No. 4, p. 252, pl. 22, figs. 3, 4 (1932).

1 ♂ : Tambon Kao Panom (c. 400 ft.), Krabi, P. Siam, 20. 3. 30 (*Dr. A. F. G. Kerr*).

*Distribution* : P. Siam.

Riley (*l.c.*) remarks that the genus *Hypochrysops* is essentially Australian and Papuan and that, apart from *kerri*, only one species, *coelisparus*, which occurs in Nias has been found west of the Moluccas.

**711. *Talicada macbethi* Ril.**

*Talicada macbethi* Riley, Journ. Siam Soc., Nat. Hist. Suppl., Vol. viii, No. 4, p. 251, pl. 22, figs. 7, 8 (1932).

1 ♀: Lat Bua Khao, on the Korat line E. Siam 7. 2. 29 (*J. J. McBeth*).

*Distribution*: Siam.

**712. *Everes potanini* (Alph.).**

*Lycaena potanini* Alphéraky, Rom. Mém. v, p. 104, pl. 5, fig. 4 (1889); Seitz (*Everes*), Macrolep. ix, p. 924 (1924).

1 ♂: Kanburi, C. Siam, 13. 9. 31.

*Distribution*: N. Burma to the Dawnas; Siam; W. China.

**713. *Jamides philatus subdita* (Moore).**

*Lampides subdita* Moore, Journ. Linn. Soc. Zool. xxi, p. 41 (1886).

*Jamides philatus subdita* Seitz, Macrolep. ix, p. 910 (1924).

1 ♂: La-un, Ranawng [Renong], P. Siam, 1.1.29 (*Dr. A. F. G. Kerr*).

*Distribution*: S. Burma; P. Siam.

The species ranges from Burma west to the Philippines, south to New Guinea.

**714. *Heliophorus androcles rubida* Ril.**

*Ilerda androcles* Doubleday and Hewitson, Gen. Di. Lep. p 487 (1851); Seitz, Macrolep. ix, p. 933 (1927).

*Heliophorus androcles rubida* Riley, Journ. Bomb. Nat. Hist. Soc. xxxiii, p. 396 (1929).

1 ♂: Doi Sutep (5500 ft.), Chiengmai Province, N. Siam, Feb. 1928 (*Prof. D. T. A. Cockerell*)

*Distribution*: Upper Burma; C. Yunnan; N. Siam.

In addition to *a. rubida* Riley recognises four other races of *androcles*—*a. corruscans* Moore (Mussoorie), *a. moorei* Hew. (Sikkim), *a. tytleri* Ril. (Naga Hills), and *a. androcles* (Sylhet).

**715. *Amblypodia horsfieldi eurysthenes* Fruhst.**

*Amblypodia horsfieldi* Pagst, Jahrb. Nassau. Ver. 43, p. 99

*Amblypodia horsfieldi eurythenes* Seitz, Macrolep. ix, p. 959 (1926).

1 ♂ : Kapong, Takuapa, P. Siam, 17. 2. 29 (Dr. A. F. G. Kerr).

*Distribution* : Dawnas to S. Burma ; P. Siam.

Other subspecies are found in Malacca, Sumatra, Borneo, and Java.

[487a] ***Amblypodia perimuta regia* Evans.**

*Amblypodia perimuta* Moore, Cat. Lep. Mus. E. I. C. p. 42 (1857) ;

Seitz, Macrolep. ix, p. 965 (1926).

*Amblypodia perimuta regia* Evans, Journ. Bomb. N. H. S. xxx, p. 635 (1925).

1 ♂ : Kampuan, Ranawng [Renong], P. Siam, 5. 2. 29 (Dr. A. F. G. Kerr).

*Distribution* : S. Burma, Mergui, P. Siam.

Replaced in N. Siam by *A. p. perimuta* Moore.

**716. *Amblypodia anniella* Hew.**

*Amblypodia anniella* Hewitson, Cat. Lyc. Br. Mus., p. 10 (1862) ; Seitz, Macrolep. ix, p. 961, fig. 150 d (1926).

1 ♂ : Kao Pawta Luang Keo (c. 1300m.), Ranawng [Renong], P. Siam, 2. 2. 29 (Dr. A. F. G. Kerr).

*Distribution* : Shan States to S. Burma ; P. Siam ; Philippine Islands, Moluccas, Borneo ; Java.

**717 *Amblypodia adriana* de Nicév.**

*Amblypodia adriana* de Nicéville, Journ. As. Soc. Bengal, 52 (2),

p. 79 ; Seitz, Macrolep. ix, p. 961, fig. 150B, b (1926) ; Evans, Ident.

Ind. Butt., p. 271 (1932).

1 ♂ : Tung Luang, Petchaburi, C. Siam, 8. 11. 31 (Dr. A. F. G. Kerr).

*Distribution* : Sikkim to the Shan States (Evans).

Riley\* informs me that, apart from the specimen recorded above, the British Museum has no examples of *adriana* from anywhere south of Assam.

**718. *Pratapa deva* (Moore).**

*Camena deva* Moore, Cat. Lep. Mus. E. I. C. p. 46 (1857) ; Seitz,

---

\* Riley, *in litt.*, 31. 12. 31.

*Macrolep.* ix, p. 971 (1926).

1 ♂ : Naung Bua Deng, Chaiyapum, E. Siam, 2. 3. 31 (*Dr. A. F. G. Kerr*)

*Distribution* : N. India to Siam, Borneo and Java.

**719. *Sithon nedymond nedymond* (Cram.).**

*Papilio nedymond* Cramer, Pap. Exot. iv, t. 299, E, F (1782).

*Sithon nedymond nedymond* Seitz, *Macrolep.* ix, p. 978 (1926).

1 ♂, 1 ♀ : Klawng Kampuam, Ranawng [Renong], P. Siam, 4. 2. 29 (*Dr. A. F. G. Kerr*).

*Distribution* : P. Siam ; Java.

Other subspecies are found in Malacca and Sumatra.

**720. *Tajuria mantra mantra* (Feld.).**

*Pseudolycaena* Felder, Wien. Ent. Mon. iv, p. 396, n. 9 (1860) ; Seitz (*Tajuria*), *Macrolep.* ix, p. 974 (1926).

1 ♀ : Kaw Libong, Trang, P. Siam, 26. 4. 31 (*Dr. A. F. G. Kerr*).

*Distribution* : S. Burma, P. Siam ; Malay Peninsula ; Malacca ; Sumatra.

Other subspecies are found in Borneo, Java and the Celebes.

**721. *Tajuria isaeus isaeus* (Hew.).**

*Myrina isaeus* Hewitson, Ill. Di. Lep. p. 44 (1865).

*Tajuria isaeus isaeus* Seitz, *Macrolep.* ix, p. 975, fig. 156 c (1926).

1 ♀ : Nang Yawn, Takuapa, P. Siam, 8. 2. 29 (*Dr. A. F. G. Kerr*).

*Distribution* : S. Burma; P. Siam; Borneo.

Other subspecies are found in Java and Nias.

**[514] *Neocheritra amrita paulina* Ril.**

*Myrina amrita* Felder, Wien. Ent. Monat. iv, p. 395 (1860) ; Seitz. (*Neocheritra*), *Macrolep.* ix, p. 991, fig. 259 a (1926).

*Neocheritra amrita paulina* Riley, Journ. Siam Soc., Nat. Hist. Suppl., Vol. viii, No. 4, p. 252, pl. 22, fig. 5 (1932)

1 ♀ : Kaw Payam, Ranawng [Renong], P. Siam, 14. 1. 29.

*Distribution* : P. Siam.

Typical *amrita* occurs in Malacca and Sumatra, and a subspecies, *theodora* Druce, is found in Borneo.

**[519a] *Cheritra freja regia* Evans.**

*Cheritra freja* Seitz, *Macrolep.* ix, p. 993 (1926).

*Cheritra freja regia* Evans, Journ. Bomb. N. H. S. xxx, p. 766 (1925).

1 ♀ : Kapôr, Ranawng [Renong], P. Siam, 17. 1. 29.

*Distribution* : Mergui, P. Siam.

Replaced elsewhere in Siam by *C. f. freja* Fabr.

**722. *Artipe eryx* (Linn.).**

*Deudorix eryx* Seitz, Macrolep. ix, p. 1001, fig. 146 f (as *anna*) (1926), Evans (*Artipe*), Ident. Ind. Butt., p. 294 (1932).

1 ♀ : Kao Libong, Trang, P. Siam, 29. 4. 30.

*Distribution* : From the Himalayas east over Hong Kong and S. China to Formosa, south to Assam and Siam.

**723. *Virachola isocrates* (Fabr.).**

*Hesperia isocrates* Fabricius, Ent. Syst. iii, 1, p. 266 (1793); Seitz (*Deudorix*), Macrolep. ix, p. 1001, fig. 161 b (1926).

3 ♂ ♂ : Tung Luang, Petchaburi, C. Siam, 8. 11. 31.

*Distribution* : Hitherto known only from the region extending from N. W. India to Ceylon and across to Orissa.

The pupae of the specimens recorded above were found by Dr. A. Kerr in the fruit of *Diospyros* sp.

Riley (*in litt.*, 21. 1. 32), remarking on the unexpected occurrence of *isocrates* in Siam, assumed that it could only have been imported somehow with plants. Dr. Kerr, however, says that he found *Diospyros* sp. growing wild near Tung Luang and other remote villages in the Petchaburi district and that neither it, nor other plants found thereabouts, could possibly have been imported.

### HESPERIIDAE.

**724. *Tagiades lavata* Butt. (= *albovittata* Moore).**

*Tagiades lavata* Butler, Trans. Linn. Soc. Lond. Zool., p. 557, pl. 69, fig. 8 (1877); Seitz, Macrolep. ix, p. 1042, fig. 165 d (1927).

*Tagiades alborivitata* Moore, Proc. Zool. Soc. Lond., p. 534 (1883).

1 ♂ : Kraburi, Ranawng [Renong], P. Siam, 23. 12. 28. (Dr. A. F. G. Kerr).

*Distribution* : S. Burma to the Malay Peninsula; Borneo; Sumatra; Natuna Isles.

The British Museum possesses only the male recorded above

and five other examples of this species.

**725. *Gangara sybrita* (Hew.).**

*Hidari sybrita* Hewitson, Ann. and Mag. Nat. Hist. ser. 4, vol. xviii, p. 451 (1876); Seitz, Macrolep. ix, p. 1071, fig. 170 b (1927).

1 ♀: Kaw Libong, Trang, P. Siam, 26. 4. 30. (*Dr. A. F. G. Kerr*).

*Distribution*: P. Siam; Malacca; (?) Borneo.

This is a very rare species; the British Museum possesses only the female recorded above and two males.

**726. *Sepa bononia* (Hew.).**

*Hesperia bononia* Hewitson, Deser. Hesp. p. 29. n. 16 (1868); Seitz (*Scobura*) Macrolep. ix, p. 1069, fig. 169 h (1927).

1 ♀: Lam Rieng, Ranawng [Renong], 27. 12. 28. (*Dr. A. F. G. Kerr*).

*Distribution*: P. Siam; Malacca; Singapore.

**727. *Baoris assamensis* (W. M. & de Nic.).**

*Parnara assamensis* Wood Mason and de Nicéville, Journ. As. Soc. Beng., p. 65 (1882); Seitz, Macrolep. ix, p. 1083 (1927); Evans (*Baoris*), Ident. Ind. Butt. p. 416 (1932).

1 ♂: Doi Angka (4500 ft.), Chiengmai Province, N. Siam,

*Distribution*: Central Provinces; Mussoorie to Burma; Siam; Malay Peninsula.

Seitz treats *assamensis* as a subspecies of *P. guttatus* Brem. Evans regards it as a separate species.

